

Europe Supplementary Material

Coordinating Lead Authors: Birgit Bednar-Friedl (Austria), Robbert Biesbroek (the Netherlands), Daniela N. Schmidt (UK/Germany)

Lead Authors: Peter Alexander (UK), Knut Yngve Børsheim (Norway), Jofre Carnicer (Spain), Elena Georgopoulou (Greece), Marjolijn Haasnoot (the Netherlands), Gonéri Le Cozannet (France), Piero Lionello (Italy), Oksana Lipka (Russian Federation), Christian Möllmann (Germany), Veruska Muccione (Switzerland/Italy), Tero Mustonen (Finland), Dieter Piepenburg (Germany), Lorraine Whitmarsh (UK)

Contributing Authors: Magnus Benzie (Sweden), Pam Berry (UK), Sara Burbi (UK), Erika Coppola (Italy), Mladen Domazet (Croatia), Frank Ewert (Germany), Federica Gasbarro (Italy), Matthias Gault (Italy), François Gemenne (Belgium), Peter Greve (Austria/Germany), Ana Iglesias (Spain), Elizabeth Kendon (UK), Heidi Kreibich (Germany), Nikos Koutsias (Greece), Anna Laine-Petäjäkangas (Finland), Dimitris Lalas (Greece), Cristina Linares Gil (Spain), Danijela Markovic (Germany), Sadie McEvoy (the Netherlands/Ireland), Ana Mijic (UK), Raya Muttarak (Austria/Thailand), Rita Nogherotto (Italy), Hans Orru (Estonia), Mark Parrington (UK), Jeff Price (UK), Kaisa Raitio (Sweden), Marta Guadalupe Rivera Ferre (Spain), Jan C. Semenza (Switzerland), Rubén Valbuena (UK), Michelle van Vliet (the Netherlands), Heidi Webber (Germany), Laura Wendling (Finland), Katherine Yates (UK), Monika Zurek (UK)

Chapter Scientists: Sadie McEvoy (the Netherlands/Ireland), Phoebe O'Brien (UK/Sweden)

Review Editors: Georg Kaser (Austria/Italy), Jose Manuel Moreno (Spain)

This Supplementary Material should be cited as:

Bednar-Friedl, B., R. Biesbroek, D.N. Schmidt, P. Alexander, K.Y. Børsheim, J. Carnicer, E. Georgopoulou, M. Haasnoot, G. Le Cozannet, P. Lionello, O. Lipka, C. Möllmann, V. Muccione, T. Mustonen, D. Piepenburg, and L. Whitmarsh, 2022: Europe Supplementary Material. In: *Climate Change 2022: Impacts, Adaptation and Vulnerability*. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Available from <https://www.ipcc.ch/report/ar6/wg2/>.

Table of Contents

SM13.1 Supplementary Material Supporting
Section 13.2 3

SM13.2 Supplementary Material Supporting
Section 13.3 8

SM13.3 Supplementary Material Supporting
Section 13.4 10

SM13.4 Supplementary Material Supporting
Section 13.5 11

SM13.5 Supplementary Material Supporting
Section 13.6 14

SM13.6 Supplementary Material Supporting
Section 13.7 32

SM13.7 Supplementary Material Supporting
Section 13.8 33

SM13.8 Supplementary Material Supporting
Section 13.10 34

SM13.9 Supplementary Material Supporting the
Feasibility and Effectiveness Assessment 36

SM13.10 Supplementary Material Supporting the Key
Risks and Burning Embers Assessments 36

SM13.11 Supplementary Material Supporting the
Development of Adaptation Pathways 48

References 49

SM13.1 Supplementary Material Supporting Section 13.2

Table SM13.1 | Literature sources used in the assessment of feasibility and effectiveness of adaptation options for water systems in Europe (Figure 13.6)

Impact type	Adaptation option	Effectiveness					Feasibility				Geophysical
							Economic	Technological	Institutional	Sociocultural	Ecological
Coastal and riverine flooding	Flood defences (Protect)	Andersson-Sköld et al. (2015); Alfieri et al. (2016a); Bollinger and Dijkema (2016); Bouwer et al. (2018); Pérez-Morales et al. (2018); Thacker et al. (2018); EEA (2019a); Straatsma et al. (2019); Dottori et al. (2020); Voudoukas et al. (2020); Unglösser et al. (2021)	Alfieri et al. (2016a); Bollinger and Dijkema (2016); Bouwer et al. (2018); Pérez-Morales et al. (2018); Thacker et al. (2018); EEA (2019a); Voudoukas et al. (2020)	Metin et al. (2018); Thacker et al. (2018); Straatsma et al. (2019); Voudoukas et al. (2020)	Bollinger and Dijkema (2016); Bubeck et al. (2017); Thacker et al. (2018); EEA (2019a)	Alfieri et al. (2016a)	Andersson-Sköld et al. (2015); Del Bello (2018); Pérez-Morales et al. (2018); Voudoukas et al. (2020)	Del Bello (2018); EEA (2019a); Straatsma et al. (2019)			
	Flood preparedness and early warning plans (Protect/ Accommodate)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)	Poussin et al. (2013); Kreibich et al. (2015); Pappenberger et al. (2015); Bubeck et al. (2017); Kreibich et al. (2017); Pérez-Morales et al. (2018); Restemeyer et al. (2018); Varrani and Nones (2018); Adedeji et al. (2019); Merz et al. (2020); Ribas et al. (2020); Kreibich et al. (2021)
	Planned relocation (Retreat)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)	Koerth et al. (2013); Harman et al. (2015); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede (2019); Rey-Valette et al. (2019); Buser (2020); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Seebauer and Winkler (2020); Thaler (2021)

		Feasibility						Geophysical
Impact type	Adaptation option	Effectiveness	Economic	Technological	Institutional	Sociocultural	Ecological	
Coastal and riverine flooding	No-build zone, restrict new developments (Avoidance)	Koerth et al. (2013); Andersson-Sköld et al. (2015); Harman et al. (2015); Thieken et al. (2016); Pérez-Morales et al. (2018); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede et al. (2019); Rey-Valette et al. (2019); Dottori et al. (2020); Lincke et al. (2020); Thaler Mayr et al. (2020); Thaler (2021)	Koerth et al. (2013); Andersson-Sköld et al. (2015); Harman et al. (2015); Thieken et al. (2016); Pérez-Morales et al. (2018); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede et al. (2019); Rey-Valette et al. (2019); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Thaler (2021)	Koerth et al. (2013); Andersson-Sköld et al. (2015); Harman et al. (2015); Thieken et al. (2016); Pérez-Morales et al. (2018); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede et al. (2019); Rey-Valette et al. (2019); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Thaler (2021)	Koerth et al. (2013); Andersson-Sköld et al. (2015); Harman et al. (2015); Thieken et al. (2016); Pérez-Morales et al. (2018); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede et al. (2019); Rey-Valette et al. (2019); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Thaler (2021)	Koerth et al. (2013); Poussin et al. (2013); Keskitalo et al. (2014); Paudel et al. (2015); Penning-Rowse and Priest (2015); Surminski et al. (2015); Hudson et al. (2016); O'Hare et al. (2016); Suykens et al. (2016); Surminski and Thieken (2017); Hanger et al. (2018); Surminski (2018)	Koerth et al. (2013); Poussin et al. (2013); Keskitalo et al. (2014); Paudel et al. (2015); Penning-Rowse and Priest (2015); Surminski et al. (2015); Hudson et al. (2016); O'Hare et al. (2016); Suykens et al. (2016); Surminski and Thieken (2017); Hanger et al. (2018); Surminski (2018)	Koerth et al. (2013); Andersson-Sköld et al. (2015); Harman et al. (2015); Thieken et al. (2016); Pérez-Morales et al. (2018); Thacker et al. (2018); Dachary-Bernard et al. (2019); Hofstede et al. (2019); Rey-Valette et al. (2019); Dottori et al. (2020); Lincke et al. (2020); Mayr et al. (2020); Thaler (2021)
		Koerth et al. (2013); Poussin et al. (2013); Keskitalo et al. (2014); Paudel et al. (2015); Penning-Rowse and Priest (2015); Surminski et al. (2015); Hudson et al. (2016); O'Hare et al. (2016); Suykens et al. (2016); Surminski and Thieken (2017); Hanger et al. (2018); Surminski (2018)	Koerth et al. (2013); Poussin et al. (2013); Keskitalo et al. (2014); Paudel et al. (2015); Penning-Rowse and Priest (2015); Surminski et al. (2015); Hudson et al. (2016); O'Hare et al. (2016); Suykens et al. (2016); Surminski and Thieken (2017); Hanger et al. (2018); Surminski (2018)	Koerth et al. (2013); Poussin et al. (2013); Keskitalo et al. (2014); Paudel et al. (2015); Penning-Rowse and Priest (2015); Surminski et al. (2015); Hudson et al. (2016); O'Hare et al. (2016); Suykens et al. (2016); Surminski and Thieken (2017); Hanger et al. (2018); Surminski (2018)	Koerth et al. (2013); Poussin et al. (2013); Keskitalo et al. (2014); Paudel et al. (2015); Penning-Rowse and Priest (2015); Surminski et al. (2015); Hudson et al. (2016); O'Hare et al. (2016); Suykens et al. (2016); Surminski and Thieken (2017); Hanger et al. (2018); Surminski (2018)	Koerth et al. (2013); Poussin et al. (2013); Keskitalo et al. (2014); Paudel et al. (2015); Penning-Rowse and Priest (2015); Surminski et al. (2015); Hudson et al. (2016); O'Hare et al. (2016); Suykens et al. (2016); Surminski and Thieken (2017); Hanger et al. (2018); Surminski (2018)	Koerth et al. (2013); Poussin et al. (2013); Keskitalo et al. (2014); Paudel et al. (2015); Penning-Rowse and Priest (2015); Surminski et al. (2015); Hudson et al. (2016); O'Hare et al. (2016); Suykens et al. (2016); Surminski and Thieken (2017); Hanger et al. (2018); Surminski (2018)	
		Flood insurance (Supporting)						
Coastal flooding	Wet and dry proofing (Accommodate)	Botzen et al. (2013); Poussin et al. (2015); Stojanov et al. (2015); Bouwer et al. (2018); Hudson et al. (2019)	Botzen et al. (2013); Poussin et al. (2015); Stojanov et al. (2015); Bouwer et al. (2018); Hudson et al. (2019)	Botzen et al. (2013); Poussin et al. (2015); Stojanov et al. (2015); Bouwer et al. (2018); Hudson et al. (2019)	Botzen et al. (2013); Poussin et al. (2015); Stojanov et al. (2015); Bouwer et al. (2018); Hudson et al. (2019)	Botzen et al. (2013); Stojanov et al. (2015)		
		Campos et al. (2016)	Temmerman et al. (2013); Campos et al. (2016); de Schipper et al. (2021); Staudt et al. (2021)	Temmerman et al. (2013); Campos et al. (2016); de Schipper et al. (2021); Staudt et al. (2021)	Campos et al. (2016); de Schipper et al. (2021); Staudt et al. (2021)	Temmerman et al. (2013); de Schipper et al. (2021); Staudt et al. (2021)	Temmerman et al. (2013); Parkinson and Ogurcak (2018); de Schipper et al. (2021); Staudt et al. (2021)	Parkinson and Ogurcak (2018); Staudt et al. (2021)
		Temmerman et al. (2013); Narayan et al. (2016); Vuik et al. (2019)	Temmerman et al. (2013); Narayan et al. (2016); Vuik et al. (2019)	Temmerman et al. (2013); Narayan et al. (2016); Vuik et al. (2019)	Temmerman et al. (2013); Narayan et al. (2016); Vuik et al. (2019)	Temmerman et al. (2013); Narayan et al. (2016); Vuik et al. (2019)	Temmerman et al. (2013); Narayan et al. (2016); Vuik et al. (2019)	Temmerman et al. (2013); Vuik et al. (2019)

		Effectiveness		Feasibility						
Impact type	Adaptation option			Economic	Technological	Institutional	Sociocultural	Ecological	Geophysical	
Riverine flooding	Wet and dry proofing (Accommodate)	Hudson et al. (2014); Kreibich et al. (2015); Poussin et al. (2015); Jones et al. (2017); Osberghaus (2018); Salinas-Rodriguez et al. (2018); Thacker et al. (2018); Haer et al. (2019); Hudson et al. (2019); Sairam et al. (2019)	Osberghaus (2017); Thacker et al. (2018)		Botzen et al. (2013); Hudson et al. (2014); Kreibich et al. (2015); Poussin et al. (2015); Stojanov et al. (2015); Jones et al. (2017); Osberghaus (2017); Bouwer et al. (2018); Salinas-Rodriguez et al. (2018); Thacker et al. (2018); Haer et al. (2019); Hudson et al. (2019); Sairam et al. (2019)	Botzen et al. (2013); Hudson et al. (2014); Kreibich et al. (2015); Poussin et al. (2015); Stojanov et al. (2015); Jones et al. (2017); Osberghaus (2017)			Jones et al. (2017)	
	Ecosystem based (e.g., floodplain restoration, widening riverbed) (Protect)	Dadson et al. (2017); Straatsma et al. (2019); Dottori et al. (2020)	Straatsma et al. (2019); Dottori et al. (2020)		Straatsma et al. (2019)		Straatsma et al. (2019); Dottori et al. (2020)		Asselman and Klijn (2016); Dadson et al. (2017); Straatsma et al. (2019); Dottori et al. (2020); European Commission (2020)	
	Retention and diversion (Accommodate)	Gocht and Meon (2016); Dadson et al. (2017); Verkerk et al. (2017); Dottori et al. (2020)	Gocht and Meon (2016); Verkerk et al. (2017)		Gocht and Meon (2016); Verkerk et al. (2017)			Verkerk et al. (2017); Dottori et al. (2020)		
Pluvial flooding	Green roofs (Accommodate)	Andersson-Sköld et al. (2015); Zölch et al. (2017); Liu et al. (2018); Babovic and Mijic (2019)	Andersson-Sköld et al. (2015); Zölch et al. (2017); Liu et al. (2018); Babovic and Mijic (2019)		European Commission (2020)		Zölch et al. (2017)	Andersson-Sköld et al. (2015); Zölch et al. (2017)	Andersson-Sköld et al. (2015); Zölch et al. (2017); Liu et al. (2018); European Commission (2020)	
	Retention parks (Accommodate)	Andersson-Sköld et al. (2015); Ambjerg-Nielsen et al. (2015); Maragno et al. (2018); Salinas-Rodriguez et al. (2018); Ribas et al. (2020)			Maragno et al. (2018); Ribas et al. (2020)		Andersson-Sköld et al. (2015); Maragno et al. (2018)	Andersson-Sköld et al. (2015); Ambjerg-Nielsen et al. (2015); Maragno et al. (2018); Salinas-Rodriguez et al. (2018); Ribas et al. (2020)	Andersson-Sköld et al. (2015); Ambjerg-Nielsen et al. (2015); Maragno et al. (2015); Maragno et al. (2018); Salinas-Rodriguez et al. (2018); Ribas et al. (2020)	
	Update drainage system and pumps (Accommodate)	Skougaard Kaspersen et al. (2017)	Skougaard Kaspersen et al. (2017); Liu and Jensen (2018); EEA (2020b); Ribas et al. (2020)		Ribas et al. (2020)	Liu and Jensen (2018); EEA (2020b)	EEA (2020b)	EEA (2020b)	Liu and Jensen (2018); EEA (2020b)	

Effectiveness		Feasibility					Geophysical	
Adaptation option	Impact type	Economic	Technological	Institutional	Sociocultural	Ecological		
Water scarcity	Supply: storage (i.e., reservoirs)	Papadaskalopoulou et al. (2015b); Kingsborough et al. (2016); Varela-Ortega et al. (2016); Bucak et al. (2017); Verkerk et al. (2017); Di Baldassarre et al. (2018); Garnier and Holman (2019)	Papadaskalopoulou et al. (2015b); Varela-Ortega et al. (2016); Verkerk et al. (2017); Garnier and Holman (2019)	Papadaskalopoulou et al. (2015b); Kingsborough et al. (2016); Varela-Ortega et al. (2018); Garnier and Holman (2019)	Papadaskalopoulou et al. (2015b); Verkerk et al. (2017)	Papadaskalopoulou et al. (2015b); Varela-Ortega et al. (2016); Di Baldassarre et al. (2018); Santos et al. (2018); Garnier and Holman (2019)	Papadaskalopoulou et al. (2015b); Kingsborough et al. (2016); Bucak et al. (2017); Di Baldassarre et al. (2018); Santos et al. (2018); Garnier and Holman (2019)	
	Supply: water diversion and transfer	Flekens et al. (2013); Collet et al. (2015); Papadaskalopoulou et al. (2015b)	Flekens et al. (2013); Collet et al. (2015); Papadaskalopoulou et al. (2015b)	Flekens et al. (2013); Collet et al. (2015); Papadaskalopoulou et al. (2015b); Garnier and Holman (2019)	Flekens et al. (2013); Papadaskalopoulou et al. (2015b)	Papadaskalopoulou et al. (2015b); Garnier and Holman (2019)	Collet et al. (2015)	
	Supply: desalination	Papadaskalopoulou et al. (2015b); Kingsborough et al. (2016); Garnier and Holman (2019); Morote et al. (2019)	Papadaskalopoulou et al. (2015b); Kingsborough et al. (2016); Garnier and Holman (2019); Morote et al. (2019)	Papadaskalopoulou et al. (2015b); Kingsborough et al. (2016); Garnier and Holman (2019); Morote et al. (2019)	Morote et al. (2019)	Papadaskalopoulou et al. (2015b); Morote et al. (2019); Papadaskalopoulou et al. (2015)	Garnier and Holman (2019); Morote et al. (2019)	
	Supply: water reuse	Papadaskalopoulou et al. (2015b); Kingsborough et al. (2016); Morote et al. (2019); De Roo et al. (2020)	Papadaskalopoulou et al. (2015b); Morote et al. (2019)	Papadaskalopoulou et al. (2015b); Kingsborough et al. (2016); Morote et al. (2019); De Roo et al. (2020)	Papadaskalopoulou et al. (2015b); Morote et al. (2019)	Papadaskalopoulou et al. (2015b); Morote et al. (2019)	Morote et al. (2019)	
	Demand: water saving and efficiency		Papadaskalopoulou et al. (2015); Kingsborough et al. (2016); Varela-Ortega et al. (2016); Iglesias et al. (2018); Manouseli et al. (2018); Papadimitriou et al. (2019); De Roo et al. (2020)	Collet et al. (2015); Papadaskalopoulou et al. (2015b); van Duinen et al. (2015); Fader et al. (2016); Kingsborough et al. (2016); Varela-Ortega et al. (2016); Rey et al. (2017); Verkerk et al. (2017); Iglesias et al. (2018); Manouseli et al. (2018)	Collet et al. (2015); Papadaskalopoulou et al. (2015b); Varela-Ortega et al. (2016); Verkerk et al. (2017); Papadimitriou et al. (2019)	Collet et al. (2015); Papadaskalopoulou et al. (2015b); Manouseli et al. (2018)	van Duinen et al. (2015); Fader et al. (2016); Varela-Ortega et al. (2016); Verkerk et al. (2017); Iglesias et al. (2018); Papadimitriou et al. (2019)	van Duinen et al. (2015); Papadimitriou et al. (2019)
			Papadaskalopoulou et al. (2015b); Manouseli et al. (2018); Garnier and Holman (2019); Teotónio et al. (2020)	Manouseli et al. (2018); Garnier and Holman (2019)	Papadaskalopoulou et al. (2015b); Manouseli et al. (2018); Garnier and Holman (2019); Teotónio et al. (2020)	Papadaskalopoulou et al. (2015b); Manouseli et al. (2018); Garnier and Holman (2019); Teotónio et al. (2020)	Papadaskalopoulou et al. (2015b); Manouseli et al. (2018); Garnier and Holman (2019); Teotónio et al. (2020)	Manouseli et al. (2018); Garnier and Holman (2019); Teotónio et al. (2020)

Impact type	Adaptation option	Effectiveness	Feasibility					Geophysical
			Economic	Technological	Institutional	Sociocultural	Ecological	
Water scarcity	Demand: economic instruments	Kayaga and Smout (2014); Wimmer et al. (2014); Esteve et al. (2015); Kahil et al. (2015); Papadaskalopoulou et al. (2015b); Varela-Ortega et al. (2016); Koopman et al. (2017); Crespo et al. (2019)	Kayaga and Smout (2014); Esteve et al. (2015); Kahil et al. (2015); Papadaskalopoulou et al. (2015b); Varela-Ortega et al. (2016); Rey et al. (2017); Garnier and Holman (2019)	Wimmer et al. (2014); Esteve et al. (2015); Kahil et al. (2015); Papadaskalopoulou et al. (2015b); Varela-Ortega et al. (2016); Koopman et al. (2017); Crespo et al. (2019)	Esteve et al. (2015); Kahil et al. (2015); Papadaskalopoulou et al. (2015b); Varela-Ortega et al. (2016); Koopman et al. (2017); Crespo et al. (2019)	Kayaga and Smout (2014); Esteve et al. (2015); Kahil et al. (2015); Papadaskalopoulou et al. (2015b)	Esteve et al. (2015); Kahil et al. (2015); Varela-Ortega et al. (2016)	Esteve et al. (2015)
	Demand: land management and cover change	Papadaskalopoulou et al. (2015b); Carvalho-Santos et al. (2016); Varela-Ortega et al. (2016); Papadimitriou et al. (2019)	Carvalho-Santos et al. (2016); Varela-Ortega et al. (2016)	Carvalho-Santos et al. (2016); Varela-Ortega et al. (2016); Verkerk et al. (2017)	Varela-Ortega et al. (2016); Verkerk et al. (2017)	Garnier and Holman (2019)	Carvalho-Santos et al. (2016); Varela-Ortega et al. (2016); Verkerk et al. (2017); Papadimitriou et al. (2019)	Carvalho-Santos et al. (2016); Varela-Ortega et al. (2016); Verkerk et al. (2017); Garnier and Holman (2019); Papadimitriou et al. (2019)
	Monitoring and operational management, drought early warning systems	Papadaskalopoulou et al. (2015b)	Papadaskalopoulou et al. (2015b); Verkerk et al. (2017); Garnier and Holman (2019)	Papadaskalopoulou et al. (2015b)	Papadaskalopoulou et al. (2015b); Verkerk et al. (2017)	Verkerk et al. (2017)	Verkerk et al. (2017)	

SM13.2 Supplementary Material Supporting Section 13.3

Table SM13.2 | Literature sources used for assessment of major impacts on, and risks for, terrestrial and freshwater ecosystems in Europe for 1.5°C and 3°C GWL (Figure 13.8)

Terrestrial and freshwater ecosystems	Supporting references of assessment
Reduction in habitat availability of cold-adapted groups	Balint et al. (2011); Dornelas et al. (2014); Hodd et al. (2014); Hubble (2014); Kovats et al. (2014); Oliver et al. (2014); McGill et al. (2015); Oliver et al. (2015); Saltré et al. (2015); Talavera et al. (2015); Barredo et al. (2016); Coll et al. (2016); Hellmann et al. (2016); Jørgensen et al. (2016); Liverpool (2016); Dapporto et al. (2017); EEA (2017b); Vermaat et al. (2017); Ciscar et al. (2018); Hillebrand et al. (2018); Sirois-Delisle and Kerr (2018); Suggitt et al. (2018); Warren et al. (2018); Habel et al. (2019); Hinojosa et al. (2019); van Strien et al. (2019); Dullinger et al. (2020); Outhwaite et al. (2020); Soroye et al. (2020); Xi (2020); Carnicer et al. (2021)
Reduction in biodiversity of cold-adapted groups	Balint et al. (2011); Stefanescu et al. (2011); Dornelas et al. (2014); Oliver et al. (2014); Zografou et al. (2014); Hill and Preston (2015); McGill et al. (2015); Oliver et al. (2015); Talavera et al. (2015); Hellmann et al. (2016); Hendriks et al. (2016); Jørgensen et al. (2016); Rizzetto et al. (2016); Stephens et al. (2016); Vodá et al. (2016); Dapporto et al. (2017); EEA (2017b); Vermaat et al. (2017); Dyderski et al. (2018); Hillebrand et al. (2018); Sirois-Delisle and Kerr (2018); Spooner et al. (2018); Suggitt et al. (2018); Warren et al. (2018); Dennis et al. (2019); Habel et al. (2019); Herrando et al. (2019); Hinojosa et al. (2019); van Strien et al. (2019); Dullinger et al. (2020); Kougiumoutzis et al. (2020); Outhwaite et al. (2020); Soroye et al. (2020); Xi (2020)
Range shifts	Parmesan et al. (1999); Wilson et al. (2007); Devictor et al. (2008); Lenoir et al. (2008); Jiguet et al. (2010); Chen et al. (2011); Scherrer and Körner (2011); Devictor et al. (2012); De Frenne et al. (2013); Lenoir et al. (2013); Kovats et al. (2014); Bowler et al. (2015); Ancillotto et al. (2016); Jørgensen et al. (2016); Stephens et al. (2016); Wiens (2016); Bowler et al. (2017); EEA (2017b); Massimino et al. (2017); Mills et al. (2017); Pearce-Higgins et al. (2017); Sáenz-Romero et al. (2017); Bowler et al. (2018); Dyderski et al. (2018); Mori et al. (2018); Rumpf et al. (2018); Sirois-Delisle and Kerr (2018); Spooner et al. (2018); Steinbauer et al. (2018); Suggitt et al. (2018); Bowler et al. (2019); Carnicer et al. (2019b); Gómez (2019); Jaime et al. (2019); Lehikoinen et al. (2019a); Pérez Navarro et al. (2019); Post et al. (2019); Termaat et al. (2019); Vilà-Cabrera et al. (2019); Margalef-Marrase et al. (2020); Pavón-Jordán et al. (2020); Soroye et al. (2020); van Klink et al. (2020a); Zellweger et al. (2020); Urvois et al. (2021)
Changes in phenology	Ovaskainen et al. (2013); Thackeray et al. (2013); Frolov et al. (2014); Garonna et al. (2014); Karlsson (2014); Plard et al. (2014); Schröder et al. (2014); van Vliet et al. (2014); Fu et al. (2015); Gill et al. (2015); Malcolm et al. (2015); Roberts et al. (2015); Gaüzère et al. (2016); Newson et al. (2016); Szabó et al. (2016); Thackeray et al. (2016a); EEA (2017b); Gauzere et al. (2017); Glushenkov (2017); Güsewell et al. (2017); Halupka and Halupka (2017); Mayor et al. (2017); Miles et al. (2017); Prokosheva (2017); Wang et al. (2017a); Asse et al. (2018); Chen et al. (2018); Chizhikova (2018); Cohen et al. (2018b); Donnelly et al. (2018); Hidalgo-Galvez et al. (2018); Posledovich et al. (2018); Vitasse et al. (2018); Wu et al. (2018); Bobretsov et al. (2019); Fraga et al. (2019); Jakoby et al. (2019b); Lehikoinen et al. (2019a); Ma et al. (2019); Macgregor et al. (2019); Peaucelle et al. (2019); Piao et al. (2019); Prislán et al. (2019); Tishkov et al. (2019); Delgado et al. (2020); Menzel et al. (2020); Orellana-Macias et al. (2020); Wang et al. (2020); Keogan et al. (2021); Rosbakh et al. (2021)
Decrease in ecosystem production	Nabuurs et al. (2003); Ciais et al. (2005); Schröter et al. (2005); Smith et al. (2005); Reichstein et al. (2007); Schulze et al. (2009); Carnicer et al. (2011); Fantappiè et al. (2011); Elmendorf et al. (2012); Carnicer et al. (2013); Coll et al. (2013); Peñuelas et al. (2013); Kovats et al. (2014); Ruiz-Benito et al. (2014); Schröter et al. (2014); Gazol et al. (2015); Keenan et al. (2016); Naudts et al. (2016); Novick et al. (2016); Polce et al. (2016); Schubert et al. (2016); Tian et al. (2016); van der Plas et al. (2016); Yigini and Panagos (2016); Ballantyne et al. (2017); Bright et al. (2017); EASAC (2017); EEA (2017b); Nabuurs et al. (2017); Peñuelas et al. (2017a); Peñuelas et al. (2017b); Ratcliffe et al. (2017); Schwalm et al. (2017); Teuling et al. (2017); Valade et al. (2017); Gazol et al. (2018); Humphrey et al. (2018); Lugato et al. (2018); Luyssaert et al. (2018); Nabuurs (2018); Sanginés de Cárcer et al. (2018); Stocker et al. (2018); Torralba et al. (2018); Verhagen et al. (2018); Carnicer et al. (2019a); Ciais et al. (2019); EASAC (2019); Fernández-Martínez et al. (2019); Green et al. (2019); Jaime et al. (2019); Lee et al. (2019); Natali et al. (2019); Pérez Navarro et al. (2019); Post et al. (2019); Stocker et al. (2019); Xu et al. (2019); Yuan et al. (2019); Zhou et al. (2019); Batllori et al. (2020); Brodribb et al. (2020); Ito et al. (2020); Krause et al. (2020); Lian et al. (2020); Margalef-Marrase et al. (2020); Schuldt et al. (2020); Wang (2020); Zhang et al. (2020); Canadell and Jackson (2021); Rocas-Díaz et al. (2021); Yu et al. (2021)
Rising incidence of fire	Moriondo et al. (2006); Bondur (2011); Dury et al. (2011); Bedia et al. (2014); Turco et al. (2014); Drobyshev et al. (2015); Jolly et al. (2015); Tedim et al. (2015); Wu et al. (2015b); Drobyshev et al. (2016); Khabarov et al. (2016); Regos et al. (2016); Turco et al. (2016); Camia (2017); de Rigo et al. (2017); Forzieri et al. (2017); Fréjaville (2017); Ruffault et al. (2017); Sitnov et al. (2017); Turco et al. (2017); Bedia et al. (2018); Filipchuk et al. (2018); Lahaye et al. (2018); San-Miguel-Ayanz et al. (2018); Sitnov and Mokhov (2018); Turco et al. (2018a); Turco et al. (2018b); Chergui (2019); Michetti and Pinar (2019); Pausas (2019); Costa (2020); Di Giuseppe et al. (2020); Dupuy et al. (2020); Royé et al. (2020)
Reduced pollination services (reduction of regulating ecosystem services)	Menéndez et al. (2006); Roberts et al. (2011); Franzén and Öckinger (2012); Carvalheiro et al. (2013); Polce et al. (2014); Kaloveloni et al. (2015); Kerr et al. (2015); Rasmont et al. (2015); Tzilivakis et al. (2015); Ipbes (2016); Petz et al. (2016); Settele et al. (2016); Radenković et al. (2017); Marshall et al. (2018); Fourcade et al. (2019); Powney et al. (2019); Steele et al. (2019); Van Dooren (2019); Soroye et al. (2020); Zattara and Aizen (2020); Vasiliev and Greenwood (2021)
Increased soil erosion (reduction of regulating ecosystem services)	Bangash et al. (2013); Mezösi et al. (2013); Routschek et al. (2014); Anaya-Romero et al. (2015); Cilek et al. (2015); European Commission (2015); Panagos et al. (2015); Serpa et al. (2015); Sobol et al. (2015); Tzilivakis et al. (2015); Adler-Nissen (2016); Borrelli et al. (2016); Grossel et al. (2016); Guerra et al. (2016); Polce et al. (2016); Li et al. (2017); Litvin et al. (2017); Panagos et al. (2017); Prins et al. (2017); Chizhikova (2018); Gusarov et al. (2018); Auerswald and Fiener (2019); Mullan et al. (2019); Pastor et al. (2019a); Berberoglu et al. (2020); Borrelli et al. (2020); Ciampalini et al. (2020); Gianinetto et al. (2020); Gusarov (2020); Luetzenburg et al. (2020); Morán-Ordóñez et al. (2020); Rodrigues et al. (2020); Svetlitchnyi (2020)

Table SM13.3 | Percentage of species per group projected to remain within their suitable climate conditions (Figure 13.9)

Plants														
Regions	1.5°C GWL	Std		2°C GWL	Std		2.7°C GWL	Std		3.2°C GWL	Std		4.5°C GWL	Std
WCE	0.77	0.08		0.70	0.09		0.58	0.10		0.53	0.10		0.39	0.10
NEU	0.85	0.18		0.81	0.19		0.76	0.20		0.74	0.21		0.67	0.23
SEU	0.75	0.09		0.67	0.11		0.53	0.12		0.47	0.12		0.31	0.12
EEU	0.75	0.12		0.69	0.13		0.58	0.14		0.52	0.15		0.35	0.14
Insects														
Regions	1.5°C GWL	Std		2°C GWL	Std		2.7°C GWL	Std		3.2°C GWL	Std		4.5°C GWL	Std
WCE	0.59	0.11		0.49	0.11		0.34	0.11		0.29	0.10		0.17	0.08
NEU	0.90	0.14		0.86	0.18		0.76	0.25		0.72	0.28		0.58	0.34
SEU	0.65	0.13		0.56	0.15		0.43	0.17		0.38	0.17		0.26	0.15
EEU	0.72	0.15		0.64	0.18		0.49	0.20		0.42	0.21		0.25	0.19
Pollinator														
Regions	1.5°C GWL	Std		2°C GWL	Std		2.7°C GWL	Std		3.2°C GWL	Std		4.5°C GWL	Std
WCE	0.59	0.14		0.50	0.15		0.39	0.15		0.35	0.14		0.26	0.11
NEU	0.84	0.18		0.78	0.23		0.65	0.29		0.59	0.31		0.43	0.33
SEU	0.75	0.13		0.69	0.14		0.57	0.16		0.52	0.16		0.38	0.15
EEU	0.69	0.21		0.60	0.23		0.43	0.25		0.36	0.24		0.19	0.19
Amphibians														
Regions	1.5°C GWL	Std		2°C GWL	Std		2.7°C GWL	Std		3.2°C GWL	Std		4.5°C GWL	Std
WCE	0.88	0.12		0.84	0.14		0.76	0.17		0.72	0.17		0.59	0.18
NEU	0.89	0.24		0.91	0.16		0.87	0.19		0.84	0.21		0.76	0.25
SEU	0.84	0.16		0.79	0.18		0.68	0.21		0.63	0.22		0.50	0.22
EEU	0.83	0.29		0.86	0.20		0.78	0.24		0.75	0.25		0.63	0.30
Reptiles														
Regions	1.5°C GWL	Std		2°C GWL	Std		2.7°C GWL	Std		3.2°C GWL	Std		4.5°C GWL	Std
WCE	0.89	0.09		0.86	0.10		0.80	0.11		0.77	0.11		0.67	0.13
NEU	0.90	0.18		0.87	0.19		0.82	0.22		0.79	0.23		0.71	0.25
SEU	0.89	0.10		0.85	0.12		0.76	0.15		0.72	0.15		0.58	0.15
EEU	0.87	0.27		0.84	0.28		0.79	0.30		0.78	0.31		0.68	0.33
Birds														
Regions	1.5°C GWL	Std		2°C GWL	Std		2.7°C GWL	Std		3.2°C GWL	Std		4.5°C GWL	Std
WCE	0.88	0.04		0.85	0.05		0.79	0.06		0.76	0.07		0.66	0.11
NEU	0.93	0.13		0.92	0.12		0.90	0.12		0.88	0.13		0.83	0.16
SEU	0.87	0.06		0.82	0.08		0.73	0.10		0.69	0.11		0.54	0.12
EEU	0.86	0.14		0.84	0.15		0.79	0.16		0.77	0.16		0.69	0.18

Mammals														
Regions	1.5°C GWL	Std		2°C GWL	Std		2.7°C GWL	Std		3.2°C GWL	Std		4.5°C GWL	Std
WCE	0.79	0.10		0.73	0.11		0.61	0.12		0.55	0.12		0.42	0.12
NEU	0.90	0.11		0.87	0.12		0.81	0.16		0.77	0.17		0.63	0.23
SEU	0.78	0.12		0.70	0.14		0.59	0.16		0.54	0.16		0.38	0.15
EEU	0.80	0.16		0.75	0.19		0.65	0.20		0.60	0.19		0.44	0.16

Notes:

WCE: Western Central Europe, NEU: Northern Europe, SEU: Southern Europe, EEU: Eastern Europe.

Species are projected to remain within their suitable climate conditions at increasing levels of climate change averaged over 21 CMIP5 climate models with standard deviation (Std) (Warren et al., 2018). There is increased loss of climatic niche with warming increasing from 1.5°C to 4.5°C GWL. Risks that appear to be lower in some groups through droughts, habitat fragmentation and loss are not considered and will exacerbate the risks, while dispersal may reduce risk.

SM13.3 Supplementary Material Supporting Section 13.4

Table SM13.4 | Literature sources used for assessment of impacts on and risks for marine and coastal ecosystems (Figure 13.11)

Impact/risk	Supporting references of assessment
Loss of habitat availability	Coma et al. (2009); Garrabou et al. (2009); Huete-Stauffer et al. (2011); Munari (2011); Kersting et al. (2013); Brodie et al. (2014); Frolov et al. (2014); Rivetti et al. (2014); Altieri and Gedan (2015); García Molinos et al. (2016); Spencer et al. (2016); Bakanev (2017); Jessen et al. (2017); Orekhova (2017); Berlinski and Popov (2018); Buonomo et al. (2018); Jokinen et al. (2018); Reusch et al. (2018); Schuerch et al. (2018); van der Spek (2018); Wang et al. (2018); Filatov et al. (2019); Garrabou et al. (2019); Saraiva et al. (2019); Spivak et al. (2019); D'Amen and Azzurro (2020); Jiang et al. (2020); Pavlova (2020); Sandø et al. (2020); Stepanyan (2020)
Shifts in ranges (including invasions), composition (taxonomic, functional), phenologies	Kortsch et al. (2015); Assis et al. (2017); Bakanev (2017); Frainer et al. (2017); Kotenev et al. (2017); Rasmussen et al. (2017); Raybaud et al. (2017); Townhill et al. (2017); Vasilakopoulos et al. (2017); Benedetti et al. (2018); Gaudin et al. (2018); Jonsson et al. (2018); Kotta et al. (2018); Minicheva et al. (2018); Townhill et al. (2018); Benedetti et al. (2019); Berdnikov et al. (2019); Casado-Amezúa et al. (2019); Chefaoui et al. (2019); de la Hoz et al. (2019); Erauskin-Extramiana et al. (2019); Filatov et al. (2019); Hjerne et al. (2019); Kröncke et al. (2019); Krovnin et al. (2019); Moullec et al. (2019); Wasmund et al. (2019); Baudron et al. (2020); Bedford et al. (2020); Clark et al. (2020); Desmit et al. (2020); Maltby et al. (2020); Martynova et al. (2020); Nohe et al. (2020); Pavlova (2020); Pecuchet et al. (2020); Pennino et al. (2020); Pyatinsky et al. (2020); Stepanyan (2020); Uriarte et al. (2021)
Reduction in growth and reproductive success	Bramanti et al. (2013); Maier et al. (2013); Gazeau et al. (2014); Hennige et al. (2015); Wall et al. (2015); Ragazzola et al. (2016); Stiasny et al. (2016); Durant and Hjermann (2017); Smoliński and Mirny (2017); Thomsen et al. (2017); Capuzzo et al. (2018); Lindegren et al. (2018); Queirós et al. (2018); Sswat et al. (2018a); Sswat et al. (2018b); Stiasny et al. (2018); Coll et al. (2019); Franz et al. (2019); Goldberg et al. (2019); Herrera et al. (2019); Hidalgo et al. (2019); Sguotti et al. (2019a); Stiasny et al. (2019); Tanner et al. (2019); Tsikliras et al. (2019); Verezemskaya et al. (2019); Vieira et al. (2019); Voss et al. (2019); Denechaud et al. (2020); Maynou et al. (2020); Mitchell et al. (2020); Tanner et al. (2020); Ikpewe et al. (2021); Polte et al. (2021)
Loss of biodiversity	Berlinski and Popov (2018); IPBES (2018); Filatov et al. (2019); Pyatinsky et al. (2020); Stepanyan (2020)
Decline in production	Maugendre et al. (2014); Arrigo and van Dijken (2015); Laufkotter et al. (2015); Holt et al. (2016); Børshheim (2017); Orekhova (2017); Capuzzo et al. (2018); Holt et al. (2018); Minicheva et al. (2018); Berdnikov et al. (2019); Bryndum-Buchholz et al. (2019); Carozza et al. (2019); Free et al. (2019); Kwiatkowski et al. (2019); Lotze et al. (2019b); Verezemskaya et al. (2019); Lewis et al. (2020); Pyatinsky et al. (2020)
Emergence of harmful algal blooms and pathogens	Frolov et al. (2014); Baker-Austin et al. (2017); Semenza et al. (2017); Minicheva et al. (2018); Riebesell et al. (2018); Roggatz et al. (2019)
Reduction in ecosystem services	Roebeling et al. (2013); Brodie et al. (2014); Carstensen et al. (2014); Kjesbu et al. (2014); Maugendre et al. (2014); Serra et al. (2015); Krumhansl et al. (2016); De los Santos et al. (2017); Gao et al. (2018); van der Spek (2018); Wang et al. (2018); Moullec et al. (2019); Sguotti et al. (2019b); Sheverdyayev (2019); Baudron et al. (2020); Maltby et al. (2020)

SM13.4 Supplementary Material Supporting Section 13.5

Table SM13.5 | Literature sources used in the assessment of feasibility and effectiveness of adaptation options for food systems in Europe (Figure 13.14)

Impact type	Adaptation option	Effectiveness	Feasibility					
			Economic	Technological	Institutional	Sociocultural	Ecological	Geophysical
Heat stress	Irrigation	Dono et al. (2013); Schaap et al. (2013); Sutton et al. (2013); Bird et al. (2016); Diogo et al. (2017); Webber et al. (2018); Feyen et al. (2020); Holzkämper, 2020); Santillán et al. (2020); Kebede et al. (2021)	Dono et al. (2013); Schaap et al. (2013); Sutton et al. (2013); Bird et al. (2016); Costa et al. (2016); Diogo et al. (2017); Holzkämper (2020); Kebede et al. (2021)	Dono et al. (2013); Schaap et al. (2013); Sutton et al. (2013); Bird et al. (2016); Costa et al. (2016); Diogo et al. (2017); Neset et al. (2019); Holzkämper (2020); Santillán et al. (2020); Kebede et al. (2021)	Schaap et al. (2013); Mandryk et al. (2015); Bird et al. (2016); Kebede et al. (2021)	Schaap et al. (2013); Sutton et al. (2013); Mandryk et al. (2015); Costa et al. (2016); Neset et al. (2019); Kebede et al. (2021)	Dono et al. (2013); Sutton et al. (2013); Bird et al. (2016); Costa et al. (2016); Siebert et al. (2017); Webber et al. (2018); Neset et al. (2019); Holzkämper (2020); Santillán et al. (2020); Kebede et al. (2021)	Dono et al. (2013); Webber et al. (2018); Neset et al. (2019); Kebede et al. (2021)
	Change of sowing/ harvest date	Schaap et al. (2013); Trnka et al. (2014); Donatelli et al. (2015); Gabaldon-Leal et al. (2015); Peltonen-Sainio et al. (2016); Diogo et al. (2017); Feyen et al. (2020); Holzkämper (2020)	Schaap et al. (2013); Trnka et al. (2014); Donatelli et al. (2015); Diogo et al. (2017); Grüneis et al. (2018)	Trnka et al. (2014); Donatelli et al. (2015); Gabaldon-Leal et al. (2015); Feyen et al. (2020); Holzkämper (2020)		Schaap et al. (2013); Donatelli et al. (2015); Mandryk et al. (2015); Peltonen-Sainio et al. (2016); Diogo et al. (2017); Grüneis et al. (2018)	Donatelli et al. (2015); Peltonen-Sainio et al. (2016); Holzkämper (2020)	Schaap et al. (2013); Diogo et al. (2017); Holzkämper (2020)
	Change of cultivars	Sutton et al. (2013); Trnka et al. (2014); Gabaldon-Leal et al. (2015); Costa et al. (2016); Peltonen-Sainio et al. (2016); Rial-Lovera et al. (2017); Webber et al. (2018); Santillán et al. (2020)	Sutton et al. (2013); Trnka et al. (2014); Rial-Lovera et al. (2017); Grüneis et al. (2018); Holzkämper (2020)	Sutton et al. (2013); Trnka et al. (2014); Gabaldon-Leal et al. (2015); Costa et al. (2016); Rial-Lovera et al. (2017); Webber et al. (2018); Feyen et al. (2020); Holzkämper (2020); Santillán et al. (2020)		Sutton et al. (2013); Costa et al. (2016); Peltonen-Sainio et al. (2016); Grüneis et al. (2018)	Sutton et al. (2013); Rial-Lovera et al. (2017); Holzkämper (2020); Santillán et al. (2020)	Rial-Lovera et al. (2017); Webber et al. (2018); Santillán et al. (2020)
	Livestock management	Vitali et al. (2015); Cox et al. (2016); Schaubberger et al. (2020)	Vitali et al. (2015); Schaubberger et al. (2020)	Morignat et al. (2014); Vitali et al. (2015); Cox et al. (2016); Schaubberger et al. (2020)		Vitali et al. (2015); Cox et al. (2016)	Morignat et al. (2014); Vitali et al. (2015); Cox et al. (2016)	

		Effectiveness	Feasibility					
Impact type	Adaptation option		Economic	Technological	Institutional	Sociocultural	Ecological	Geophysical
Drought	Irrigation	Dono et al. (2013); Schaap et al. (2013); Sutton et al. (2013); Bird et al. (2016); Diogo et al. (2017); Stańczuk-Galwiczek et al. (2018); Webber et al. (2018); Harmanny and Malek (2019); Holzkämper (2020); Santillán et al. (2020)	Dono et al. (2013); Schaap et al. (2013); Sutton et al. (2013); Costa et al. (2016); Diogo et al. (2017); Harmanny and Malek (2019); Holzkämper (2020); Kebede et al. (2021)	Dono et al. (2013); Schaap et al. (2013); Sutton et al. (2013); Mandryk et al. (2015); van Duinen et al. (2015); Bird et al. (2016); Costa et al. (2016); Diogo et al. (2017); Stańczuk-Galwiczek et al. (2018); Harmanny and Malek (2019); Feyen et al. (2020); Holzkämper (2020); Santillán et al. (2020); Kebede et al. (2021)	Schaap et al. (2013); Mandryk et al. (2015); Bird et al. (2016); Grüneis et al. (2018); Kebede et al. (2021)	Schaap et al. (2013); Sutton et al. (2013); Mandryk et al. (2015); van Duinen et al. (2015); Costa et al. (2016); Grüneis et al. (2018); Stańczuk-Galwiczek et al. (2018); Harmanny and Malek (2019); Kebede et al. (2021)	Dono et al. (2013); Sutton et al. (2013); Bird et al. (2016); Costa et al. (2016); Stańczuk-Galwiczek et al. (2018); Webber et al. (2018); Harmanny and Malek (2019); Holzkämper (2020); Santillán et al. (2020); Kebede et al. (2021)	Dono et al. (2013); Diogo et al. (2017); Harmanny and Malek (2019); Santillán et al. (2020); Kebede et al. (2021)
	Change of sowing/ harvest date	Schaap et al. (2013); Trnka et al. (2014); Donatelli et al. (2015); Gabaldon-Leal et al. (2015); Peltonen-Sainio et al. (2016); Diogo et al. (2017); Parent et al. (2018); Lamichhane et al. (2019); Feyen et al. (2020); Holzkämper (2020)	Schaap et al. (2013); Trnka et al. (2014); Donatelli et al. (2015); Diogo et al. (2017); Grüneis et al. (2018); Lamichhane et al. (2019)	Trnka et al. (2014); Donatelli et al. (2015); Gabaldon-Leal et al. (2015); Feyen et al. (2020); Holzkämper (2020)		Schaap et al. (2013); Donatelli et al. (2015); Mandryk et al. (2015); Peltonen-Sainio et al. (2016); Diogo et al. (2017); Grüneis et al. (2018)	Donatelli et al. (2015); Peltonen-Sainio et al. (2016); Holzkämper (2020)	Schaap et al. (2013); Diogo et al. (2017); Holzkämper (2020)
	Change of cultivars	Sutton et al. (2013); Trnka et al. (2014); Gabaldon-Leal et al. (2015); Costa et al. (2016); Peltonen-Sainio et al. (2016); Rial-Lovera et al. (2017); Parent et al. (2018); Webber et al. (2018); Santillán et al. (2020)	Sutton et al. (2013); Trnka et al. (2014); Rial-Lovera et al. (2017); Grüneis et al. (2018); Holzkämper (2020)	Sutton et al. (2013); Trnka et al. (2014); Gabaldon-Leal et al. (2015); Costa et al. (2016); Rial-Lovera et al. (2017); Webber et al. (2018); Feyen et al. (2020); Holzkämper (2020); Santillán et al. (2020)		Sutton et al. (2013); Costa et al. (2016); Peltonen-Sainio et al. (2016); Grüneis et al. (2018)	Sutton et al. (2013); Rial-Lovera et al. (2017); Holzkämper (2020); Santillán et al. (2020)	Rial-Lovera et al. (2017); Webber et al. (2018); Santillán et al. (2020)
	Soil management	Schönhart et al. (2014); Rial-Lovera et al. (2017); Hamidov et al. (2018); Wiréhn (2018); Jørgensen et al. (2020); Wiréhn et al. (2020)	Schönhart et al. (2014); Rial-Lovera et al. (2017); Hamidov et al. (2018); Wiréhn (2018); EEA (2019b); Wiréhn et al. (2020)	Schönhart et al. (2014); Rial-Lovera et al. (2017); Hamidov et al. (2018); Wiréhn (2018); Wiréhn et al. (2020)	Rial-Lovera et al. (2017); Wiréhn (2018); Jørgensen et al. (2020)	Rial-Lovera et al. (2017); Wiréhn (2018)	Rial-Lovera et al. (2017); Hamidov et al. (2018); Wiréhn (2018)	Rial-Lovera et al. (2017); Hamidov et al. (2018); Wiréhn (2018)
Flooding	Change of sowing/ harvest date	Sutton et al. (2013); Rial-Lovera et al. (2017); Stańczuk-Galwiczek et al. (2018); Neset et al. (2019)	Rial-Lovera et al. (2017); Neset et al. (2019); Kebede et al. (2021)	Sutton et al. (2013); Papadaskalopoulou et al. (2016); Rial-Lovera et al. (2017); Stańczuk-Galwiczek et al. (2018); Neset et al. (2019)			Sutton et al. (2013); Stańczuk-Galwiczek et al. (2018); Neset et al. (2019); Kebede et al. (2021)	Sutton et al. (2013); Stańczuk-Galwiczek et al. (2018); Neset et al. (2019); Kebede et al. (2021)

Impact type	Adaptation option	Effectiveness		Feasibility				
			Economic	Technological	Institutional	Sociocultural	Ecological	Geophysical
Compound and extreme weather	Plant and livestock breeding	Trnka et al. (2014); Macholdt and Honermeier (2017); Rial-Lovera et al. (2017); Costa et al. (2019a); Senapati et al. (2019); Wreford and Topp (2020)	Trnka et al. (2014); Macholdt and Honermeier (2017); Rial-Lovera et al. (2017); Costa et al. (2019a); Senapati et al. (2019); Wreford and Topp (2020)	Rial-Lovera et al. (2017); Costa et al. (2019a); Senapati et al. (2019); Wreford and Topp (2020)			Wiréhn (2018); Costa et al. (2019a); Holzkämper (2020); Wreford and Topp (2020)	Rial-Lovera et al. (2017); Wreford and Topp (2020)
	Mixed use: agroecology and agroforestry	Lüscher et al. (2014); Moraine et al. (2014); Himanen et al. (2016); Hernández-Morcillo et al. (2018)	Lüscher et al. (2014); Moraine et al. (2014); Himanen et al. (2016); Rojas-Downing et al. (2017); Hernández-Morcillo et al. (2018)	Moraine et al. (2014); Hernández-Morcillo et al. (2018)	Moraine et al. (2014); Hernández-Morcillo et al. (2018)	Moraine et al. (2014); Hernández-Morcillo et al. (2018)	Moraine et al. (2014); Hernández-Morcillo et al. (2018); Öllerer et al. (2019); Oggioni et al. (2020)	Prem et al. (2014); Fornara et al. (2019); Oggioni et al. (2020); Ford et al. (2021)
	Agricultural policy changes	Papadaskalopoulou et al. (2016); Erjavec et al. (2017); McVittie et al. (2018); Muenzel and Martino (2018); Faria and Morales (2020)	Buelow and Cradock-Henry (2018); Grüneis et al. (2018); McVittie et al. (2018); Muenzel and Martino (2018); Sneessens et al. (2019); Faria and Morales (2020)	Li et al. (2017); Wiréhn (2018); EEA (2019b); Szumelda (2019)	Reidsma et al. (2015); Papadaskalopoulou et al. (2016); Li et al. (2017); Grüneis et al. (2018); McVittie et al. (2018); Muenzel and Martino (2018); Wiréhn (2018); Sneessens et al. (2019); Szumelda (2019); Faria and Morales (2020); Jørgensen et al. (2020); Mitter et al. (2020)	Li et al. (2017); Buelow and Cradock-Henry (2018); Nguyen et al. (2019); Sneessens et al. (2019); Szumelda (2019)	McVittie et al. (2018); Muenzel and Martino (2018); Faria and Morales (2020)	McVittie et al. (2018); Muenzel and Martino (2018); Faria and Morales (2020)
	Training and information	Li et al. (2017); Rial-Lovera et al. (2017); Buelow and Cradock-Henry (2018); Nguyen et al. (2019)	Rial-Lovera et al. (2017); McVittie et al. (2018); Szumelda (2019)		Li et al. (2017); Rial-Lovera et al. (2017); McVittie et al. (2018)	Nguyen et al. (2019); Szumelda (2019)	Li et al. (2017); Rial-Lovera et al. (2017); McVittie et al. (2018)	Li et al. (2017); Rial-Lovera et al. (2017); McVittie et al. (2018)
	Crop selection changes	Lüscher et al. (2014); Trnka et al. (2015); Rial-Lovera et al. (2017); Li et al. (2018); Harmanny and Malek (2019)	Lüscher et al. (2014); Reidsma et al. (2015); Wiréhn (2018); Wiréhn et al. (2020)	Lüscher et al. (2014); Trnka et al. (2015); Rial-Lovera et al. (2017); Li et al. (2018); Wiréhn et al. (2020)		Himanen et al. (2016); Ricart et al. (2019)	Rial-Lovera et al. (2017); Li et al. (2018); Harmanny and Malek (2019); Wiréhn et al. (2020)	Rial-Lovera et al. (2017); Li et al. (2018); Harmanny and Malek (2019); Wiréhn et al. (2020)
	Land cover change (including agricultural land abandonment)	Leclère et al. (2013); Dunford et al. (2015); Kebede et al. (2021)	Dunford et al. (2015); Alexander et al. (2018); Kebede et al. (2021)	Leclère et al. (2013); Mandryk et al. (2015); Alexander et al. (2018)	Mandryk et al. (2015)	Leclère et al. (2013); Mandryk et al. (2015); Neset et al. (2019)	Dunford et al. (2015); Rabin et al. (2020); Kebede et al. (2021)	Dunford et al. (2015); Rabin et al. (2020); Kebede et al. (2021)

		Effectiveness	Feasibility					
Impact type	Adaptation option		Economic	Technological	Institutional	Sociocultural	Ecological	Geophysical
Disease pathogens and vectors	Plant and livestock breeding	Hoffmann (2013)	Hoffmann (2013)	Hoffmann (2013)	Hoffmann (2013); Grüneis et al. (2018)	Hoffmann (2013); Neset et al. (2019)		
	Management (including high-frequency rotations)	Maclachlan and Guthrie (2010); Skuce et al. (2013); Moraine et al. (2014)	Maclachlan and Guthrie (2010); Morgan and van Dijk (2012); Wiréhn et al. (2020)	Dórea et al. (2016); Harrus and Baneth (2005); Moraine et al. (2014); Pascual-Linaza et al. (2014); Skuce et al. (2013)	Harrus and Baneth (2005); Moraine et al. (2014); Roberts et al. (2014); Dórea et al. (2016)	Morgan and van Dijk (2012); Wiréhn et al. (2020)	Acevedo et al. (2010); Maclachlan and Guthrie (2010); Martínez-López et al. (2014); Moraine et al. (2014); Rose Vineer et al. (2020)	Maclachlan and Guthrie (2010); Moraine et al. (2014); Paz (2015); Tjaden et al. (2018); Rose Vineer et al. (2020)
Combined impacts on productivity	International trade changes	Dunford et al. (2015); Holman et al. (2016); Alexander et al. (2018); EEA (2019b)				Dunford et al. (2015); Holman et al. (2016); Mitter et al. (2020); Kebede et al. (2021)	Dunford et al. (2015); Holman et al. (2016); Kebede et al. (2021)	Alexander et al. (2018); EEA (2019b)
	Consumer shifts in consumption		Dunford et al. (2015); Mitter et al. (2020)			Dunford et al. (2015); Mitter et al. (2020)		

SM13.5 Supplementary Material Supporting Section 13.6

Table SM13.6 | Sign of future change in onshore wind power potential under global warming levels (Figure 13.16)

Sub-region	Area in study	Onshore wind power potential (sign of change)					
		1.5°C		2°C		≥3°C	
Northern Europe							
Davy et al. (2018)	Europe, North Africa, Middle East	–		–	No model agreement over large areas	–	No model agreement over large areas
Moemken et al. (2018)	Europe	–		+/–	+ only over a minor part of Scandinavia W: + in central Scandinavia, – in the rest S: – except northern Norway	+/–	+ over northern Scandinavia W: + in almost all areas S: –
Devis et al. (2018)	Europe	+	W: + S: + except in the Baltic				
Tobin et al. (2018a)	EU and Switzerland	–	Some ensemble members project increases up to 5%	–	Some ensemble members project increases up to 4%	–	Some ensemble members project increases up to 4%
Reyers et al. (2016)	Europe (without Russia)			+	W: +, S: –	+	W: +, S: –
Carvalho et al. (2017b)	Europe (without Russia)	–	– in southern UK and northern Norway, rest: n. s. W: few –, rest: n. s. S: few –, most of Scandinavia: n. s.	+/–	+ in southern Finland, in southern Norway: n. s. W: few –, rest: n. s. S: +/–, most of Scandinavia: n. s.	+/–	+ in southern Finland, in southern Norway: n. s. W: few –, rest: n. s. S: +/–, most of Scandinavia: n. s.
Tobin et al. (2016)	Europe	–		–			
Després and Adamovic (2020)	EU	n.c.		n.c.		–	

Sub-region	Area in study	Onshore wind power potential (sign of change)					
		1.5°C		2°C		≥3°C	
Western Central Europe							
Reyers et al. (2016)	Europe (without Russia)			n.c./+/-	n.c. over half of France, – in the rest of France	n.c./+/-	n.c. over half of France, – in the rest of France
Moemken et al. (2018)	Europe	–		–	W: +/- (and opposite signs of change between RCMs over large areas) S: +/-	+/-	+ over coastal Poland W: +/- (and opposite signs of change between RCMs over most areas) S: +/- (but – in most areas)
Davy et al. (2018)	Europe, North Africa, Middle East	–	No model agreement over France, Belarus, Ukraine	–	No model agreement over Poland, Belarus	+/-	No model agreement over large areas; parts of Ukraine and Belarus with +
Carvalho et al. (2017b)	Europe (without Russia)	–	Changes n. s. over large areas W: –, S: –	–	Changes n. s. over large areas W: –, S: –	–	Changes n. s. over large areas W: –, S: –
Devis et al. (2018)	Europe	+/-	– in Belarus, Ukraine and most of France W: +/-, S: +/-				
Tobin et al. (2018a)	EU and Switzerland	–		–		–	
Tobin et al. (2016)	Europe	+/-	– for Poland	+			
Després and Adamovic (2020)	EU	n.c.		n.c.		+	
Southern Europe							
Solaun and Cerdá (2020)	Spain	+/-		+/-			
Katopodis et al. (2019)	Greece	+/-		+/-			
Reyers et al. (2016)	Europe			+/-	+ only over a few areas in Turkey and Greece; high robustness in – sign W: –, S: +/-	–	High robustness in sign W: –, S: +/-
Moemken et al. (2018)	Europe	+/-	+ in Turkey, large parts of Italy and Greece, and southern France	+/-	+ in Turkey W: – except southern France and central Italy S: + in Turkey and most of Spain, – in the rest	+/-	+ in coastal Turkey W: – except Croatia, central Italy and southern France where models disagree S: +/-
Davy et al. (2018)	Europe, North Africa, Middle East	+/-	+ over coastal Turkey and a few more locations	+/-	+ over coastal Turkey and a few more locations	+/-	+ over a small part of coastal Turkey
Devis et al. (2018)	Europe	–	W: –, S: + in southern Iberia, in the rest – in the day and + in the night				

Sub-region	Area in study	Onshore wind power potential (sign of change)					
		1.5°C		2°C		≥3°C	
Carvalho et al. (2017b)	Europe	–	In Turkey, changes n. s. W: few –, rest: n. s. S: +/-, most: n. s.	–	In Turkey, changes n. s. W: few –, rest: n. s. S: +/-, most: n. s.	+/-	+ over northern Turkey W: –, Turkey mostly n. s. S: +/- (+ in Turkey and northern Iberia)
Tobin et al. (2018a)	EU and Switzerland	+/-	+ only for Greece	+/-	+ only for Greece	+/-	+ only for Greece
Tobin et al. (2016)	Europe	–		–			
Després and Adamovic (2020)	EU	n.c.		n.c.		+	
Eastern Europe							
Devis et al. (2018)		+	W: +, S: +/-				
Carvalho et al. (2017b)		–	W: –, S: –	–	W: –, S: –	–	W: –, S: –
Davy et al. (2018)	Europe, North Africa, Middle East	–		–	No model agreement over a large part of the sub-region	–	No model agreement over most of the sub-region
Moemken et al. (2018)		–		–	W: – over southwest Russia, disagreement between RCMs on northwest Russia S: –	–	W: – no model agreement between RCMs on sign S: –

Notes:

+: increase, –: decrease, +/-: increase in some regions and decrease in others, n. s.: not statistically significant, n.c.: no change, W: winter, S: summer

For those studies not reporting a global warming level but only an RCP scenario–timeline combination, the latter was associated with the relevant global mean temperature increase.

Table SM13.7 | Magnitude of future change in onshore wind power potential under global warming levels (Figure 13.16)

	Onshore wind power potential (magnitude of change)			
Sub-region	1.5°C	2°C	≥3°C	Measurement unit in study
Northern Europe				
Davy et al. (2018)	−3 to 0%	−8 to 0%	−10 to 0%	Wind power density
Moemken et al. (2018)	−2 to 0%	−8 to 4% (W: −6 to 6%, S: −12 to 6%)	−8 to 6% (W: −4 to 14%, S: −20 to −4%)	Wind energy output
Devis et al. (2018)	4 to 8% (W: up to 7%, S: −7 to 6%)			Mean power output
Tobin et al. (2018a)	−2to 0%	−2 to −1%	−6 to −2.5%	Wind power production
Reyers et al. (2016)		0 to 1% (W: 0 to 3%, S: −3 to 0%)	1 to 4% (W: 0 to 8%, S: −7 to 0%)	Wind energy output
Carvalho et al. (2017b)	−15 to 0% (W: −15 to −5%, S: −15 to −5%)	−20 to 20% (W: −15 to −5%, S: −15 to 20%)	−20 to 30% (W: −15 to 0%, S: −30 to 15%)	Wind energy density
Tobin et al. (2016)	−2 to −1%	−3 to −1%		Annual energy yield
Després and Adamovic (2020)	0%	0%	−5 to −1%	Electricity production from wind
Western Central Europe				
Reyers et al. (2016)		−1 to 1%	−1 to 2%	Wind energy output
Moemken et al. (2018)	−2 to 0%	−4 to 0% (W: −4 to 12%, S: −14 to 0%)	−6 to 4% (W: −4 to 18%, S: −18 to 4%)	Wind energy output
Davy et al. (2018)	−3 to 0%	−3 to 0%	−5 to 2%	Wind power density
Carvalho et al. (2017b)	−30 to 0% (W: −40 to −5%, S: −30 to −5%)	−12 to 0% (W: −40 to −5%, S: −30 to −5%)	−12 to 0% (W: −40 to −5%, S: −30 to −5%)	Wind energy density

Sub-region	Onshore wind power potential (magnitude of change)			Measurement unit in study
	1.5°C	2°C	≥3°C	
Devis et al. (2018)	–3 to 5% (W: –5 to 5%, S: –8 to –5%)			Mean power output
Tobin et al. (2018a)	–2.5 to 0%	–4 to –1%	–4 to –1%	Wind power production
Tobin et al. (2016)	–0.5 to 1%	–3 to 2%		Annual energy yield
Després and Adamovic (2020)	0%	0%	0 to 2%	Electricity production from wind
Southern Europe				
Solaun and Cerdá (2020)	–8.2 to 5%	–8 to 6.5%		Production
Katopodis et al. (2019)	–15 to 8%	–5 to 8%		Wind potential
Reyers et al. (2016)		–2 to 0.8% (W: –2 to 0%, S: –6 to 1%)	–4 to 0% (W: –7 to –2%, S: –8 to 2%)	Wind energy output
Moemken et al. (2018)	–4 to 4%	–6 to 4% (W: –6 to 4%, S: –8 to 16%)	–14 to 12% (W: –16 to 16%, S: –12 to 18%)	Wind energy output
Davy et al. (2018)	–5 to 7%	–8 to 6%	–17 to 15%	Wind power density
Devis et al. (2018)	–12 to –2% (W: –12 to –6%, S: –8 to 6%)			Mean power output
Carvalho et al. (2017b)	–10 to –5% (W: –15 to –5%, S: –10 to –5%)	–10 to –5% (W: –15 to –5%, S: –20 to 20%)	–15 to 5% (W: –15 to –5%, S: –30 to 30%)	Wind energy density
Tobin et al. (2018a)	–4 to 1%	–4 to 1%	–8 to 3%	Wind power production
Tobin et al. (2016)	–2 to –1%	–6 to –4%		Annual energy yield
Després and Adamovic (2020)	0%	0%	0 to 2%	Electricity production from wind
Eastern Europe				
Devis et al. (2018)	–10 to 0% (W: –8 to 2%, S: –10 to 2%)			Mean power output
Carvalho et al. (2017b)	–30 to –5% (W: –20 to –10%, S: –30 to –5%)	–10 to 0% (W: –20 to –5%, S: –10 to –5%)	–20 to –5% (W: –20 to –5%, S: –20 to –5%)	Wind energy density
Davy et al. (2018)	–5 to –3%	–8 to –3%	–15 to –5%	Wind power density
Moemken et al. (2018)	–2 to 0%	–2 to 0% (W: –4 to 6%, S: –6 to 0%)	–4 to 0% (W: –4 to 8%, S: –12 to –4%)	Wind energy output

Notes:

In many studies, changes are provided in map format and thus figures in this table are our own interpretation of these map-based findings. Ranges illustrate variations within sub-regions. Values correspond to ensemble means, unless characterised otherwise.

Table SM13.8 | Sign of future change in offshore wind power potential under global warming levels (Figure 13.16)

Sub-region	Area in study	Offshore wind power potential (sign of change)					
		1.5°C		2°C		≥3°C	
Northern Europe							
Davy et al. (2018)	Europe, North Africa, Middle East	+/-	+ in a few parts of Baltic Sea, – in upper North Sea	+/-	+ in most of Baltic Sea, – in lower North Sea	+/-	+ in Baltic Sea, – in North Sea
Moemken et al. (2018)	Europe	+/-	+ in Baltic Sea, +/- in North Sea	+/-	+ in Baltic Sea, – in North Sea W: + S: + in Baltic Sea, – in North Sea	+/-	+ in Baltic Sea, – in North Sea W: +/- S: + in Baltic Sea, – in North Sea
Reyers et al. (2016)	Europe			+/-	+/- in Baltic Sea, +/- in the rest W: +, S: –	+/-	+/- in Baltic Sea, +/- in the rest W: +, S: –
Devis et al. (2018)	Europe	+	W: +, S: +				
Southern Europe							
Katopodis et al. (2019)	Greece	+/-	– in Ionian Sea and most of Aegean Sea	+/-	– in Ionian Sea and most of Aegean Sea		

Sub-region	Area in study	Offshore wind power potential (sign of change)					
		1.5°C		2°C		≥3°C	
Reyers et al. (2016)	Europe			+/-	+/- in Aegean Sea, – in the rest W: n.a. S: n.a.	–	– in almost all of Aegean Sea and in the rest W: n.a. S: n.a.
Koletsis et al. (2016)	Mediterranean and Black seas	+/-	+ in Aegean Sea, – in the rest			+/-	+ in Aegean Sea, – in the rest
Moemken et al. (2018)	Europe	+/-	+ in Aegean and Adriatic seas, – in the rest	+/-	+ in Aegean and Adriatic seas, – in the rest W: +/- S: + in Aegean and Adriatic seas, – in the rest	+/-	+ in most of Aegean Sea, – in the rest W: – S: + in Aegean and Adriatic seas, – in the rest
Devis et al. (2018)	Europe	–	– in most of Mediterranean Sea, n.c. in Aegean Sea W: – S: + in most of Aegean Sea, +/- in the rest				
Davy et al. (2018)	Europe, North Africa, Middle East	+/-	+ in most of Aegean Sea, – in the rest	+/-	+ in most of Aegean Sea, – in the rest	+/-	+ in most of Aegean Sea, – in the rest
Alvarez and Lorenzo (2019)	EU	+/-	+ in most of Aegean Sea, +/- in the rest	+/-	+ in most of Aegean Sea, – in most of the rest		

Notes:

+: increase, -: decrease, +/-: increase in some regions and decrease in others, n.c.: no change, W: winter, S: summer, n.a.: not available

For those studies not reporting a global warming level but only an RCP scenario–timeline combination, the latter was associated with the relevant global mean temperature increase.

Table SM13.9 | Magnitude of future change in offshore wind power potential under global warming levels (Figure 13.16)

	Offshore wind power potential (magnitude of change)			
Sub-region	1.5°C	2°C	≥3°C	Measurement unit in study
Northern Europe				
Davy et al. (2018)	−3 to 3%	−1 to 5%	−10 to 7%	Wind power density
Moemken et al. (2018)	−2 to 4%	−2 to 4% (W: −2 to 4%, S: 4 to 8%)	−4 to 4% (W: −4 to 4%, S: −8 to 6%)	Wind energy output
Reyers et al. (2016)		−1 to 0.5% (W: 0 to 2%, S: −2 to 0%)	−1 to 1% (W: 0 to 2%, S: −2 to 0%)	Wind energy output
Devis et al. (2018)	0 to 3% (W: 0 to 3%, S: 0 to 3%)			Mean power output
Southern Europe				
Katopodis et al. (2019)	−15 to 5%	−15 to 5%		Wind potential
Reyers et al. (2016)		−1 to 1% (W: 0 to 2%, S: −2 to 0%)	−1 to 1% (W: 0 to 2%, S: −2 to 0%)	Wind energy output
Moemken et al. (2018)	−2 to 4%	−4 to 2% (W: 0 to 4%, S: −4 to 8%)	−8 to 4% (W: −4 to 4%, S: −8 to 8%)	Wind energy output
Koletsis et al. (2016)	>5%		< −5% to >5%	Wind power
Devis et al. (2018)	−4 to 0%			Mean power output
Davy et al. (2018)	−6 to 6%	−10 to 6%	−15 to 12%	Wind power density

Notes:

In many studies, changes are provided in map format and thus figures in this table are our own interpretation of these map-based findings. Ranges illustrate variations within sub-regions. Values correspond to ensemble means, unless characterised otherwise.

Table SM13.10 | Sign and magnitude of future change in solar power potential under global warming levels (Figure 13.16)

Sub-region	Area in study	Solar power potential (magnitude of change)								
		1.5°C			2°C			≥3°C		
Northern Europe										
Tobin et al. (2018a)	EU and Switzerland	–		–4 to –1%	–		–6 to –2%	–		–8 to –3%
Jerez et al. (2015)	Europe				–	Higher decrease in northern Scandinavia	–10 to 0%	–	Higher decrease in northern Scandinavia	–20 to 0%
Gutiérrez et al. (2020)	Europe				+/–	– in southern Norway and northern UK in one RCM	–5 to 20%			
					+/–	+ in southern UK and southern Sweden	–20 to 5%			
Muller et al. (2019)	Europe	+/–	+ in southern UK and Denmark	–3 to 2%	+/–	+ in southern UK and Denmark	–6 to 2%			
Després and Adamovic (2020)	EU	n.c.			n.c.					
Western Central Europe										
Tobin et al. (2018a)	EU and Switzerland	–		–3 to 0%	–		–3 to –1%	–		–4 to –1%
Jerez et al. (2015)	Europe				–	Negligible changes over France	–10 to 0%	–	Uncertain changes over western France	–10 to 0%
Gutiérrez et al. (2020)	Europe				+	Two RCMs (including aerosols); in RACMOE22, increases to 15%	5 to 30%			
					+/–	Four RCMs + over France	–15 to 5%			
Muller et al. (2019)	Europe	+		1 to 2.5%	+		0 to 3%			
Després and Adamovic (2020)	EU	n.c.			n.c.					
Southern Europe										
Tobin et al. (2018a)	EU and Switzerland	–		–1 to 0%	–		–1 to 0%	–		–1 to 0%
Jerez et al. (2015)	Europe				–	Uncertain changes over most of the Iberian Peninsula	–5 to 0%	–	Negligible changes over most of the Iberian Peninsula	–5 to 0%
Gutiérrez et al. (2020)	Europe				+	Two RCMs (including aerosols)	5 to 25%			
					+/–	Four RCMs; + over most of the area	–5 to 5%			
Muller et al. (2019)	Europe	+		0.5 to 2.5%	+		1 to 3%			
Després and Adamovic (2020)	EU	n.c.			n.c.					
Eastern Europe										
Jerez et al. (2015)	Europe				–		–10 to –5%	–	Higher decrease in northwest Russia	–15 to –5%
Gutiérrez et al. (2020)	Europe				+	Two RCMs (including aerosols)	5 to 20%			
					–	Four RCMs	–15 to –5%			

Sub-region	Area in study	Solar power potential (magnitude of change)								
		1.5°C			2°C			≥3°C		
Muller et al. (2019)	Europe	–		–3 to 0%	–		–6 to –2%			
Després and Adamovic (2020)	EU	n.c.			n.c.			n.c.		

Notes:

+: increase, –: decrease, +/-: increase in some regions and decrease in others, n.c.: no change

(a) In many studies, changes are provided in map format and thus figures in this table are our own interpretation of these map-based findings. Ranges illustrate variations within sub-regions. Values correspond to ensemble means, unless characterised otherwise.

(b) Measurement units for magnitude of change are as follows:

Tobin et al. (2018): PV power production; Jerez et al. (2015): PV potential production; Gutiérrez et al. (2020): surface solar radiation (and hence PV potential production); Müller et al. (2019): yearly PV production; Després and Adamovic (2020): electricity production from PV.

(c) The sign and magnitude presented in the first line for Gutiérrez et al. (2020) corresponds only to the range values for the two RCMs which include the effect of aerosols (i.e., ALADIN53 and RACMO22E), while the results in the second line corresponds to the range values for the other four RCMs explored in this study.

Table SM13.11 | Sign and magnitude of future change in hydropower potential under global warming levels (Figure 13.16)

Sub-region	Area in study	Hydropower potential (magnitude of change)								
		1.5°C			2°C			≥3°C		
Northern Europe										
Tobin et al. (2018a)	EU and Switzerland	+	Lowest in UK and Ireland	3 to 17%	+	Lowest in UK and Ireland	3 to 15%	+	Lowest in UK and Ireland; some models predict >30%	5 to 21%
van Vliet et al. (2016a)	Global (including Europe)				+/-	5 to 20% in northern Scandinavia, ±5% in most of the rest	-20 to 20%	+/-	5–20% in most of Norway, Sweden, northern Finland and northern UK; -20 to -5% in very few areas	-20 to 20%
van Vliet et al. (2016c)	Global (including Europe)				+/-	High increase in northern Scandinavia	-10 to >15%			
Després and Adamovic (2020)	EU	+		0 to 4%	+		1 to 7%	+		2 to 13%
Western Central Europe										
Tobin et al. (2018a)	EU and Switzerland	+		3 to 9%	+/-	– only in France; highest + in Czech Republic and Poland	-1 to 15%	+/-	– only in France; highest + in Czech Republic and Poland	-5 to 12%
van Vliet et al. (2016a)	Global (including Europe)				+/-	-5 to 5% only in Germany; -40 to -20% in Bulgaria, Romania and parts of Ukraine	-40 to 5%	+/-	-5 to 5% only in coastal Germany and Poland; -60 to -20% in most areas	-60 to 5%
van Vliet et al. (2016c)	Global (including Europe)				-		-15 to -5%			
Anghileri et al. (2018)	Alps	-		-27%						
Bombelli et al. (2019)	Italian Alps	+	-9.16% (CCSM4), 1.85% (EC-EARTH), 14.65% (ECHAM6), → 2.5% (mean)			+	0.03% (CCSM4), 10.72% (EC-EARTH), 8.33% (ECHAM6), → 6.4% (mean)		-	-6.3% (CCSM4), 0.23% (EC-EARTH), -7.23% (ECHAM6), → -4.4% (mean)
Patro et al. (2018)	Italian Alps	+/-	+ when glacier area is <10% of total basin area	-32 to 5%	+/-	+ when glacier area is <10% of total basin area	-23 to 1%	+/-	+ when glacier area is <10% of total basin area	-40 to 6%

Sub-region	Area in study	Hydropower potential (magnitude of change)							
		1.5°C			2°C			≥3°C	
Stocchi et al. (2019)	Italian Alps	–	–25% (CCSM4), –14% (EC-EARTH), –10% (ECHAM6), → –16.3% (mean)		–	–28% (CCSM4), –10% (EC-EARTH), –13% (ECHAM6), → –17% (mean)		–	–27% (CCSM4), –23% (EC-EARTH), –33% (ECHAM6), → –28% (mean)
Adynkiewicz-Piragas and Miszuk (2020)	Germany, Poland				–		–10 to 0%	–	–34 to –16%
Després and Adamovic (2020)	EU	+		0 to 1% (median values)	+		2 to 4% (median values)	+	1% (median value)
Southern Europe									
Tobin et al. (2018a)	EU and Switzerland	+/-	+ only in Italy and Slovenia; decrease >10% in Greece, Spain and Portugal in some models	–7 to 5%	+/-	+ only in Italy and Slovenia	–10 to 2%	+/-	+ only in Slovenia; decrease >20% in Greece, Spain and Portugal in some models
van Vliet et al. (2016a)	Global (including Europe)				–	–20 to –5% only in Italy	–40 to –5%	–	–60 to –40%
van Vliet et al. (2016c)	Global (including Europe)				–		< –15 to –10%		
Lobanova et al. (2016)	Portugal (Tagus)	–		–50 to –10%	–		–50 to –10%	–	–60 to –40%
Solaun and Cerdá (2017)	Spain	–		–25 to –5%	–		–30 to –10%	–	–49 to –30%
Després and Adamovic (2020)	EU	–		–1% (median value)	–		–3% (median value)	–	–2% (median value); may reach –13% under extreme events)
Eastern Europe									
van Vliet et al. (2016a)	Global (including Europe)				+/-	>5% in northern Russia, < –5% in western Russia	–20 to 20%	+/-	–5 to 5% in most of the areas
van Vliet et al. (2016c)	Global (including Europe)				+/-	+ in northern Russia	–10 to 15%		
Akentieva et al. (2014)	Russia	+/-		–8 to 14%	+/-		–10 to 18%		

Notes:

+: increase, –: decrease, +/-: increase in some regions and decrease in others, n.c.: no change

(a) In many studies, changes are provided in map format and thus figures in this table are our own interpretation of these map-based findings. Ranges illustrate variations within sub-regions. Values correspond to ensemble means, unless characterised otherwise.

(b) Measurement units for magnitude of change are as follows:

Tobin et al. (2018): hydropower production; van Vliet et al. (2016a): gross hydropower potential; van Vliet et al. (2016b): annual mean usable capacity of current hydropower plants; Després and Adamovic (2020): electricity production from existing hydropower plants; Anghileri et al. (2018): electricity production; Bombelli et al. (2019): yearly average energy production; Patro et al. (2018): water volume used for energy production in hydropower plants; Stocchi et al. (2019): energy production; Adynkiewicz-Piragas and Miszuk (2020): energy production; Després and Adamovic (2020): electricity production; Lobanova et al. (2016): hydropower produced; Akentieva et al. (2014): average annual electricity production.

Table SM13.12 | Sign and magnitude of future change in bioenergy crops (rapeseed) potential under global warming levels (Figure 13.16)

Sub-region	Area in study	Bioenergy potential (magnitude of change)				
		1.5°C		2°C	≥3°C	
Western Central Europe						
Jaime et al. (2018)	Global (including Europe)	–	EU countries: –8.5% (CSIRO Mk2), –9% (Hadley CM3), –10.2% (MPI ECHAM4), → –9.2% (mean)		–	EU countries: –7.6% (CSIRO Mk2), –22.2% (Hadley CM3), –24.9% (MPI ECHAM4), → –18.2% (mean)
Cronin et al. (2020)	Global (including Europe)				+	4% (no land restrictions both in past and future)
						27% (SSP5, where urban, food agricultural land and protected areas are excluded both in past and future)
						1% (SSP5, but forest areas are not excluded both in past and future)
Eastern Europe						
Jaime et al. (2018)	Global (including Europe)	+	Rest of Europe (mainly Russia): 32.5% (CSIRO Mk2), 41.6% (Hadley CM3), 54.4% (MPI ECHAM4), → 42.8% (mean)		+	Rest of Europe (mainly Russia): 15.8% (CSIRO Mk2), 44.2% (Hadley CM3), 58.8% (MPI ECHAM4), → 39.6% (mean)
Cronin et al. (2020)	Global (including Europe)				+	Russia (whole): 210% (no land restrictions both in past and future)
						Russia (whole): 219% (SSP5, where urban, food agricultural land and protected areas are excluded both in past and future)
						Russia (whole): 431% (SSP5, but forest areas are not excluded both in past and future)

Notes:

+/: increase, –/: decrease, +/–/: increase in some regions and decrease in others, n.c.: no change

(a) Measurement units for magnitude of change are as follows: Jaime et al. (2018): percentage of change of land suitable for rapeseed cultivation (to produce biofuels) compared with 1996; Cronin et al. (2020): percentage of change of total land that is moderately or highly suitable for energy crops cultivation (15 crops) compared with 1980–2009.

(b) Jaime et al. (2018) present results for EU countries as a total. According to the study, most suitable land in the EU is, and will remain, in Western Central Europe, and therefore the whole percentage change in the EU has been allocated to Western Central Europe. Note that that ‘Eastern Europe’ in the study corresponds to ‘Western Central Europe’ in the present IPCC Assessment Report.

(c) The study also presents results for the whole suitable area for rapeseed cultivation, and thus by subtracting the EU figures, the change in the rest of Europe and Western Asia can be estimated. The figures for Eastern Europe correspond to the total change in the rest of Europe and Western Asia and thus are overestimated.

(d) Cronin et al. (2020) present results for different sub-regions of Europe which, however, do not match the definitions adopted in this IPCC chapter; therefore, the reference’s results—calculated from the data provided on the reference’s supplementary material—for Russia are presented under Eastern Europe, while EEU, FSU and WEU are presented as an aggregated figure under Western Central Europe.

Table SM13.13 | Sign and magnitude of future change in thermoelectric power capacity under global warming levels (Figure 13.16)

Sub-region	Area in study	Thermoelectric power capacity (magnitude of change)								
		1.5°C			2°C			≥3°C		
Northern Europe										
Tobin et al. (2018a)	EU and Switzerland	–		–7 to –2%	–		–10 to –3%	–		–15 to –7%
Després and Adamovic (2020)	EU	–		–4 to 0%	–		–6 to 0%	–		–15 to 0%
van Vliet et al. (2016c)	Global (including Europe)				–		<–15%			
Byers et al. (2020)	UK				–	Cumulative impacts over 32 plants	–18% (p5), –20% (p50), –45% (p95)	–	Cumulative impacts over 32 plants	–36% (p5), –41% (p50), –58% (p95)

Sub-region	Area in study	Thermoelectric power capacity (magnitude of change)								
		1.5°C			2°C			≥3°C		
Western Central Europe										
Tobin et al. (2018a)	EU and Switzerland	–		–8 to –5%	–		–13 to –6%	–		–18 to –11%
Després and Adamovic (2020)	EU	n.c.		0%	–		~0%	–	– for nuclear plants	–2%
van Vliet et al. (2016c)	Global (including Europe)				–		<–15%			
Southern Europe										
Tobin et al. (2018a)	EU and Switzerland	–		–7 to –1%	–	All countries except Portugal <–8%	–12 to –1%	–	All countries except Portugal <–14%	–18 to –1%
Després and Adamovic (2020)	EU	–	– for nuclear plants; figure results from impacts on whole energy system	–2 to 0%	–	– for nuclear plants; figure results from impacts on whole energy system; –13% under extreme events	–5 to –1% (and may reach –12% under extreme events)	–	– for nuclear plants; figure results from impacts on whole energy system; –12% under extreme events	–8 to –2% (and may reach –12% under extreme events)
van Vliet et al. (2016c)	Global (including Europe)				–		<–15%			
Payet-Burin et al. (2018)	Iberian Peninsula	–	Average for all freshwater-cooled plants: –18 to –6% (annual) W: –12 to –1%, S: –30 to –16%							
			Once-through cooling: –16 to –11% (annual) S: –40 to –31% closed-circuit cooling: <–0.5%							
Eastern Europe										
van Vliet et al. (2016c)	Global (including Europe)				–		–10 to –5%			
Klimenko et al. (2018)	Russia				–	Efficiency reduction by 0.2–0.6% per +1°C	–1.2 to –0.4% (if a +2°C is considered)	–	Efficiency reduction by 0.2–0.6% per +1°C	Up to –2% (if a +3°C is considered)

Notes:

+: increase, –: decrease, +/-: increase in some regions and decrease in others, n.c.: no change, S: summer, W: winter, p5: 5th percentile, p50: 50th percentile (median), p95: 95th percentile.

(a) In many studies, changes are provided in map format and thus figures in this table are our own interpretation of these map-based findings. Ranges illustrate variations within sub-regions. Values correspond to ensemble means, unless characterised otherwise.

(b) Measurement units for magnitude of change are as follows:

Tobin et al. (2018): thermoelectric power production; van Vliet et al. (2016b): annual mean usable capacity of thermoelectric power plants; Després and Adamovic (2020): electricity production from existing thermoelectric plants (including nuclear); Anghileri et al. (2018): electricity production; Bombelli et al. (2019): yearly average energy production; Patro et al. (2018): water volume used for energy production in hydropower plants; Stucchi et al. (2019): energy production; Adynkiewicz-Piragas and Miszuk (2020): energy production; Després and Adamovic (2020): electricity production; Lobanova et al. (2016): hydropower produced; Akentieva et al. (2014): average annual electricity production; Payet-Burin et al. (2018): available power plant capacity of freshwater-cooled thermal power plants; Byers et al. (2020): available power plant capacity at 99th percentile extreme day; Klimenko et al. (2018): power plant efficiency.

Table SM13.14 | Examples of adaptation options for reducing sectoral climate-change risks (Figure 13.19)

Sector	Risk and cause	Examples of adaptation options	Elements related to the implementation of adaptation options
Thermoelectric power	Reduction/interruption of operation due to water cooling constraints	Replacement of once-through cooling by cooling towers; dry air-cooling; seawater cooling (for coastal plants); replacement by new, better-adapted plants (Gasbarro et al., 2016; Behrens et al., 2017; EEA, 2019a; IAEA, 2019)	The choice of electricity producers between options is guided mainly by the economics of adaptation technologies, and less by the supply of information on future climate change (Bogmans et al., 2017). Costs for retrofitting cooling are site specific and increase with the distance to water bodies, needs for additional structures and the plant's age (Sieber, 2013). Dry air-cooling for new plants is 3–4 times higher than wet recirculating system and 4–5.5 times higher than once-through cooling (IAEA, 2019), and could result in 10% efficiency losses (EEA, 2019a).
		Switching to alternative-generation technologies with low water use, (e.g., wind, solar PV) (Gasbarro et al., 2016; Porfiriev et al., 2017; EEA, 2019a)	Fragmentation of policy frameworks for energy and water make cohesive energy and water management difficult (Byers et al., 2015; Behrens et al., 2017). Ignoring the impact of climate-induced water constraints may significantly increase the energy system costs (Khan et al., 2016).
		Inter-basin water transfer (Koch et al., 2014)	
		Use of 'non-traditional' water sources (e.g., recirculation of water from oil and gas fields or coal mines, treated wastewater from nearby cities, desalination, water reuse) (Sieber, 2013; Gasbarro et al., 2016)	Nearby sources of 'non-traditional' waters may not exist or may be of insufficient capacity.
		Shift part of power production to estuaries or coasts (Byers et al., 2015)	Its implementation to high-demand areas could bring significant nationwide water reductions (Byers et al., 2015), provided that recipient plants can undertake the extra production.
		Improved electricity interconnections (Behrens et al., 2017)	There are high investment costs.
		Demand management measures (e.g., smart grid/meters, pricing options, energy efficiency) to reduce the economic losses during power curtailment (Gasbarro et al., 2016)	This can be an effective option, particularly under low climate-change scenarios (Hanski et al., 2018).
		Institutional measures (e.g., water temperature cap, heat-load plan, contract between environmental regulator and electricity producers)	The efficiency of institutional adaptation options may differ depending on the increase in heatwave intensity, frequency or both (Eisenack, 2016). Stakeholder engagement (e.g., increase awareness of decreasing water flows, cooperation) to prevent the conflicts with the local communities in case of water curtailments is helpful (Gasbarro et al., 2016)
Hydropower	Reduced production due to lower streamflow	Adjusted hydropower management (Gaudard et al., 2013; EEA, 2019a)	Optimising the hydraulic head and the turbine schedule with respect to the prices could reduce power losses by up to 35% (Gaudard et al., 2013).
	Increased risk of damage from flooding	Adjusted hydropower management (Aparicio, 2017; Ranzani et al., 2018)	Adaptive strategies in the management of reservoirs could reduce (but not prevent) revenue losses (Ranzani et al., 2018).
		Recalibration of spillways (e.g., through piano-key wires, concrete or metal fuse gates)	Recalibration systems have been implemented successfully in hydropower facilities in Europe and outside Europe (EEA, 2019a).
		Increase the capacity of existing hydropower plants (i.e. increase installed turbine capacity, increase reservoir storage)	This option has been implemented in some northern European hydropower plants (EEA, 2016a).
		Hydropower operational warning systems, monitoring of snowpack and river flows, forecast of high water flows	Hydropower forecasting faces key challenges related to integration of state-of-the-art weather services, data assimilation schemes, links between forecast quality and value, and enhancement of risk-based decision making (Boucher and Ramos, 2018).

Sector	Risk and cause	Examples of adaptation options	Elements related to the implementation of adaptation options
Electricity transmission and distribution	Power outages due to damage to transmission lines and power stations from extreme winds, storm surges, floods and very high temperatures	Construction of new substations and overhead lines (providing additional paths to transfer power in case of a transmission line failure); improve old overhead lines and substations (Gasbarro et al., 2016; Fu et al., 2018)	Efficiency increases when construction of new lines and stations occurs to decentralised power systems, while improvement of existing lines and stations is less efficient (Fu et al., 2018). The contribution of substations' refurbishment in building flood resilience depends on the degree of protection of critical substations which may not necessarily be the most vulnerable to high water levels (Bollinger and Dijkema, 2016). The willingness to pay (WTP) to avoid power outage is higher for older people, women and urban residents; risk perceptions are greatly influenced by current regional temperatures; and under strong warming the WTP increases in summer and decreases in winter in all countries, particularly in the north (Cohen et al., 2018a). Experience with previous power cuts significantly enables resilience (Ghanem et al., 2016; Cohen et al., 2018a).
		Vegetation management	New technologies (e.g., Lidar) allow to reduce the need for time-consuming and labour-intensive traditional verification and to provide reliable input for guiding tree trimming; however, their implementation may require significant changes of management practices at the utility level.
		Turning, partially or totally, of overhead lines into underground cables (Wang et al., 2016; Ciasca et al., 2017)	There is high installation cost and potentially long implementation time, depending on the area covered and the length of cables (EEA, 2019a). Selective undergrounding of line sections exposed to higher risk or harder to access can be a cost-effective adaptation strategy (Wang et al., 2016). For new lines, there are potentially long permit processes and public opposition (Wang et al., 2016; Ciasca et al., 2017).
		Locating assets above flood level, flood barriers, relocation of assets (Bollinger and Dijkema, 2016; Wang et al., 2016; Thacker et al., 2018)	There are high investment costs for existing assets (Wang et al., 2016) and long implementation time (EEA, 2020c). transmission system operators often prefer to combine investments in flood defences with major renovations or refurbishments to substations, and consequently prioritisation often occurs based on factors (e.g., its age) unrelated to a substation's criticality (Bollinger and Dijkema, 2016; Wang et al., 2016). Assets' relocation is almost always not cost beneficial (Thacker et al., 2018).
		Distribution circuit segregation and automation (Wang et al., 2016).	
		Increase in the height of poles supporting power lines, installation of conductors with hotter operating limits, use of 'low-sag' conductors (EEA, 2019a)	There are legal requirements on minimum pole-height-support adaptation (EEA, 2019a).
	Reliability problems in electricity networks due to increased peak load for cooling	Strict efficiency standards for cooling equipment (EEA, 2019a; Palkowski et al., 2019)	Reliability can be increased through measures reducing cooling demand, such as improved building design, water cooling technologies for thermoelectric generation that do not use electricity (e.g., heat-driven absorption cooling), or direct utilisation of cooling water where available (EEA, 2019a).
		Increase in transmission capacity, including international linkages	
		Increase in backup capacity	
Transport	Reduction/interruption of transportation due to damaged infrastructure and/or traffic disruption as a result of intense rain, flooding and heatwaves	Broad range of options (EEA, 2014; Frolov et al., 2014; Burbidge, 2015; Stamos et al., 2015; van Slobbe et al., 2016; Bachner, 2017):	Particularly in road transport, measures have also economy-wide feedback effects which must be considered when assessing adaptation benefits (Bachner, 2017). Nevertheless, as it is difficult to quantify the benefits and costs of adaptation measures in transport, cost-benefit analyses need to be performed on a case study level (Doll et al., 2014).
		(a) Infrastructure construction/retrofitting (e.g., enlargement of drainage systems, measures to reduce slippery roads, raise links above flood level)	As 'soft' adaptation options (e.g., ICT) have already been implemented to a large extent in railways, investments in advanced protection systems (e.g., tunnels, protection walls and enlarged drainage) are necessary to support proactive maintenance strategies (Doll et al., 2014). Improving drainage or elevating critical road links can be cost-effective but requires analysis at the city level (Pregolato et al., 2017). Adaptations in vessel design may reduce the vulnerability to low depth, but with a trade-off with performance in times of sufficient discharge (van Slobbe et al., 2016).
		(b) Improved maintenance (e.g., vegetation management, visual road inspection)	Network maintenance can be a more cost-efficient way to reduce short- and medium-term damage risks (Doll et al., 2014).
		(c) ICT and users (e.g., early warning on adverse weather, weather forecasting)	Lack of coherence between the climate adaptation plans of companies operating major transport infrastructure and their neighbouring municipalities may reduce the effectiveness of adaptation actions undertaken by the transport sector (EEA, 2014).
		(d) Modal shifts	
		(e) Technological innovations (e.g., heat-resistant pavement materials, materials designed for a greater number of cycles of freezing and thawing, logistic chains)	Some of the new pavement materials may increase noise levels in urban areas (Enriquez-de-Salamanca (2019).

Sector	Risk and cause	Examples of adaptation options	Elements related to the implementation of adaptation options
Transport	Reduction/ interruption of transportation due to damaged infrastructure and/or traffic disruption as a result of intense rain, flooding and heatwaves	(f) Revising operational guidelines and standards	The location-specific nature of weather impacts requires analysis and response also at the route level to ensure that investments in flood protection are cost-effective (EEA, 2020c). Dynamic heat management can reduce the heat-related disruption from unnecessary emergency speed restrictions on railway networks (Ferranti et al., 2016).
	Reduction of thermal comfort of passengers in railways and metro lines due to higher temperatures	Saloon cooling, cooling of platforms and tunnels (Jenkins et al., 2014a)	Saloon cooling alone may not be sufficient to maintain comfortable thermal conditions for some lines under high emission scenarios (Jenkins et al., 2014b).
Winter tourism	Reduction/ interruption of operation due to lack of snow	Snowmaking (including application of automated snowmaking systems)	High investment cost (i.e., for development of water supply systems, purchase and installation of snow cannons) and increased operational costs (Campos Rodrigues et al., 2018; Scott et al., 2019) are potential barriers. Increased snowmaking can maintain snow reliability under low warming but is not sufficient under high-end warming (Steiger and Scott, 2020). Snowmaking was found to reduce guest loyalty of skiing destinations in some customer segments by affecting the natural scenery and raising prices (Bausch et al., 2019).
		Protection and conservation of snowpack (e.g., water drainage, modification of the ski-run slopes, protection from avalanches, protection or storage of snow during the non-ski seasons); snow farming (Steiger and Scott, 2020)	Several techniques are available.
		Expansion of skiable area	Need of substantial investments and free areas has adverse impacts (e.g., land-use conflicts, impacts of construction on natural areas, impacts on the landscape quality, increased water and energy use) (Campos Rodrigues et al., 2018).
		Nocturnal skiing (Campos Rodrigues et al., 2018)	Nocturnal skiing is already offered at some ski resorts, but it can compensate for a small part of potential losses due to adverse weather and safety limitations (Campos Rodrigues et al., 2018).
		Shift to other ski destinations (spatial substitution)	The attachment ('loyalty') of glacier skiers to their favourite leisure destination, gender, demographics (i.e., age) and perceptions towards environmental sustainability were found to be important in guiding adaptation preferences (Demiroglu et al., 2018).
		Diversification of snow-based activities	Diversification may be too costly for some resorts. Skiing in indoor slopes may be the least preferred option for skiers, while other winter activities (e.g., snowboarding, downhill skiing) may not be effective adaptation options as skiing is perceived as a necessity in some countries (Falk, 2015; Falk and Scaglione, 2018).
		Transformation to multi-recreational mountain resorts, compensating non-snow activities	Transformation can take place by diversifying the offer in winter towards nature-based activities or place-bound products beyond winter sports (Bausch and Gartner, 2020) and by developing year-round tourism and snow-independent tourism products (Steiger and Scott, 2020). Transformation may be too costly for some resorts (Campos Rodrigues et al., 2018). Non-snow activities are not appealing to people for whom skiing is the main activity of their winter holiday (Steiger and Scott, 2020). Cultural differences affect the effectiveness of compensating activities (Landauer et al., 2013).
		Management options (e.g., grouping of resorts, pricing strategies) (Campos Rodrigues et al., 2018)	Price discounts are effective under less severe warming scenarios (Steiger et al., 2020). Price discounts in ski lift tickets may not be enough to attract foreign visitors (Falk and Scaglione, 2018).

Sector	Risk and cause	Examples of adaptation options	Elements related to the implementation of adaptation options
Coastal and summer tourism; other forms of tourism	Loss of beaches/coasts due to SLR and increased erosion	Hard defences (e.g., seawalls)	It is a measure that has been widely applied in Europe, but generally with no concern for future climate-change impacts. It also requires high investments, proper maintenance (which is costly) and can affect sediment transport and coastal erosion (Pranzini et al., 2015).
		Soft measures (e.g., artificial beach nourishment, dune planting) (Pranzini et al., 2015; Jiménez et al., 2017)	Selective sand nourishment is common in Europe (Pranzini et al., 2015), as in Spain where more than 22 million m ³ of sand have been deposited on the Catalan coast during the past 30 years (Jiménez and Valdemoro, 2019). There is often reduced availability and there are high costs of fill material. Potential governance difficulties also exist (e.g., lack of well-defined roles in coordinating nearshore activities, division of costs between government, private owners and local communities).
		Inland shift of tourism activities	This may not be possible due to land-use and financial constraints, as well as environmental and administrative regulations.
	Disruption of tourism activities due to higher temperatures	Temporal shift/extension of recreational activities outside the summer period (e.g., transition-time tourism, year-round tourism)	Potential barriers include organisational issues and the inability or reluctance of clientele (Mourey et al., 2020). Limiting factors to be considered include cost, school holidays or work (Brosy et al., 2014).
	Disruption of tourism activities due to water scarcity	Water supply and demand measures (e.g., desalination, rainwater harvesting, water remediation and water reuse, water saving devices) (Michailidou et al., 2016)	Identification of the suitable alternatives implies consideration of several and often conflicting criteria and stakeholder involvement (Michailidou et al., 2016).
Business	Reduction/interruption of operation due to extreme events; asset damage; reduced sales due to lower demand; health and safety risks for staff	Individual adaptation measures at the enterprise and cluster levels: general measures (e.g., risk assessment and monitoring, technical solutions, reduction of exposure through geographical decisions, shifting and sharing of risks, disaster relief and business continuity, portfolio diversification, and cooperation) (Pinkse and Gasbarro (2019)	Potential barriers include a lack of human or financial resources and scientific or technical knowledge to understand climate-change risks (Aguinaldo et al., 2019); and also limited knowledge on scale of assessment, evidence base, adaptation response, scope of impacts, interdependencies and public policy (Surminski et al., 2018). The cluster approach allows to overcome the lack of resources and knowledge to address climate risks and adaptation options, which characterise particularly small and medium-size enterprises (Aguinaldo et al., 2019).
		Individual adaptation at the enterprise and cluster levels: measures against heatwaves (e.g., shielding or reflective surfaces, use of 'cool' materials, green walls, green parking and draining floors, energy management system, changes in working practices, appropriate clothing) (Ciscar et al., 2018; IRIS LIFE project, 2019)	Changes in working practices for outdoor workers: more frequent and longer breaks during the hottest parts of the day; earlier starts and/or later ends to the working day; and night working. Other changes: provide drinking water; scheduling heavy work during the cooler parts of the day or reducing work during the hottest part of the day; alternate work and rest periods; wearing appropriate clothing; and employees' education. Shifting the working hours from daytime to nighttime could entail side effects such as chronic fatigue, anxiety, depression and noise pollution for nearby residents (Ciscar et al., 2018).
		Individual adaptation at the enterprise and cluster levels: measures against storms	Potential measures include closing tunnel connection in case of strong wind, tree pruning, limiting storage of materials outside the building, shielding surfaces and roof anchorage (IRIS LIFE project, 2019).
		Individual adaptation at the enterprise and cluster levels: measures against heavy rains and floods	Potential measures include alert systems, anti-reflux valves, temporary meteoric water storage areas, anti-flooding bulkheads and storage of materials at a safe height from the ground (IRIS LIFE project, 2019).
		Individual adaptation at the enterprise and cluster levels: measures against drought	Drought causes reduced water availability for the processes within the factories. Potential measures include treatment and reuse of wastewater as well as sector-specific measures for reducing water consumption (Alkayal et al., 2015). In food and beverage, options include implementation of Best Available Techniques, sector-specific measures for meat and poultry, fish, fruits and vegetables, dairy products and drink processing sectors (Valta et al., 2016).
		Corporate adaptation	Corporate strategies to climate risks still predominantly focus on mitigation (Sakhel (2017; Pinkse and Gasbarro, 2019). It is dominated by surveillance of climatic changes, climate-proofing production facilities and assets, and supply chain management (Sakhel (2017).
Banking and finance	Risk of instability of the financial system due to damage caused by climate extremes	Extension of financial regulations and requirements for risk monitoring towards climate-related risks, such as climate-related stress tests (TCFD, 2017; D'Orazio and Popoyan, 2019)	Only in a few European countries are such regulations in place, but they are voluntary (D'Orazio and Popoyan, 2019).
Insurance	Risk of insurance default	Public reinsurance	
		Extension of risk monitoring towards climate-related risks	

Sector	Risk and cause	Examples of adaptation options	Elements related to the implementation of adaptation options
Cities	Reduced indoor and outdoor thermal comfort and power outages due to heatwaves ^a	Passive and active cooling measures in buildings (e.g., air conditioning, ventilation, shading)	Coping appraisal is a strong predictor for citizens' motivation to adapt, while elderly people are less motivated (Murtagh et al., 2019). Though necessary, natural ventilation alone cannot fully mitigate overheating (Dodoo and Gustavsson, 2016; Heracleous and Michael, 2018; Dino and Meral Akgül, 2019; Shen et al., 2020), and its effectiveness is strongly affected by occupants' behaviour (Tillson et al., 2013; van Hooff et al., 2014), while a combination of shading and ventilation (including nighttime) is needed (Ibrahim and Pelsmakers, 2018). Installation of air conditioning may be too costly for many households (Thomson et al., 2019). Large increase of air conditioning in densely populated areas may exacerbate the urban heat island and thus overheating (Kingsborough et al., 2017), and increases pressure on electricity systems.
		Interventions at the buildings' shell (e.g., improving insulation, increasing thermal mass, use of phase-change materials)	Addition of insulation in poorly ventilated and shaded buildings may increase overheating. Altering the thermal mass is much harder in older buildings (Tillson et al., 2013). Phase-change materials could significantly reduce the cooling load but is a relatively new technology to the construction industry, with many uncertainties (e.g., future prices, long-term durability, energy cost) (Sajjadian et al., 2015). Material changes to buildings are often prohibited by restrictive tenancy relations (Thomson et al., 2019).
		Green, blue and grey infrastructure (e.g., green areas, green roofs and walls, cool roofs and façades, cool pavements)	Climatic conditions may affect performance of options (Ward et al., 2016). On-site water reuse systems can provide supplementary water to green roofs and walls, gardens and other smaller-scale urban nature-based solutions on an as-needed basis. The cooling potential of plants in green roofs or walls is influenced by the choice of plant species (Cameron et al., 2014). Cool roofs are less expensive and easier to apply than green roofs (Carvalho et al., 2017a). People's willingness to pay for green infrastructure (GI) has been found to be mostly related to income and ethnicity, while citizens are willing to support climate adaptation through GI as long as the GI is multifunctional (i.e., comes with recreational and aesthetic benefits) (Derkzen et al., 2017). Urban governance mechanisms and institutional barriers to GI planning need additional research (Emmanuel and Loconsole, 2015). Cool roofs are an established technology, but this is not yet the case for cool pavement materials which may cause glare problems or excessive illumination levels (Carnielo and Zinzi (2013). Options are not easily transferable between countries or even cities (Hintz et al., 2018).
		Update building standards to consider the expected increase of extreme summer temperatures and the consequent increase of energy demand for cooling	Standards considering climate change and outdoor climate conditions, and realistic assumptions in terms of occupant adaptations, are needed (Mulville and Stravoravdis, 2016; Sánchez-García et al., 2020; Shen et al., 2020). Standards should consider regional differences (Frolov et al., 2014).
		Watering of roads and pavements	Emergency options are needed during heatwaves, but not a long-term adaptation option (Hendel et al., 2017; Enriquez-de-Salamanca, 2019). Optimisation of possible watering methods has only rarely been conducted, while water consumption is an issue (Hendel et al., 2015).
		Escape to nearby mountainous regions (Juschten et al. (2019b)	Mostly motivated by social and subjective norms, past experience with heat stress, outdoor sports as a travel motive, previous visits to the destination, positive media coverage and perceived behaviour control (Juschten et al., 2019a).
	Reduced water supply due to drought	Water demand management (Buurman et al., 2017)	Voluntary or enforced water conservation is a potential measure. There is a high degree of uncertainty regarding effectiveness, but when coupled with new water reuse infrastructure, it could keep the probability of exceeding the target frequency of an emergency drought order below 0.01 in London under severe drought by 2100 for a medium-emissions scenario and high population growth (Kingsborough et al., 2016).
		Expansion of water supply (Buurman et al., 2017)	Water reuse, new water reservoirs, inter-basin transfers, desalination and artificial aquifers are potential measures. This is less flexible as it requires commitment to supply infrastructure, which can be maladaptive under increasing water demand (Kingsborough et al., 2016).
	Corrosion of buildings due to permafrost and thaw melting	Use of materials with proper resistance to freezing–thawing cycles (Frolov et al., 2014)	Development of an assessment methodology and database on the durability of materials under various climatic conditions is needed to support the selection of optimal materials under the future climate. Regular updating of regulatory parameters based on observational data is also needed (Frolov et al., 2014).
		Increase in corrosion resistance of structural elements (Frolov et al., 2014)	
		Design solutions that prohibit an increase in moisture content in building structures (Frolov et al., 2014)	

Sector	Risk and cause	Examples of adaptation options	Elements related to the implementation of adaptation options
Cities	Damage to settlements and infrastructure due to flooding	Building new flood defences (e.g., dikes)	Costs for maintaining the baseline flood protection level under climate change (and introducing a minimum of 100 years of protection) through defences would not outweigh benefits for many countries before 2030 under different RCP or SSP scenarios, while in 2080 adaptation benefits would exceed costs in almost all countries (Bouwer et al., 2018). Update and increased maintenance of storm barriers comes along with concerns on their environmental impacts as in St Petersburg (Rodionov, 2016). Decision making under deep-uncertainty approaches can be applied to deal with the uncertainties of climatic projections (Babovic et al., 2018). Information such as a high-resolution European and Mediterranean flash floods dataset can provide input to impact assessment, modelling and forecast (Amponsah et al., 2018).
		Heightening and/or strengthening of existing dikes, dams and levees; widening of river floodplains and reduction of obstructions in floodplains (Bouwer et al., 2018); updating the urban drainage system (Bodoque et al., 2019)	
		Flood protection measures at the building and household level, such as dry-proofing (e.g., sealing walls with waterproof coatings, impermeable layering of masonry, sealants for openings), wet-proofing (e.g., building elevation, use of water-resistant materials), emergency measures (e.g., mobile flood barriers, sandbags), securement of sources of contamination	There is <i>high confidence</i> that past experience of damage strongly affects risk perception and hence motivation for adaptation (Baron and Petersen, 2015; Lujala et al., 2015; Osberghaus, 2015; Madsen et al., 2019); however, protection may be motivated mostly by coping and threat appraisal as well as trust in public institutions (Bamberg et al., 2017). Dry-proofing is costly and thus usually applied to new buildings (Bouwer et al., 2018). The level of wet-proofing differs significantly between locations (Koerth et al., 2013; Stojanov et al., 2015). Perceptions of flood risks, expected climate impacts, risk attitudes and geographical characteristics were found to be the most important determinants in the decision to invest in elevating houses (Botzen et al., 2013).
		Nature-based Solutions (NbS) to manage water runoff (e.g., multifunctional green spaces, wetlands, retention/detention and infiltration basins, rain gardens and green roofs)	Natural ecosystems remain under threat from changing climatic conditions. Potential barriers for development of green infrastructure for flood risk management include coordinating and convincing stakeholders, limitations of the existing legislations and difficulty in assessing non-monetary benefits (Liu and Jensen, 2018).
		Facilitation of recovery after climate extremes	The location and composition of urban green spaces is key for effective adaptation (García Sánchez et al., 2018).
		Increase in flood risk standards, land-use planning, risk zoning, dedicated flood management legislation	Measures that work well in one region may not be effective in another region facing different flood hazards, and thus building codes and all other flood risk management policies have to be region specific (Poussin et al., 2015). Legislation to delimit non-suitable land for urbanisation often shows a slow implementation (Pérez-Morales et al., 2018), although it can be very effective in reducing risks under climate change (Thieken et al., 2016). Under high SLR, risk zoning can be more effective than hard defences (Andersson-Sköld et al., 2015).
		Emergency plans, training for evacuations, early warning systems	Require that the role and responsibilities of the different administrative departments and organisations involved be well defined, and that there be a clear plan on how to manage the different stages in the recovery process (Adedeji et al., 2019).
		Planned relocation	Higher resistance was found among seafront residents, second-home occupants, homeowners, elderly and retired people, and multi-generation households (Dachary-Bernard et al., 2019; Rey-Valette et al., 2019; Seebauer and Winkler, 2020).
	Clay-related subsidence due to increased or extreme drought	Deeper foundations, trees or terraces around light buildings to keep humidity in soils and prevent ground motions	Clay-related subsidence risks can be managed by appropriate adaptation measures at building scales (<i>medium confidence</i>) (Pritchard et al., 2015).

Note:

(a) Heatwave warnings and heat action plans as means for adaptation are discussed in Section 13.7.2.

Table SM13.15 | Literature sources used in the assessment of feasibility and effectiveness of adaptation options for cities, settlements and key infrastructures in Europe (Figure 13.20)

Impact type	Adaptation option	Effectiveness	Feasibility					
			Economic	Technological	Institutional	Sociocultural	Ecological	Geophysical
Reduction of thermal comfort due to increasing temperatures and extreme heat	Interventions in the building shell	Tillson et al. (2013); Sajjadian et al. (2015); Ibrahim and Pelsmakers (2018); Domínguez-Amarillo et al. (2019)	Tillson et al. (2013); Sajjadian et al. (2015); Murtagh et al. (2019)	Sajjadian et al. (2015)	Ibrahim and Pelsmakers (2018); Murtagh et al. (2019)	Ibrahim and Pelsmakers (2018); Murtagh et al. (2019)		Tillson et al. (2013); Ibrahim and Pelsmakers (2018); Domínguez-Amarillo et al. (2019)
	Ventilation	Tillson et al. (2013); van Hooff et al. (2014); Dodoo and Gustavsson (2016); Mulville and Stravaravdis (2016); Hamdy et al. (2017); Zinzi et al. (2017); Heracleous and Michael (2018); Ibrahim and Pelsmakers (2018); Dino and Meral Akgül (2019); Thomson et al. (2019)	van Hooff et al. (2014); Murtagh et al. (2019)	van Hooff et al. (2014)	Tillson et al. (2013); Mulville and Stravaravdis (2016); Murtagh et al. (2019)	Tillson et al. (2013); van Hooff et al. (2014); Mulville and Stravaravdis (2016); Ibrahim and Pelsmakers (2018); Murtagh et al. (2019); Thomson et al. (2019)		Tillson et al. (2013); van Hooff et al. (2014); Ibrahim and Pelsmakers (2018)
	Air conditioning	Jenkins et al. (2014a); Dodoo and Gustavsson (2016); Dino and Meral Akgül (2019)	Ferrara and Fabrizio (2017); Thomson et al. (2017)			Thomson et al. (2019)		Jenkins et al. (2014a)
	Shading	Tillson et al. (2013); van Hooff et al. (2014); Dodoo and Gustavsson (2016); Zinzi et al. (2017); Ibrahim and Pelsmakers (2018)	Tillson et al. (2013); van Hooff et al. (2014); Murtagh et al. (2019)	van Hooff et al. (2014)	Murtagh et al. (2019); Thomson et al. (2019)	Tillson et al. (2013); Ibrahim and Pelsmakers (2018); Murtagh et al. (2019)		van Hooff et al. (2014); Thomson et al. (2019)
	Green roofs, green walls	Cameron et al. (2014); van Hooff et al. (2014); Virk et al. (2015); Carvalho et al. (2017a); de Munck et al. (2018)	Cameron et al. (2014); Virk et al. (2015); Carvalho et al. (2017a); de Munck et al. (2018)	Cameron et al. (2014); Carvalho et al. (2017a); de Munck et al. (2018)	Cameron et al. (2014); Virk et al. (2014)	Cameron et al. (2014); van Hooff et al. (2014); Virk et al. (2015); Carvalho et al. (2017a); Derksen et al. (2017)	Virk et al. (2015)	van Hooff et al. (2014)
	Urban green spaces	Emmanuel and Loconsole (2015); Ward et al. (2016); Carvalho et al. (2017a); de Munck et al. (2018)	Carvalho et al. (2017a); de Munck et al. (2018)	Carvalho et al. (2017a); de Munck et al. (2018)	Emmanuel and Loconsole (2015)	Carvalho et al. (2017a); Derksen et al. (2017); Thomson et al. (2019)	de Munck et al. (2018)	Emmanuel and Loconsole (2015); Carvalho et al. (2017a); de Munck et al. (2018); Thomson et al. (2019)
	Use of 'cool' paints and coatings	Carnielo and Zinzi (2013); van Hooff et al. (2014); Virk et al. (2014); Virk et al. (2015); Zinzi (2016); Carvalho et al. (2017a)	van Hooff et al. (2014); Virk et al. (2015); Carvalho et al. (2017a); Murtagh et al. (2019)	Carnielo and Zinzi (2013); Carvalho et al. (2017a)	Virk et al. (2014)	Carnielo and Zinzi (2013); Virk et al. (2015); Murtagh et al. (2019)		Zinzi (2016)
	Escape to nearby non-urban destinations					Juschten et al. (2019a); Juschten et al. (2019b)		

Impact type	Adaptation option	Effectiveness	Feasibility					
			Economic	Technological	Institutional	Sociocultural	Ecological	Geophysical
Loss of critical services due to heatwaves and drought	Improvements in cooling systems	Jenkins et al. (2014a); Koch et al. (2014); Byers et al. (2015); Ferranti et al. (2016); Kingsborough et al. (2016); van Vliet et al. (2016b); Behrens et al. (2017); Bogmans et al. (2017)	Koch et al. (2014); van Vliet et al. (2016b); Behrens et al. (2017); Bogmans et al. (2017); EEA (2019a)	Sieber (2013); Ferranti et al. (2016)	Jenkins et al. (2014a); Koch et al. (2014); Byers et al. (2015); Hendel et al. (2016); Kingsborough et al. (2016); Behrens et al. (2017)			Sieber (2013); Koch et al. (2014); van Vliet et al. (2016b); Behrens et al. (2017)
	Shifting production to less water-intensive plants	Khan et al. (2016)	Khan et al. (2016); Behrens et al. (2017)	Khan et al. (2016)	Behrens et al. (2017)			
	Regulatory measures	Eisenack (2016)	Eisenack (2016)		Eisenack (2016)			
	Management measures	Gaudard et al. (2013); Hendel et al. (2015); Ferranti et al. (2016); Kingsborough et al. (2016); Ranzani et al. (2018); Wang et al. (2019)	Ferranti et al. (2016); Ranzani et al. (2018); Wang et al. (2019)	Hendel et al. (2015); Kingsborough et al. (2016)	Ferranti et al. (2016); EEA (2019a); Palkowski et al. (2019)		Hendel et al. (2015)	Gaudard et al. (2013); Hendel et al. (2015)
	Use of heat-resilient materials	Carnielo and Zinzi (2013); EEA (2019a); Wang et al. (2019)	Carnielo and Zinzi (2013); EEA (2019a); Wang et al. (2019)	Carnielo and Zinzi (2013); EEA (2019a); Wang et al. (2019)	Carnielo and Zinzi (2013); EEA (2019a); Wang et al. (2019)	Carnielo and Zinzi (2013); EEA (2019a); Wang et al. (2019)		
	Replacement of vulnerable infrastructure with resilient one	van Slobbe et al. (2016); Wang et al. (2019)	van Slobbe et al. (2016); Wang et al. (2019)					

Table SM13.16 | Reported adaptation limits for cities, settlements and key infrastructure in Europe (Figure 13.21)

Technical limits Physical characteristics of the existing housing stock prevent high ventilation (Tillson et al., 2013). There is limited efficacy of hard defences for high SLR (i.e., >1 m) and rapid rates of sea level rise (e.g., above 1 cm yr⁻¹) (see Box 13.1) (Umgiesser, 2020). Natural ventilation is limited by safety, noise and air pollution concerns (Tillson et al., 2013; van Hooff et al., 2014; Mulville and Stravaravdis, 2016). Too large sediment volumes are needed for beach nourishment (Galofré et al., 2016; Jiménez and Valdemoro, 2019). Wet-bulb temperature below 2°C is needed for snowmaking (Spandre et al., 2016; Hartl et al., 2018). Seawater cooling is feasible only for coastal plants (Behrens et al., 2017). Management optimisation is not applicable to run-of-river hydropower plants (Gaudard et al., 2013). Automation for flood discharge is not suitable for certain hydropower dams (EEA, 2020c). Water temperature caps can reduce thermal power availability and cause blackouts (Eisenack, 2016).
Socioeconomic limits Low flood probability prohibits the payoff of costly investments in home flood proofing (Poussin et al., 2015). Energy poverty limits households' capacity to adapt to overheating (Sanchez-Guevara et al., 2019; Thomson et al., 2019). High investments are needed for upgrading current drainage to new standards (EEA, 2020c). High installation costs exist for applying flood-proofing measures beyond critical substations (EEA, 2020a). There are no adaptation benefits from turning aerial transmission cables into underground cables in flood-prone areas (Sieber, 2013).
Environmental and regulatory limits Minimum Energy Performance standards cover only residential air conditioners (Palkowski et al., 2019). Space constraints exist on green infrastructure for flood management (Liu and Jensen, 2018). There is limited or no availability of free areas in higher altitudes, and there are orographic constraints for expanding skiable corridors Campos Rodrigues et al. (2018). There are limited water resources for increasing snowmaking (Spandre et al., 2016; Scott et al., 2019; Steiger et al., 2020). An impossible inland shift of tourism and settlements exists due to coastal urbanisation or geomorphology (Toimil et al., 2018). There is a lack of nearby alternative non-freshwater sources for plant cooling (Sieber, 2013).

Table SM13.17 | Present status of planned and implemented adaptation in cities, the energy sector, tourism, transport and industry in Europe (Table 13.1)

Sector	References
Cities	Reckien et al. (2015); EEA (2016b); Geneletti and Zardo (2016); Buurman et al. (2017); Davis et al. (2018); Gedikli and Balaban (2018); Reckien et al. (2018); CoM (2019); Pietrapertosa et al. (2019); Bertoldi et al. (2020)
Energy	EEA (2018); EEA (2019a); Gasho (2019)
Tourism	Gómez-Martín et al. (2014); Haanpää et al. (2015); Damm et al. (2017); Campos Rodrigues et al. (2018); Joye (2018); Landauer et al. (2018); Tirado et al. (2019)
Transport	Rotter et al. (2016); Battiston et al. (2017)
Industry and business	Averchenkova et al. (2016); Herrmann and Guenther (2017); Halkos et al. (2018); Schiemann and Sakhel (2018); Aguinaldo et al. (2019); D'Orazio and Popoyan (2019); CDSB (2020); de Bruin et al. (2020); Feridun and Güngör (2020); ECB (2021a); ECB (2021b)

SM13.6 Supplementary Material Supporting Section 13.7

Table SM13.18 | Literature sources used in the assessment of climate-sensitive infectious diseases (Figure 13.23)

Climate-sensitive infectious disease	References
Tick-borne encephalitis and borreliosis (Lyme)	Jaenson and Lindgren (2011); Rizzoli et al. (2011); Estrada-Pena et al. (2012); Jaenson et al. (2012); Medlock et al. (2013); Alkishe et al. (2017); Sykes and Makiello (2017); Tokarevich et al. (2017); Daniel et al. (2018); Medlock et al. (2018); Semenza and Suk (2018); Waits et al. (2018); Estrada-Pena and Fernandez-Ruiz (2020); Nah et al. (2020); Rogovskyy et al. (2020); Vandekerckhove et al. (2021)
West Nile virus	Semenza and Menne (2009); Medlock et al. (2015); Paz (2015); Proestos et al. (2015); Marini et al. (2016); Semenza et al. (2016); Vogels et al. (2017); Haussig et al. (2018); Vlaskamp et al. (2020); Marini et al. (2021); Rodriguez-Alarcon et al. (2021); Young et al. (2021)
Dengue, chikungunya, zika viruses	Caminade et al. (2012); Bouzid et al. (2014); Schaffner and Mathis (2014); Semenza et al. (2014); Kraemer et al. (2015); Cunze et al. (2016); Liu-Helmersson et al. (2016); Nsoesie et al. (2016); Rezza (2016); Shepard et al. (2016); Stanaway et al. (2016); Caminade et al. (2017); Tjaden et al. (2017); Mascarenhas et al. (2018); Medlock et al. (2018); Semenza and Suk (2018); Solimini et al. (2018); Tjaden et al. (2018); Liu-Helmersson et al. (2019); Messina et al. (2019); Metelmann et al. (2019); Ryan et al. (2019); Blagrove et al. (2020); Bruguera et al. (2020); Iwamura et al. (2020); Liu et al. (2020); Colon-Gonzalez et al. (2021); Oliveira et al. (2021); Ryan et al. (2021); Zeng et al. (2021)
Malaria	Caminade et al. (2014); Murdock et al. (2016); Piperaki and Daikos (2016); Hertig (2019); Fischer et al. (2020b); Karypidou et al. (2020)
Vibriosis	Baker-Austin et al. (2013); Escobar et al. (2015); Baker-Austin et al. (2017)

Table SM13.19 | Literature sources used in the assessment of feasibility and effectiveness of adaptation options for mortality, morbidity, exposure and stress from heat in Europe (Figure 13.24)

Impact type	Adaptation option	Effectiveness	Feasibility					
			Economic	Technological	Institutional	Sociocultural	Ecological	Geophysical
Heat	Behaviour-change measures	Khare et al. (2015); Hendel et al. (2017); Schuster et al. (2017)	Hendel et al. (2017)	Khare et al. (2015)	Khare et al. (2015)	Khare et al. (2015); Schuster et al. (2017)		
	Natural cooling	Alessandrini et al. (2019)		Alessandrini et al. (2019)		Alessandrini et al. (2019)		
	Building interventions	Åström et al. (2017); Taylor et al. (2018); Macintyre and Heaviside (2019); Murtagh et al. (2019)	Macintyre and Heaviside (2019)	Åström et al. (2017); Taylor et al. (2018); Macintyre and Heaviside (2019)		Åström et al. (2017); Macintyre and Heaviside (2019)		Fallmann et al. (2013)
	Green infrastructures	Richter (2016); Taylor et al. (2018); Rotzer et al. (2019); Venter et al. (2020)	Venter et al. (2020)	Richter (2016); Taylor et al. (2018); Rotzer et al. (2019); Venter et al. (2020)	Venter et al. (2020)	Taylor et al. (2018); Venter et al. (2020)	Rotzer et al. (2019); Venter et al. (2020)	Fallmann et al. (2013); Taylor et al. (2018); Rotzer et al. (2019); Venter et al. (2020)

Impact type	Adaptation option	Effectiveness	Feasibility					
			Economic	Technological	Institutional	Sociocultural	Ecological	Geophysical
Heat	Heat-proof land management	Fallmann et al. (2013); Åström et al. (2017); Montazeri et al. (2017)		Åström et al. (2017); Montazeri et al. (2017)	Donner et al. (2015); Åström et al. (2017)		Åström et al. (2017)	Fallmann et al. (2013); Donner et al. (2015); Montazeri et al. (2017)
	Heat health action plans	Gasparrini et al. (2015); Carmona et al. (2016); De'Donato et al. (2018)		Gasparrini et al. (2015); Carmona et al. (2016); De'Donato et al. (2018); Reischl et al. (2018); Morabito et al. (2019)	De'Donato et al. (2018); Reischl et al. (2018)	Gasparrini et al. (2015); De'Donato et al. (2018); Reischl et al. (2018); Morabito et al. (2019)	De'Donato et al. (2018); Casanueva et al. (2019); Morabito et al. (2019)	
	Bundle of options	Åström et al. (2017); Guo et al. (2018); Díaz et al. (2019)	Díaz et al. (2019)	Díaz et al. (2019)				Díaz et al. (2019)

SM13.7 Supplementary Material Supporting Section 13.8

Table SM13.20 | References supporting the examples of losses and damages to vulnerable livelihoods in Europe, differentiating for different categories of non-economic loss and damage (Table 13.2)

Example of losses and damages	Reference
Loss of livelihood, culture, health and well-being of the Sámi and the Nenets	Arctic Council (2013); Forbes et al. (2016); Hayashi (2017); Huntington et al. (2017); Mikhaylova (2018); Mustonen (2018); Inuit Circumpolar Council (2020); Feodoroff (2021)
Loss of key species in high-Arctic freshwater habitats, proliferation of introduced species and disruption of local food systems in Greenland, Finland, Sweden, northwest Russia and Scotland	Mustonen et al. (2018); Post et al. (2019); Frainer et al. (2020); Mustonen and Huusari (2020); Feodoroff (2021); Mustonen et al. (2021a); Mustonen et al. (2021b)
Warmer winters leading to loss of income from ice fishing and cultural heritage in Finland	Mustonen (2014); Mustonen and Huusari (2020)
Changes to marine food web resulting in loss of Indigenous knowledge and food insecurity in Greenland	Hayashi (2017); Pecl et al. (2017); Hayashi and Walls (2019); Inuit Circumpolar Council (2020)
Reduced yields on managed alpine grasslands decreasing the self-sufficiency of pastoral livestock farming in the Austrian, French and Swiss Alps	Brunner et al. (2019); Deléglise et al. (2019); Lavorel et al. (2019); Lavorel et al. (2020)
Reduced yields on semi-natural grasslands, compromising livestock feeding in winter, and ultimately decreasing viability of pastoralism in the Spanish Pyrenees	López-i-Gelats et al. (2016); Fernández-Giménez and Ritten (2020)
Retreating glaciers and changes in the landscape leading to loss of identity, culture and self-reliance in the Italian Alps (Alto Adige)	Jurt et al. (2015a); Jurt et al. (2015b)
Drought resulting in a reduction of provisioning (water) and regulating services (protection against floods) in the Western and Eastern Alps, Iberian Mountains and Dinaric Mountains	Leitinger et al. (2015); Mina et al. (2017); Schirpke et al. (2017); Strasser et al. (2019)
Increase of sea temperature leads to shifts in distribution of cold water species, reducing productivity at lower latitudes. Artisanal fisheries in Southern European coastal areas (Mediterranean) that rely on local, nearshore stocks can have difficulties to adapt	Lloret et al. (2018)

Table SM13.21 | References supporting Figure Box 13.2.1 'Cumulative impacts of climate and land-use change on reindeer herding as a traditional, semi-nomadic Sámi livelihood'

Indicator	References
Boundaries of reindeer herding areas in Sweden	Sámi Parliament in Sweden (2020)
Amount of snow	Ranasinghe et al. (2021)
Unstable ice conditions	Forbes et al. (2016); Mallory and Boyce (2018)
Frequent freeze-thaw cycles	Johansson et al. (2011); Hansen et al. (2014); Bokhorst et al. (2016); Rasmus et al. (2018)
Late snow melting during spring	Meredith et al. (2019)

Indicator	References
Heatwaves during summer	Skarin et al. (2010); Furberg et al. (2011); Löf (2013); Löf (2014); Meredith et al. (2019); Rosqvist et al. (2021)
Spread of new diseases	Omazic et al. (2019)
Insect harassment	Mallory and Boyce (2018); Tryland et al. (2019)
Psychological stress	Kaiser et al. (2010); Furberg et al. (2011); Stoor (2016)
Workload and costs	Furberg et al. (2011); Löf (2013); Löf (2014); Rosqvist et al. (2021)
Conflicts between herding communities and developers, authorities and members of local communities about the desirability of competing land uses	Lawrence (2014); Sehlén MacNeil (2015); Lawrence and Kløcker Larsen (2017); Persson et al. (2017); Beland Lindahl et al. (2018)
Self-determination and adaptive capacity	Brännlund and Axelsson (2011); Löf (2013); Brännström (2017); Allard (2018); Larsen and Raitio (2019)
Mining	Herrmann et al. (2014b); Eftestøl et al. (2019); Lawrence and Kløcker Larsen (2019); Österlin and Raitio (2020)
Hydropower	Össbo and Lantto (2011); Össbo (2018)
Forestry	Kivinen et al. (2012); Sandström et al. (2016); Fischer et al. (2020a)
Wind power	Skarin et al. (2015); Skarin and Alam (2017); Österlin and Raitio (2020)
Tourism	Nellemann et al. (2000); Skarin and Åhman (2014); Olsen (2016)

SM13.8 Supplementary Material Supporting Section 13.10

Table SM13.22 | Detected changes and attribution of climate-related impacts on land and in the ocean (Figure 13.27)

Assessment statement	Supporting references
Forest growth and production have been influenced by temperature and moisture conditions combined over recent centuries. The consequences of climate change have differed regionally, especially along the south-to-north axis.	Pretzsch et al. (2014); Reyer et al. (2014); Seidl et al. (2014); Gazol et al. (2015); Keenan et al. (2016); Reich et al. (2016); Tian et al. (2016); Alrahaleh et al. (2017); Ballantyne et al. (2017); Zlatanov et al. (2017); Humphrey et al. (2018); Marqués et al. (2018); Stocker et al. (2018); Vitali et al. (2018); Carnicer et al. (2019a); Ciais et al. (2019); Green et al. (2019); Yuan et al. (2019); Brodribb et al. (2020)
Tundra vegetation growth rate and shrub height have been accelerated by climate change.	Belonovskaya et al. (2016); Martin et al. (2017)
Drought consequences in the Mediterranean region have shown a significant increase in adverse effects, and outside the southern region the effects of drought have varied considerably.	Fantappiè et al. (2011); Giuntoli et al. (2013); Yigini and Panagos (2016); Potopová et al. (2017); Stagge et al. (2017); Samaniego et al. (2018); García-Herrera et al. (2019); Spinoni et al. (2019); Zhou et al. (2019).
Crops have decreased due to temperature-related regional changes with variable regional impact in Europe, and optimal conditions of some crops have moved northward.	García-Mozo et al. (2015); Long et al. (2016); Ceglar et al. (2017); Potopová et al. (2017); Zhao et al. (2017); Pérez-Domínguez and Fellmann (2018); Webber et al. (2018); Di Lena et al. (2019)
River floods have had increasingly damaging effects in Central Europe but have decreased in other regions.	Alfieri et al. (2015a); Polemio and Lonigro (2015); Ljungqvist et al. (2016); Blöschl et al. (2017); Kundzewicz et al. (2017); Paprotny et al. (2018); Berghuijs et al. (2019); Blöschl et al. (2019); Ganguli and Merz (2019); Lenderink et al. (2019); Umgiesser (2020)
Wildfire effects are jointly influenced by climate variables such as drought and temperature, but they are also highly influenced by management.	Moriendo et al. (2006); Moreno et al. (2014); Turco et al. (2014); Jolly et al. (2015); Tedim et al. (2015); Turco et al. (2016); de Rigo et al. (2017); Turco et al. (2017); Turco et al. (2018b); Michetti and Pinar (2019)
Marine heatwaves have induced mass mortality of sessile life forms, and such episodes have increased in frequency.	Garrabou et al. (2009); Munari (2011); Rivetti et al. (2014); Smale et al. (2015); Rubio-Portillo et al. (2016); Oliver et al. (2018); Damaraki et al. (2019); Holbrook et al. (2019); Smale et al. (2019a)
Terrestrial species relocation rates towards higher latitude and altitude have increased.	Scherrer and Körner (2011); Oliver et al. (2015); Melero et al. (2016); Stephens et al. (2016); Wiens (2016); Bowler et al. (2017); Spooner et al. (2018); Lehtikoinen et al. (2019a); van Klink et al. (2020a)
Marine species relocation from warm waters to previously colder, but now warming, waters has increased.	Fossheim et al. (2015); Hiddink et al. (2015); Montero Serra et al. (2015); van der Kooij et al. (2016); Chivers et al. (2017); García-Molinos et al. (2017); Cozzi et al. (2019); Vilà-Cabrera et al. (2019)
Coastal flood damaging effects have increased.	Haigh et al. (2011); Wahl et al. (2015); Malagon Santos et al. (2017); Garnier et al. (2018); Fernández-Montblanc et al. (2020); Umgiesser (2020)
Phenology changes were well documented in AR5, and subsequent studies have confirmed the trends.	Hassall et al. (2007); Visser et al. (2012); Karlsson (2014); Thackeray et al. (2016b); Mayor et al. (2017); Cohen et al. (2018b); Lehtikoinen et al. (2019b); Ettinger et al. (2020); Menzel et al. (2020)

Assessment statement	Supporting references
Vector-borne diseases have expanded northward.	Daniel et al. (2003); Jaenson et al. (2012); Medlock et al. (2013); Jore et al. (2014); Tokarevich et al. (2017); Semenza and Suk (2018)
Winter tourism has experienced decreased potential due to reduced snow cover and reliability of natural snow, with severity of loss highest at low altitudes.	Falk (2015); Falk and Vanat (2016); Klein et al. (2016); Beniston et al. (2018); Falk and Lin (2018); Schöner et al. (2019). Rain-on-snow event frequency has increased (Beniston and Stoffel, 2016).
Damage from thaw of permafrost has been detected in a large range of societally important infrastructures, such as buildings and roads.	Stoffel et al. (2014); Porfiriev et al. (2017); Raveland et al. (2017); Beniston et al. (2018); Duvillard et al. (2019)
Energy demand for cooling has increased and for heating has decreased due to increasing temperatures.	De Rosa et al. (2015); Spinoni et al. (2015); EEA (2017a policies and practices)
Macroeconomic damage for Europe has been detected.	Burke et al. (2015); Diffenbaugh and Burke (2019)
Shoreline erosion has been detected, but the literature is limited.	Castelle et al. (2018); Mentaschi et al. (2018)
Aquatic species relocation includes expansion northward, which in the southern region implies tropicalisation.	Zhang et al. (2017); Monchamp et al. (2018); Kärcher et al. (2019); van Klink et al. (2020a)
Heatwaves have induced mortality at increasing frequency and severity.	Shaposhnikov et al. (2015); Morabito et al. (2017); Vogel et al. (2019); Vicedo-Cabrera et al. (2021)
Ocean acidification combined with warming affects several aspects of marine commercial gain.	Lacoue-Labarthe et al. (2016); Fernandes et al. (2017)
Fisheries specimen size distribution has changed. The frequency of small specimen size has increased in southern regions of European waters.	Fortibuoni et al. (2015); Gamito et al. (2015); Teixeira et al. (2016); Ding et al. (2017); Ojea et al. (2017); Free et al. (2019); Stecf (2019)
Miscellaneous effects with <i>limited evidence</i> are not included in Figure 13.29. Several isolated examples of effects that can be attributed to climate change have been adequately reported.	For example, increase in groundwater heavy metal contamination from fractured aquifers, Bondu et al. (2016); effects in livestock, Handisyde et al. (2017) and Rojas-Downing et al. (2017); pathogen sensitivity, McIntyre et al. (2017) and Moretti et al. (2019); and heat damage to railway tracks, Ferranti et al. (2018)

Note:

This assessment is based on peer-reviewed literature in this chapter that reported observed evidence with at least 90% significance and usually with 95% significance or more.

Table SM13.23 | References for assessment of macroeconomic damages and gains for selected climate risks, measured by GDP and welfare for 1.5°C and 3°C GWL relative to no additional warming (Figure 13.33)

Risks	GWL	References
Change in agricultural yields	1.5°C	Szewczyk et al. (2018); Bosello et al. (2020); Szewczyk et al. (2020)
	3°C	Roson and Sartori (2016); Aaheim et al. (2017) ^a ; Szewczyk et al. (2018); Dellink et al. (2019) ^a ; Bosello et al. (2020); Szewczyk et al. (2020)
Change in labour productivity	1.5°C	Szewczyk et al. (2018); Bosello et al. (2020); Knittel et al. (2020) ^b ; Orlov et al. (2020) ^a
	3°C	Roson and Sartori (2016); Takakura et al. (2017) ^a ; Szewczyk et al. (2018); Bosello et al. (2020); Orlov et al. (2020) ^a ; García-León et al. (2021)
Change in energy demand	1.5°C	Szewczyk et al. (2018); Bosello et al. (2020)
	3°C	Szewczyk et al. (2018); Dellink et al. (2019) ^a ; Bosello et al. (2020)
Change in mortality due to heat	1.5°C	Szewczyk et al. (2018); Szewczyk et al. (2020)
	3°C	Roson and Sartori (2016); Aaheim et al. (2017) ^a ; Szewczyk et al. (2018); Dellink et al. (2019); Szewczyk et al. (2020)
Damage to economic sectors from water scarcity and drought	1.5°C	Faust et al. (2015) ^b ; Koopman et al. (2017) ^b ; Szewczyk et al. (2020); Teotónio et al. (2020) ^b
	3°C	Szewczyk et al. (2020); Teotónio et al. (2020) ^b
Change in energy supply	1.5°C	Bosello et al. (2020); Szewczyk et al. (2020)
	3°C	Aaheim et al. (2017) ^a ; Bosello et al. (2020); Szewczyk et al. (2020)
Damage to infrastructure from coastal flooding	1.5°C	Pycroft et al. (2016); Szewczyk et al. (2018); Bosello et al. (2020); Szewczyk et al. (2020)
	3°C	Pycroft et al. (2016); Roson and Sartori (2016); Aaheim et al. (2017); Szewczyk et al. (2018); Dellink et al. (2019) ^a ; Bosello et al. (2020); Parrado et al. (2020) ^a ; Szewczyk et al. (2020)
Damage to infrastructure from inland flooding	1.5°C	Dottori et al. (2018) ^a ; Szewczyk et al. (2018); Koks et al. (2019); Bosello et al. (2020); Parrado et al. (2020) ^a ; Szewczyk et al. (2020)
	3°C	Aaheim et al. (2017); Dottori et al. (2018) ^a ; Szewczyk et al. (2018); Dellink et al. (2019); Koks et al. (2019); Bosello et al. (2020); Szewczyk et al. (2020)

Notes:

- (a) Only larger European sub-regions are covered.
 (b) Single country or subset of countries.

SM13.9 Supplementary Material Supporting the Feasibility and Effectiveness Assessment

Background

The assessment aimed to provide an overview of the current knowledge of the feasibility and effectiveness (F&E) of selected adaptation options for key climate risks in Europe, and inform the design of the illustrative adaptation pathways in Section 13.10.2 (see SM13.11). Feasibility is understood as the potential for an adaptation option to be implemented. Feasibility depends on geophysical, environmental–ecological, technological, economic, sociocultural and institutional factors that enable or constrain the implementation. Effectiveness refers to the potential the option has to reduce risk compared with a baseline (see AR6-WGII Annex II).

Figure SM13.1 presents the main steps taken in the F&E assessment. The evidence presented in Figures 13.6, 13.14, 13.20 and 13.24 is the result of the available scientific evidence presented in the Supplementary Material (Tables SM13.1, SM13.5, SM13.15, SM13.19) and expert interpretation of these data. The approach was largely similar to the IPCC AR6 Special Report: Global Warming of 1.5°C (Chapter 4, de Coninck et al., 2018), and is further detailed in Singh et al. (2020).

Scientific articles were collected from a range of sources including the Global Adaptation Mapping Initiative (Berrang-Ford et al., 2021), focused literature searches by the author team, including contributing authors, and references suggested by reviewers. The assessment focused on those studies that empirically assessed (i.e., using case studies, models, experiments) the feasibility and effectiveness of adaptation options, thereby excluding conceptual and opinion articles. Only articles that explicitly considered measures to adapt to the observed impacts or projected risks of climate change were included to ensure conceptual consistency of the assessment, as is common in most systematic assessments on climate-change adaptation (Berrang-Ford et al., 2015). Moreover, articles were only included when they reported a clear link between the adaptation option and one or several of the feasibility and effectiveness assessment criteria. Articles listing general enabling and constraining factors to adaptation within a sector in general, for example, were not included in the F&E assessment. Grey literature and articles not in the English language were not included due to time and resource constraints. These articles may have created a bias in the datasets used for the assessment.

Detailed methods of the assessment approach and calculation of the scores for individual indicators and aggregation to the six feasibility dimensions can be found in Singh et al. (2020). The results visualised in the figures is the combination of scientific evidence and discussions among experts in the author team. Contributing authors were invited to participate when expertise within the Lead Author team was limited.

SM13.10 Supplementary Material Supporting the Key Risks and Burning Embers Assessments

The detailed methodology for the key risk assessment is described in Chapter 16. Lead and Contributing Authors in Chapter 13 have been assigned to assess the key risks literature based on their expertise and contributions to the chapter. Data were extracted from the references listed in Table SM13.22 and entered in five different spreadsheets (two spreadsheets for key risk 1 and one for each of the other key risks): reference, IPCC sub-regions, RCPs or any other corresponding climate scenarios, time periods, warming (GSAT) from pre-industrial levels derived according to the common climate dimensions (see Cross Chapter Box CLIMATE in Chapter 1), climate-impact drivers (Ranasinghe et al., 2021), SSPs (if available), climate models and type of simulations, impact models, sector(s) affected, risk metric, risk consequences and comments.

Following the data extraction, the key risk teams participated in expert elicitation workshops to reach consensus on the risk and temperature transition levels to feed into the construction of the burning ember diagrams in Figures 13.29–13.32 (see Zommers et al., 2020). The methodology is provided in Section SM13.11 and was slightly adapted for the assessments in this chapter. Results are summarised in Tables SM13.25–SM13.31.

General workflow for assessing effectiveness and feasibility of adaptation options




Figure SM13.1 | General workflow for assessing effectiveness and feasibility of adaptation options

Table SM13.24 | List of references supporting the assessment of the key risks (Figure 13.28)

Key Risk	References
KR1: Risks of human mortality and heat stress, and of ecosystems disruptions, due to heat extremes and increase in average temperatures	
Human health	Frolov et al. (2014); Jenkins et al. (2014b); Barbosa et al. (2015); Åström et al. (2017); Forzieri et al. (2017); Gasparrini et al. (2017); Hamdy et al. (2017); Hunt et al. (2017); Kendrovski et al. (2017); Mora et al. (2017); Cellura et al. (2018); Forzieri et al. (2018); Guo et al. (2018); Heracleous and Michael (2018); Arnell et al. (2019); Díaz et al. (2019); Revich et al. (2019); Rohat et al. (2019); Casanueva et al. (2020); Lee et al. (2020); Vanos et al. (2020); Ebi et al. (2021)
Marine ecosystems	Roebeling et al. (2013); Albouy et al. (2014); Brodie et al. (2014); Carstensen et al. (2014); Frolov et al. (2014); Herrmann et al. (2014a); Maugendre et al. (2014); Hennige et al. (2015); Serra et al. (2015); Wall et al. (2015); Cheung et al. (2016); García Molinos et al. (2016); Holt et al. (2016); Krumhansl et al. (2016); Lam et al. (2016); Ragazzola et al. (2016); Spencer et al. (2016); Stiasny et al. (2016); Thackeray et al. (2016b); Fernandes et al. (2017); Galli et al. (2017); Narita and Rehdanz (2017); Semenza et al. (2017); Thomsen et al. (2017); Townhill et al. (2017); Wang et al. (2017b); Benedetti et al. (2018); Corrales et al. (2018); Gao et al. (2018); Jokinen et al. (2018); Mangi et al. (2018); Riebesell et al. (2018); Sswat et al. (2018b); van der Spek (2018); Wang et al. (2018); Chefaoui et al. (2019); de la Hoz et al. (2019); Durant et al. (2019); Edelgeriev (2019); Herrera et al. (2019); Lotze et al. (2019b); Moullec et al. (2019); Petrik et al. (2019); Richon et al. (2019); Roggatz et al. (2019); Spivak et al. (2019); Bryndum-Buchholz et al. (2020); Clark et al. (2020); Maltby et al. (2020); Xi et al. (2021)
Terrestrial ecosystems	Gallego-Sala et al. (2010); Perch-Nielsen et al. (2010); Dury et al. (2011); San-Miguel-Ayanz et al. (2012); Filipe et al. (2013); Matzarakis et al. (2013); Steiger and Stötter (2013); Bedia et al. (2014); Frolov et al. (2014); Markovic et al. (2014); Matzarakis et al. (2014); Oliver et al. (2015); Urban (2015); Wu et al. (2015a); Brambilla et al. (2016); Grillakis et al. (2016); Polce et al. (2016); Scott et al. (2016); Thackeray et al. (2016b); Camia et al. (2017); EEA (2017a); Pecl et al. (2017); Sáenz-Romero et al. (2017); Vazquez et al. (2017); Vermaat et al. (2017); Dyderski et al. (2018); Ceglar et al. (2019); Ferretto et al. (2019); Jakoby et al. (2019a); Lotze et al. (2019b); Berberoglu et al. (2020); Feyen et al. (2020); Gianinetto et al. (2020); Qiu et al. (2020); Wamelink et al. (2020); Urvois et al. (2021); Xi et al. (2021)

Key Risk	References
KR2: Risk of losses in crop production, due to compound heat and dry conditions, and extreme weather	
	Caffarra et al. (2012); Sutton et al. (2013); Deryng et al. (2014); Donatelli et al. (2015); Reidsma et al. (2015); Bird et al. (2016); Castellanos-Frias et al. (2016); Knox et al. (2016); Webber et al. (2016); Diogo et al. (2017); Holman et al. (2017); Nielsen et al. (2017); Popp et al. (2017); Siebert et al. (2017); Williges et al. (2017); Ben-Ari et al. (2018); Parent et al. (2018 with an appropriate use of the genetic variability of flowering time); Ruiz-Ramos et al. (2018); Szewczyk et al. (2018); Webber et al. (2018); Ceglar et al. (2019); Chen et al. (2019); EEA (2019b); Grillakis (2019); Moretti et al. (2019); Papadimitriou et al. (2019); Toreti et al. (2019); Feyen et al. (2020); Jäger et al. (2020); Huttunen et al. (2021); Pedde et al. (2021)
KR3: Risk of water scarcity to multiple interconnected sectors	
	Blauhut et al. (2015); Collet et al. (2015); Kebede et al. (2015); Papadaskalopoulou et al. (2015b); Blauhut et al. (2016); Gain et al. (2016); Gampe et al. (2016); Schleussner et al. (2016); Stahl et al. (2016); Koopman et al. (2017); van Vuuren et al. (2017); Byers et al. (2018); Greve et al. (2018); Iglesias et al. (2018); Manousseli et al. (2018); Naumann et al. (2018); Tobin et al. (2018b); Arnell et al. (2019); Garnier and Holman (2019); Harrison et al. (2019); Koutroulis et al. (2019); Morote et al. (2019); Papadimitriou et al. (2019); Bisselink et al. (2020); Cammalleri et al. (2020); Teotónio et al. (2020); Kebede et al. (2021); Naumann et al. (2021)
KR4: Risks to people, economies and infrastructures due to coastal and inland flood hazards	
Coastal flooding	Ciscar et al. (2014); Marzeion and Levermann (2014); Clark et al. (2016); Pycroft et al. (2016); Reimann et al. (2018); Voudoukas et al. (2018); Arnell et al. (2019); Harrison et al. (2019); Voudoukas et al. (2020); Haasnoot et al. (2021)
Inland flooding	Jongman et al. (2014); Alfieri et al. (2015b); Alfieri et al. (2016a); Alfieri et al. (2016b); Alfieri et al. (2017); Alfieri et al. (2018); Ciscar et al. (2018); Dottori et al. (2018); Paprotny et al. (2018); Arnell et al. (2019); Harrison et al. (2019); Papadimitriou et al. (2019); Dottori et al. (2020); Hosseinzadehtalaei et al. (2020); Lange et al. (2020); Merz et al. (2021)

Table SM13.25 | Transition and confidence levels for the risks of human mortality and heat stress due to heat extremes and increase in average temperature (Figures 13.28, 13.29)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
Risk level with present to medium adaptation, including SSP2–SSP4			
Undetectable to moderate risk	0.7–1.0	high	The range and confidence are supported by evidence from the detection and attribution assessment (Figure 13.27; Table SM13.1; Vicedo-Cabrera et al., 2021). The range reflects warming between 1995 and 2014 as reported in WGI Cross-Chapter Box 2.3 in Chapter 2. Furthermore, the climate-impact drivers (both mean air temperature and extremes) emerged already in the historical period with <i>medium confidence</i> (WGI Figure 12.7). Detected risk to date (selected from the D&A assessment in Section 13.10.1): Shaposhnikov et al. (2015); Morabito et al. (2017); Vogel et al. (2019); Vicedo-Cabrera et al. (2021)
Moderate to high risk	1.6–2.2	high	The magnitude of consequences attributed to a number of risk metrics, such as mortality and population exposure to heat extremes, all increase beyond 1.5°C GWL. Naumann et al. (2020a) show an increase already at 1.5°C GWL, but there is less evidence and consensus that a transition happens before or at 1.5°C GWL. The risk also increases less rapidly for every indicator, or at least there is little consensus in the literature. It is higher and more likely in southern regions but not everywhere. Magnitude of risk consequences reported in the literature to support this transition: – Excess mortality attributable to climate change: 1.5–1.67% (NEU, WCE, SEU) (Gasparrini et al., 2017) – Proportion of the European population at risk of heat stress: 161 million (Rohat et al., 2019) in EU-27 plus the UK – Additional attributable deaths in Europe: 30,867 (with no climate change: 17,384) – Increase in heatwave excess deaths in SEU, WCE and NEU with the highest increase in NEU (Guo et al., 2018) – People annually exposed to a once-in-50-year heatwave: 176 million (present value: 9.6 million) and heat fatalities per year: 52,182 (present value: 2752) (Naumann et al., 2020a) Literature supporting this transition: Forzieri et al. (2017); Gasparrini et al. (2017); Hunt et al. (2017); Kendrovski et al. (2017); Guo et al. (2018); Rohat et al. (2019); Lee et al. (2020); Naumann et al. (2020b)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
High to very high risk	2.9–3.5	<i>medium</i>	Some studies point to a transition at higher levels and others at lower levels; however, the risk becomes more persistent and increases in every region as well as its likelihood. The ability to adapt is not completely excluded, but assuming current adaptation conditions or SSP2/SSP4, very high risk will be realised under 3°C GWL. Magnitude of risk consequences reported in the literature to support this transition: At 3°C GWL: – Proportion of the European population at risk of heat stress compared with the reference period 1982–2005 (in parentheses): 216 million / 48.4% (2.1 million / 0.4%) (Rohat et al., 2019) – Extreme-heat-related deaths per 10 million inhabitants (1981–2010): 2817 (54.7) (Forzieri et al., 2017) – Heat fatalities per year: 90 thousand (Naumann et al., 2020a) – Extreme-heat-related deaths per 10 million inhabitants: 2816.9 in EU-27 plus the UK (Forzieri et al., 2017) At 3.5–4°C GWL: – Attributable death: 117,333 (with no climate change: 16,303) (Kendrovski et al., 2017) – Increase in heatwave-related deaths compared with the present warming levels: 200–280% (highest increase in NEU) Literature supporting this transition: Forzieri et al. (2017); Kendrovski et al. (2017); Rohat et al. (2019); Casanueva et al. (2020); Naumann et al. (2020b)
Risk level with high adaptation, including SSP1			
Undetectable to moderate risk	0.7–1.0	<i>high</i>	The range and confidence level are supported by evidence from the detection and attribution assessment (Figure 13.29; Table SM13.1; Vicedo-Cabrera et al., 2021). The ranges reflect warming between 1995 and 2014 as reported in WGI Cross-Chapter Box 2.3 in Chapter 2. Furthermore, the climate-impact drivers (both mean air temperature and extremes) emerged already in the historical period with <i>medium confidence</i> (Figure 12.7; Ranasinghe et al., 2021). Detected risk to date (from D&A assessment in Section 13.10.1): Shaposhnikov et al. (2015); Morabito et al. (2017); Vogel et al. (2019); Vicedo-Cabrera et al. (2021); additional evidence from the F&E assessment is given in Tables SM13.5.3 and SM13.6.2.
Moderate to high risk	2.5–3.8	<i>medium</i>	This transition is based on a body of literature which considers the adaptation needed to maintain the number of deaths at the current level or SSP1 conditions. It confirms that the mortality rates are smaller and the number of people at risk of heat stress are much fewer than those with only present to moderate adaptation (Åström et al., 2017; Guo et al., 2018; Diaz et al., 2019; Rohat et al., 2019). For example, Åström et al. (2017) calculate the rate of adaptation needed at around 2°C GWL to maintain the number of deaths at the present level and found this rate to be over 75%. Various caveats concern the assumptions of full acclimatisation above 3°C and the existence of possible physiological limits (Vanos et al., 2020). Ebi (2021) have found that the burning ember for heat-related mortality and morbidity with high adaptation (e.g., SSP1) has a transition between 2 and 3°C GWL. Considering that this is a global study and taking into account the evidence on high adaptation, we conclude that the transition level is somewhere above 2°C but below 4°C (Åström et al., 2017; Guo et al., 2018; Diaz et al., 2019). Additional evidence from the F&E assessment is given in Tables SM13.5.3 and SM13.6.2.
High to very high risk	Does not reach this level	n.a.	There is little evidence that the risk will reach this level assuming full adaptation because the adaptation potential is still higher in high-adaptation scenarios in several regions (NEU), e.g., and in agreement with, the burning embers in Ebi et al. (2021). We concluded that under a high-adaptation scenario, the risks do not reach this threshold.

Table SM13.26 | Transition and confidence levels for the risks of marine ecosystems disruptions due to heat extremes and increase in average temperatures (Figure 13.28)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the risk-level transition	Explanation and references
Undetectable to moderate	0.9–1.7	medium	A comprehensive summarising assessment across all impacts and risks on or for marine systems is impaired by the uneven distribution of available information on their levels and development driven by RCP scenarios at 1.7°C, 2.5°C and above 4°C GWL, making the determination of transitions challenging. The transition from undetectable to moderate starts around or less than 1°C to 1.7°C with <i>medium confidence</i>. Detected to date (selected from the D&A assessment in Section 13.10.1): – Marine heatwaves that have increased in frequency and induced mass mortality (Garrahou et al., 2009; Munari, 2011; Rivetti et al., 2014; Rubio-Portillo et al., 2016; Smale et al., 2019b) – Marine species relocation (Fosheim et al., 2015; Hiddink et al., 2015; Montero Serra et al., 2015; van der Kooij et al., 2016; Chivers et al., 2017; Cozzi et al., 2019) – Detected changes in range expansion (Birchenough et al., 2015), species distribution and species phenologies leading to a change from plankton communities (Burrows et al., 2019) to subtropicalisation of European pelagic fish communities (Montero Serra et al., 2015) – Change in size of catch of fisheries (Fortibuoni et al., 2015; Gamito et al., 2015; Teixeira et al., 2016; Ding et al., 2017; Ojea et al., 2017; Free et al., 2019; Stecf, 2019) Projected risk (further information in Figure 13.11): At 1.7°C GWL: – Increasing risks across systems (from habitat losses to biodiversity and ecosystem services); while observed impacts were or are most severe first in the Mediterranean, at this GWL risks projected to extend to transitional waters – 13% diversity losses YB19 (25–40% loss in animal biomass) (Bryndum-Buchholz et al., 2019; Lotze et al., 2019a) – 20% loss coastal wetland, 7% loss in ecosystem value (Roebeling et al., 2013) – 60% habitat reduction for fish in the Mediterranean (Clark et al., 2020), impacting commercially important fish (e.g., anchovy) (Raybaud et al., 2017) – Loss of 20–40% change in catch potential of shellfish and 20% in fish (Lam et al., 2016; Fernandes et al., 2017)
Moderate to high	2–2.2	low	Note: Confidence level is <i>low</i> because there is a distinct gap in information on risk development for levels between 1.7 and 2.5°C GWL. At 2°C GWL: – Increase in area suitable for viruses (Semenza et al., 2017) – Increase in likelihood of marine heatwaves (Frölicher et al., 2018)
High to very high	2.5–3	high	At 2.5°C GWL: – Strong risks across shallow-water ecosystems due to the combined warming and SLR: projected 28% loss in coastal wetlands and 10% ecosystem value (Roebeling et al., 2013); regionally up to 80% loss in wetland (Spencer et al., 2016) – Drowning of intertidal flats in TEUS (van der Spek, 2018) – Changes in species dominance (Moullec et al., 2019) – Diversity loss of 20% in TEUS and 40% in NEUS (García Molinos et al., 2016) – Losses in shellfish biomass in TEUS increasing from 40 to 50% locally and up to 80% in response to warming, and resulting habitat loss with significant economic losses (Fernandes et al., 2017; Galli et al., 2017; Mangi et al., 2018), 30% loss of fin fish (Cheung et al., 2016; Lam et al., 2016) and loss of habitat for economically important shellfish in the SEUS (Mangi et al., 2018) – Decrease in potential catch of cod and other economically important species (Maltby et al., 2020); needs changing in consumer behaviour and fishing At 3°C GWL: – Loss of important habitat formers in coastal systems (Brodie et al., 2014) – Decrease in primary productivity (Maugendre et al., 2014) Above 4°C GWL – 1.5 to 50% losses in macroinvertebrates, – Doubling of mortality of larva of economically important species, – 30–50% diversity losses, – 25–70% lower marine animal biomass – shellfish production collapses, – loss of habitat of species important for climate change mitigation, such as wetlands, limits their contribution

Table SM13.27 | Transition and confidence levels for the risks of terrestrial ecosystems disruptions due to heat extremes and increase in average temperatures (Figure 13.28)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
Undetectable to moderate risk	1–1.5	<i>medium</i>	There is increasing evidence across systems, but impacts are not yet considered severe and widespread based on the evidence. Detected to date (selected from the D&A assessment in Section 13.10.1): – Forest growth and production has been impacted by temperature and moisture over the past century (Pretzsch et al., 2014; Seidl et al., 2014; Gazol et al., 2015; Tian et al., 2016; Alrahahleh et al., 2017; Zlatanov et al., 2017; Marqués et al., 2018; Carnicer et al., 2019a; Ciais et al., 2019; Yuan et al., 2019; Brodribb et al., 2020). – Compound events of heat and drought have impacted forest growth with legacy effects into the next year (Schuldt et al., 2020). – Tundra vegetation shrub encroachment has been accelerated by climate change (Belonovskaya et al., 2016; Martin et al., 2017). – Drought consequences in the Mediterranean region (Fantappiè et al., 2011; Stagge et al., 2017; García-Herrera et al., 2019). – Terrestrial species relocation towards higher latitude and altitude have increased in rate (Scherrer and Körner, 2011; Oliver et al., 2015; Melero et al., 2016; Stephens et al., 2016; Wiens, 2016; Bowler et al., 2017; Spooner et al., 2018; Lehtikoinen et al., 2019a; van Klink et al., 2020b). – Range shifts have altered boreal and alpine tundra (Elmhagen et al., 2015; Post et al., 2019; Mekonnen et al., 2021) and greened polar deserts (Myers-Smith et al., 2020). – Climate-induced biodiversity declines have been detected in thermosensitive groups (Hellmann et al., 2016; Habel et al., 2019; Harris et al., 2019; Crossley et al., 2020; Soroye et al., 2020) including many freshwater plants, molluscs, flying insects, amphibians, reptiles, birds and fishes (Myers et al., 2017; Jarić et al., 2019; Kärcher et al., 2019; Seibold et al., 2019; van Strien et al., 2019; van Klink et al., 2020a). – Phenology changes (Hassall et al., 2007; Visser et al., 2012; Karlsson, 2014; Szabó et al., 2016; Thackeray et al., 2016b; Mayor et al., 2017; Asse et al., 2018; Cohen et al., 2018b; Bobretsov et al., 2019; Peaucelle et al., 2019; Menzel et al., 2020; Rosbakh et al., 2021). Warming between 1.5°C and 2°C GWL is projected to increase the impact including: – Expansion of the subtropical vegetation in SEU (Feyen et al., 2020) and large-scale threats for forest areas WCE (Dyderski et al., 2018) – Loss of 50% of the tundra biome in NEU and 40% WCE alpine regions (Feyen et al., 2020) – Loss in peatland area in NEU (Qiu et al., 2020) – Some losses in grassland in WCE (Vermaat et al., 2017) – Extinction risk of 5% of species in Europe (Urban (2015) – Loss of 6% of common and 77% of rare freshwater species extinction (Markovic et al., 2014) – 50% of protected area loss of mollusc and freshwater fish (Markovic et al., 2014) – Increasing changes in phenology, creating consumer primary producer mismatch (Thackeray et al., 2016a) – Habitat loss for pollinators in NEU (Rasmont et al., 2015), increasing risk to sensitive pollinators; uncertain net impacts on the pollination service – Increases in burned area (Wu et al., 2015a) – Increase in soil erosion by 8% in SEU (Berberoglu et al., 2020), increase in rainfall erosion especially in NEU (Panagos et al., 2017) and increase in soil loss from fire (Pastor et al., 2019b); however, soil erosion is made of many different components (e.g., precipitation, soil type, topography, land use and land management, and hence human impacts and vegetation responses to climate change are difficult to separate from climatic effects – Autonomous adaptation allowing species to shift ranges (if protected areas are sufficiently large and corridors are available, freshwater bodies are connected and, for plants, if soil types and hydrology are appropriate); opportunity for adaptation and reversibility which currently reduces the risk
Moderate to high risk	2–2.5	<i>medium</i>	The risk is increasing for a larger number of ecosystems, and the extent of loss and impact is becoming irreversible: – Increase in wildfire risk in the SEU (Bedia et al., 2014) and increase in burnt area (Camia (2017) – Expansion of temperate flora in 30% of NEU, 60% of tundra NEU and 50% of alpine tundra (Feyen et al., 2020) – 50% carbon loss in blanket bogs NEU (Ferretto et al., 2019) – 12% loss of meadows (Wamelink et al., 2020) – Increase in pests, such a bark beetle, in WCE (Jakoby et al., 2019b) – Increased extinction risk (Warren et al., 2018) Above 2.5°C GWL: – 70% range contraction in birds in the Alps (Brambilla et al., 2016) – Increase in threatened forest area by 46% in WCE (Dyderski et al., 2018) – Up to doubling in burned area (Wu et al., 2015a; Camia, 2017; EEA, 2017b)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
High to very high risk	3.3–3.8	high	The magnitude of several indicators (including losses) are projected to increase very strongly above 3°C GWL: – NEU and SEU tundra nearly completely lost, 70% of alpine tundra (Feyen et al., 2020) – Strong expansion of temperate flora into NEU and subtropical flora into WCE (Feyen et al., 2020) – Peat turning into carbon source (Qiu et al., 2020) – 8–16% of species at risk of extinction (Urban, 2015; Warren et al., 2018) Above 3.5°C GWL: – Large expansion of tree pest species in MED and WCE (Urvois et al., 2021) – 25% increase in soil erosion in SEU (Berberoglu et al., 2020) – Large habitat losses and diversity loss of 10–48% of pollinators NEU (Rasmont et al., 2015) – Species risk of extinction of 16% (Urban (2015) – Strong increases of the burned areas in Europe (Wu et al., 2015a; EEA, 2017b) – WCE threatened forest area 59% (Dyderski et al., 2018)

Table SM13.28 | Transition and confidence levels for risk of losses in crop production due to compound heat and dry conditions, and extreme weather (Figures 13.28, 13.30)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
Risk level with low adaptation			
Undetectable to moderate risk	0.8–1.1	medium	This transition is supported by the D&A assessment in Section 13.10.1 and references therein. This transition is placed around the 1°C GWL range. The impacts have been mainly attributed in the case of the extremes, where events in years such as 2010, 2016 and 2018 indicate the future on agricultural yields given the expected higher frequency and severity of these events. The average losses across Europe are not currently sufficiently large for risk to be widespread. At 1.5°C–1.7°C GWL: – Likelihood of compound events which has led to recent large wheat losses are projected to become 12% more frequent, challenging farming systems and yield forecasting systems (Ben-Ari et al., 2018) – Maize yield losses across Europe of 10–25% (Deryng et al., 2014; Webber et al., 2018; Feyen et al., 2020), though strongest in SEU – Wheat productivity not impacted at 1.7°C GWL (Webber et al., 2018) – Changes in SEU as growing regions shift northward for melons (Bisbis et al., 2019), tomatoes and grapevines reaching NEU and EEU (Hannah et al., 2013; Litskas et al., 2019) – Beginning of abandonment of agricultural land in SEU (Holman et al., 2017) – Warming that causes range expansion and alters host pathogen association of pests, diseases and weeds (Caffarra et al., 2012; Pushnya and Shirinyan, 2015; Latchininsky, 2017) – Regionally predicted reduction in rainfall which can lead to carryover of herbicides (Karkanis et al., 2018)
Moderate to high risk	2.5–3	medium	At 2°C GWL: – 30–50% yield reduction of rain-fed maize in SEU (EEA, 2019b; Feyen et al., 2020) – Increase in agricultural droughts (Toreti et al., 2019) – Locally strong changes in yield in arable systems (Diogo et al., 2017) – Increase in toxins in maize in SEU and emergence in WCE (Moretti et al., 2019) – 15% yield loss for sugar beet in rain-fed system in SEU (EEA, 2019b) At 2.7°C GWL: – Impacts becoming more widespread as agroclimatic zones move (Ceglar et al., 2019), though there are still regional gains in parts of WCE and NEU (Szewczyk et al., 2018) – Increasing uncertainty of the impact on wheat and rapeseed yields in SEU and WCE (Donatelli et al., 2015) – 108% increase in expanse of damaging weed for winter crops (Castellanos-Frias et al., 2016) – Heat stress on wheat in EEU (Ceglar et al., 2019) – Negative impact on pollinators resulting in reduced visits of bees (Nielsen et al., 2017) (see also Section 13.3.1.3) – Between 2.5°C–2.7°C and 3°C GWL, more crops being affected, though there is uncertainty of the extent of potential losses with some gains possible

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
High to very high risk	3.8–4	low	At 3°C–4°C GWL: Most of the evidence focuses on temperatures below 3°C GWL and above 4°C GWL with <i>limited evidence</i> of impacts between 2.7 and 4°C GWL. Given that only a very small number of papers are available at these warming levels, the confidence level is <i>low</i> .
			At 3.3°C GWL: – Loss of tomato production in parts of SEU (Bird et al., 2016) – 170% increase in expanse of damaging weed for winter crops (Castellanos-Frias et al., 2016) – NEU becoming a suitable climate for the wheat pest <i>Lolium rigidum</i> (Castellanos-Frias et al., 2016) At 4°C GWL and beyond: – Constant drought conditions similar to spring/summer 2018 (Ben-Ari et al., 2018) – Maize yield losses across Europe of 50–100% (Deryng et al., 2014; Webber et al., 2018; Feyen et al., 2020) – Losses in spring wheat (Deryng et al., 2014), though gains projected for EEU – Reduction in grassland biomass and start seeing losses in NEU (Jäger et al., 2020) – Increasing losses in agricultural yield SEU and WCE (Szewczyk et al., 2018) – Increased asynchrony between the larvae-resistant growth stages of grapevine and larvae occurrence (Caffarra et al., 2012)
Risk level with high adaptation			
Undetectable to moderate risk	0.8–1.1	medium	This transition is supported by the D&A assessment in Section 13.10.1 and the references therein (Table SM13.1), and is placed around the 1°C GWL range. High levels of implementation of multiple adaptation options are assumed for this high-adaptation burning ember. Agricultural production risks can be reduced through irrigation given the availability water resources and suitable infrastructure (Siebert et al., 2017; Ruiz-Ramos et al., 2018; Feyen et al., 2020). The ability to adapt through irrigation is constrained as increasing water is needed to reduce impacts of heat and reduce yield losses with higher temperature, and water availability is increasingly limited: – Irrigation can reduce projected heat and drought stress (e.g., for wheat and maize) (Siebert et al., 2017; Ruiz-Ramos et al., 2018; Feyen et al., 2020). – Irrigation of wheat reverses yield losses across Europe at 2°C GWL to become gains, while yield losses in maize in SEU are reduced from as much as 80 to 11% (Feyen et al., 2020). – Changes to cultivars and sowing dates can reduce yield losses, but are insufficient to fully ameliorate losses projected at 3°C warming and above, with an increase of risk from north to south and for crops, such as maize, growing later in the season (Ruiz-Ramos et al., 2018; Feyen et al., 2020). – Use of longer-season varieties can compensate for heat stress on maize in WCE and lead to yield increases for NEU (Siebert et al., 2017; Ceglar et al., 2019). – Moving the growth cycle towards a cooler part of year reduces the period for photosynthesis and grain filling (Ruiz-Ramos et al., 2018; Holzkämper, 2020). – Physiological constraints, such as lack of sufficient light in winter and spring, might hinder exploitation of changes in phenology and hence the potential for longer growing seasons. – Crop breeding for drought and heat tolerance can improve sustainability of agricultural production under future climate (Costa et al., 2019b), particularly in SEU at 3°C GWL (Senapati et al., 2019). – Multifunctional land use can reduce the dependency on one crop and source of income (Holman et al., 2017). – Agricultural water management adaptation, such as irrigation, reallocating water to other crops, improving use efficiency and soil water conservation practices, reduces risks (Iglesias and Garrote, 2015). – In-season forecasts of climate impacts on yield were used during the 2018 drought for European wheat to inform policy actions (van der Velde et al., 2018). – There is greater need for pesticides to control and maintain production, due to range expansion and altered host pathogen association of pests, diseases and weeds affecting the health of European crops (Caffarra et al., 2012; Pushnya and Shirinyan, 2015; Latchinsky, 2017). – Simplifying procedures for obtaining subsidies and insurance premiums and interest rates can incentivise adoption of climate-friendly agricultural methods (Garrote et al., 2015; Iglesias and Garrote, 2015; Zakharov and Sharipova, 2017; Hamidov et al., 2018; Wiréhn, 2018). Water availability and competing uses is considered in KR3. Additional evidence from the F&E assessment is given in Table SM13.5.
Moderate to high risk	Does not reach this threshold	medium	Given high adaption, with the adoption of multiple options, including the availability of sufficient irrigation water and infrastructure, there is little evidence that the risk will reach this level. KR3 considers the availability of water and competing demand, but sufficient water availability required to supply crop irrigation is assumed for the risk level with high adaptation of KR2. See undetectable to moderate risk for the evidence above for evidence of adaption options (Table SM13.5), including irrigation. We concluded that under a high adaptation scenario, the risks do not reach this threshold.
High to very high risk	Does not reach this threshold	n.a.	As per moderate to high risk

Table SM13.29 | Transition and confidence levels for the risks of water scarcity to multiple interconnected sectors (Figures 13.28, 13.31)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
Level of risk due to water scarcity in SEU with low adaptation			
Undetectable to moderate risk	0.7–1.1	high	There is already a moderate risk of water scarcity (according to the D&A in Figure 13.27). At 0.6°C, 6 months of drought duration are estimated. In 1981–2010 (–0.7°C), 48.1 million people were exposed to moderate water scarcity. Literature supporting this transition: Blauhut et al. (2015); Kebede et al. (2015); Blauhut et al. (2016); Gain et al. (2016); Gampe et al. (2016); Stahl et al. (2016); Bisselink et al. (2018); Byers et al. (2018); Naumann et al. (2018); Arnell et al. (2019); Harrison et al. (2019); Koutroulis et al. (2019); Bisselink et al. (2020); Cammalleri et al. (2020); Teotónio et al. (2020); Kebede et al. (2021); Naumann et al. (2021)
Moderate to high risk	1.3–1.7	high	– At 1.5°C, 55.7 million (i.e., +7.4 million) people under moderate water stress – At 1.8°C, Water Exploitation index increase of 230% and transition from moderate to high – At 2°C, doubling of drought duration – At 2°C, 54% of population water stressed – At 2°C, drought duration 10 months Literature supporting this transition: Kebede et al. (2015); Gampe et al. (2016); Schleussner et al. (2016); Bisselink et al. (2018); Byers et al. (2018); Naumann et al. (2018); Tobin et al. (2018a); Arnell et al. (2019); Harrison et al. (2019); Bisselink et al. (2020); Cammalleri et al. (2020); Feyen et al. (2020); Teotónio et al. (2020); Kebede et al. (2021); Naumann et al. (2021)
High to very high risk	2.5–3.5	medium	– At 1.5°C, 55.7 million people (i.e., +7.4 million) under moderate water stress – at 1.8°C, Water Exploitation index increase of 230% and transition from moderate to high – At 2°C, doubling of drought duration – At 2°C, 54% of population water stressed – At 2°C, drought duration 10 months Literature supporting this transition: Kebede et al. (2015); Gampe et al. (2016); Bisselink et al. (2018); Byers et al. (2018); Naumann et al. (2018); Tobin et al. (2018a); Arnell et al. (2019); Harrison et al. (2019); Koutroulis et al. (2019); Bisselink et al. (2020); Cammalleri et al. (2020); Teotónio et al. (2020); Kebede et al. (2021); Naumann et al. (2021)
Level of risk due to water scarcity in SEU with high adaptation			
Undetectable to moderate risk	0.9–1.3	medium	Improvements in water efficiency and in behaviour are very effective (>25% of damage avoided) in some SSP scenario improvements compared with baseline (Papadimitriou et al., 2019). There is some adaptation deficit now that can be addressed (see F&E water assessment in Table SM13.1). Literature supporting this transition: Papadaskalopoulou et al. (2015b); van Vuuren et al. (2015); Greve et al. (2018); Iglesias et al. (2018); Garnier and Holman (2019); Morote et al. (2019); Papadimitriou et al. (2019); and papers used in the F&E assessment (Table SM.13.1)
Moderate to high risk	1.8–2.2	medium	Investment in large water infrastructure and advanced technologies (including storage), water transfer, water recycling and reuse, and desalination (Papadaskalopoulou et al., 2015a; Greve et al., 2018) is needed. This will buy a bit of time in terms of coping with an increase of 0.5°C GWL. Literature supporting this transition: Papadaskalopoulou et al. (2015b); van Vuuren et al. (2015); Greve et al. (2018); Iglesias et al. (2018); Garnier and Holman (2019); Morote et al. (2019); Papadimitriou et al. (2019); and papers used in the F&E assessment (Table SM.13.1)
High to very high risk	2.8–3.8	low	Transformational adaptation is needed, but ultimately planned relocation of industry and development of alternative livelihoods may be needed. Also, there are trade-offs with other adaptation options in need of water (Papadaskalopoulou et al., 2015a; Greve et al., 2018). Adaptation will buy a bit of time in terms of coping with an increase of 0.2°C GWL. Literature supporting this transition: Papadaskalopoulou et al. (2015b); van Vuuren et al. (2015); Greve et al. (2018); Iglesias et al. (2018); Garnier and Holman (2019); Morote et al. (2019); Papadimitriou et al. (2019); and papers used in the F&E assessment (Table SM.13.1)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
Level of risk due to water scarcity in WCE with low adaptation			
Undetectable to moderate risk	0.6–1.8	medium	– At 0.6°C, 5.5–6.8 months of drought – At 1.5°C, 6.8–8.3 months of drought – In 1981–2010 (~0.7°C), 4.3 million people exposed to moderate water scarcity – At 1.5°C, 4.7 million people (i.e., +0.7 million) under moderate water stress – Number of people exposed is 1 order of magnitude smaller than in SEU (tens of millions) – Increase in exposure to moderate, but not severe, water scarcity (whereas severe water scarcity in SEU) – Population decline in continental Europe but population increase in France; overall reduced number of people exposed at 1.8°C; Water Exploitation index increase of 60–200% – Low risk today (according to D&A) Literature supporting this transition: Blauhut et al. (2015); Kebede et al. (2015); Blauhut et al. (2016); Gain et al. (2016); Schleussner et al. (2016); Stahl et al. (2016); Bisselink et al. (2018); Byers et al. (2018); Naumann et al. (2018); Tobin et al. (2018a); Arnell et al. (2019); Harrison et al. (2019); Koutroulis et al. (2019); Bisselink et al. (2020); Cammalleri et al. (2020); Teotónio et al. (2020); Kebede et al. (2021); Naumann et al. (2021)
Moderate to high risk	2–3	medium	– At 2°C, 16% of population exposed to at least moderate water stress – At 2°C, 7–10 months of drought – At 3°C, 8.8–14 months of drought – At 2°C and 2.5°C, a significant number of people exposed to drought in continental Europe and EEU – At 3°C, significant drought losses also in WCE (50% of European GDP losses), particularly in Atlantic region Literature supporting this transition: Kebede et al. (2015); Schleussner et al. (2016); Bisselink et al. (2018); Byers et al. (2018); Naumann et al. (2018); Tobin et al. (2018a); Arnell et al. (2019); Harrison et al. (2019); Koutroulis et al. (2019); Bisselink et al. (2020); Cammalleri et al. (2020); Teotónio et al. (2020); Kebede et al. (2021); Naumann et al. (2021)
High to very high risk	4–4.5	low	– More strongly at 4°C GWL, a significant number of people exposed to drought in continental Europe and EEU – At 4.5°C GWL, Water Exploitation index increase of 200–310% Literature supporting this transition: Kebede et al. (2015); Bisselink et al. (2018); Byers et al. (2018); Naumann et al. (2018); Tobin et al. (2018a); Arnell et al. (2019); Harrison et al. (2019); Koutroulis et al. (2019); Bisselink et al. (2020); Cammalleri et al. (2020); Teotónio et al. (2020); Kebede et al. (2021); Naumann et al. (2021)
Level of risk due to water scarcity in WCE with high adaptation			
Undetectable to moderate risk	1.5–2.5	low	– There is high potential of water-efficiency improvements and water savings (>25% of damage avoided) in 50% of scenario improvements compared with baseline. – There is some adaptation deficit now that can be addressed (assuming an increase of 0.5°C GWL). Literature supporting this transition: Collet et al. (2015); van Vuuren et al. (2015); Koopman et al. (2017); Greve et al. (2018); Manouseli et al. (2018); Garnier and Holman (2019); Papadimitriou et al. (2019); and papers used in the F&E assessment (Table SM.13.1)
Moderate to high risk	3–4	low	– There is considerable potential for investment in large water infrastructure and advanced technologies (including storage), water transfer, water recycling and reuse, and desalination. – As there is less of such infrastructure in place compared with SEU, there is high potential and effectiveness – (assuming an increase of 1°C GWL). Literature supporting this transition: Collet et al. (2015); van Vuuren et al. (2015); Koopman et al. (2017); Greve et al. (2018); Manouseli et al. (2018); Garnier and Holman (2019); Papadimitriou et al. (2019); and papers used in the F&E assessment (Table SM.13.1)
High to very high risk	Not reached	n.a.	

Table SM13.30 | Transition and confidence levels for the risks of people, economies and infrastructures due to coastal flooding (Figures 13.28, 13.32)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
Level of risk due to coastal flooding with low to medium adaptation (keeping coastal protection as it is now)			
Undetectable to moderate risk	0.8–1.5	<i>medium</i>	Based on the D&A assessment in Figure 13.27, coastal impacts of climate change are starting to be detected in Europe (see Box 13.1), consistent with SROCC (IPCC, 2019).
Moderate to high risk	1.5–2	<i>medium</i>	Expected annual damage is projected to rise by a factor of at least 20 for 1.5–2.1°C GWL in EU-27 plus the UK (Vousdoukas et al., 2018); T values bounded by upper scenarios; supported by references in Sections 13.2 and 13.6; consistent with AR5 (Kovats et al., 2014) and SROCC (IPCC, 2019).
High to very high risk	2–3	<i>medium</i>	Expected annual damage is projected to rise by two orders of magnitude in EU-27 plus the UK above 2–3°C GWL (Vousdoukas et al., 2018; Vousdoukas et al., 2020); supported by references in Sections 13.2 and 13.6; consistent with AR5 (Kovats et al., 2014) and SROCC (IPCC, 2019).
Level of risk due to coastal flooding with high adaptation			
Undetectable to moderate risk	0.8–1.5	<i>medium</i>	Based on the D&A assessment in Figure 13.27 and the F&E assessment in Figure 13.11 and Table SM13.1; coastal impacts of climate change are starting to be detected in Europe (see Box 13.1); consistent with SROCC.
Moderate to high risk	1.5–3	<i>medium</i>	Expected annual damage is projected to rise by a factor of 10 above 2°C and by a factor of 20 above 3°C in EU-27 plus the UK (Vousdoukas et al., 2018; Vousdoukas et al., 2020). Transboundary risks can be limited with high adaptation (Mandel et al., 2021). See also Sections 13.2 and 13.6, Table SM 13.1 and references therein. Data are consistent with AR5 (Kovats et al., 2014) and SROCC (IPCC, 2019).
High to very high risk	n.a.	n.a.	Does not reach this threshold
Level of risk from delayed impacts of SLR to cultural heritage and long-living infrastructure			
Undetectable to moderate	0.5–1.2	<i>low</i>	Except in Fennoscandia, European sea levels are projected to rise at rates and magnitudes close to the global average (Fox-Kemper et al., 2021), including Mediterranean Sea levels (SMCCP4.2, SMCCP4.4). Pre-industrial temperatures led to steady sea levels (Fox-Kemper et al., 2021), and historical greenhouse gas emissions are projected 0.7 to 1.1 m of SLR by 2300 (Fox-Kemper et al., 2021), based on Nauels (2019). Compared with other regions, Europe and particularly SEU is characterised by a very high number of UNESCO World Heritage sites exposed to SLR (Marzeion and Levermann, 2014; Reimann et al., 2018). This includes Venice, where high-tide flooding has increased consistently with relative sea level changes (see Box 13.1).
Moderate to high	1.2–1.5	<i>low</i>	Sea levels are projected to rise between 0.3 and 3.1 m by 2300 under SSP1–2.6 (<i>low confidence</i>) (Fox-Kemper et al., 2021). Sea levels are projected to rise by 2–3 m after 2000 years, and by 6–7 m after 10,000 years, for 1.5°C GWL, but these long-term projections incorporate processes in which there is <i>low confidence</i> (Fox-Kemper et al., 2021). The number of UNESCO World Heritage sites exposed to flooding (erosion) increases with SLR, and so do flood frequencies, intensities and erosion rates (Reimann et al., 2018).
High to very high	1.5–2	<i>medium</i>	Sea levels are projected to rise between 0.3 and 3.1 m by 2300 under SSP1–2.6 (<i>low confidence</i>) (Fox-Kemper et al., 2021). Sea levels are projected to rise by 2–6 m after 2000 years, and by 8–13 m after 10,000 years, for 2°C GWL, but these long-term projections incorporate processes in which there is <i>low confidence</i> (Fox-Kemper et al., 2021). In at least 12 countries in Europe, stabilisation of global warming at about 2°C would lead to drowning areas, after millennia, where at least 10% of the population currently live (Clark et al., 2016).

Table SM13.31 | Transition and confidence levels for the risks of people, economies and infrastructures due to inland flooding (Figures 13.28, 13.32)

Risk level	Range of temperature transitions (°C GWL)	Confidence level for the transition	Explanation and references
Level of risk due to inland flooding with low adaptation			
Undetectable to moderate risk	0.5–1.3	high	The same references were used as for Figure 13.27 (Table SM13.22); additional evidence from the trends in the hazards (pluvial floods) (Ranasinghe et al., 2021; Seneviratne et al., 2021) and based on independent expert assessment of the literature of two lead authors and additional support of two contributing authors. The upper value for this transition is 1.3°C GWL because it is within the likely range for observed global warming from pre-industrial levels (see Cross-Chapter Box 2.3 in Gulev et al., 2021) and to distinguish from the impacts at 1.5°C GWL, which have been assessed to be well within the moderate risk level. Namely, the lead authors considered the potential for severe consequences to happen sooner rather than later (i.e., before reaching 1.5°C) given the persistence of certain conditions in the hazards (Ranasinghe et al., 2021; Seneviratne et al., 2021) and exposure (high population density along river banks and urban infrastructures difficult to change). For example, 22% of flood damage losses were in Europe (second after Asia) between 1976 and 2005 (Dottori et al., 2018); 8–26 billion EUR in economic losses; affected potential population per year: 156–679 10⁴ (Alfieri et al., 2018); and at 1°C GWL, the population exposed to 100-year return discharge increases by 3–30% in some grid cells in WCE and NEU (Lange et al., 2020). However, there is also evidence that vulnerability is decreasing. For example, there has been an increase in annually inundated area and the number of people affected since 1870 but no significant trend in normalised financial losses from 1970 to 2006; substantial decline in fatalities since 1950; contrast between NEU and SEU; and increased hazard but decreased vulnerability of population and assets (Paprotny et al., 2018).
Moderate to high risk	1.5–2.5	medium	The level of risk is based on quantitative flood hazard projections (Ranasinghe et al., 2021), flood risk projections (Ciscar et al., 2014; Jongman et al., 2014; Alfieri et al., 2017; Ciscar et al., 2018; Arnell et al., 2019; Harrison et al., 2019; Dottori et al., 2020; Hosseinzadehtalaei et al., 2020; Lange et al., 2020; Merz et al., 2021), considering the potential for cascading risks as part of the criteria for key risk severity (Chapter 16) and considering that limited adaptation has put human lives at risk from flooding.
High to very high risk	2.5–3	medium	The level of risk is based on projections suggesting that flood hazard and risks continue to increase above 3°C GWL (Sections 13.2, 13.6; Alfieri et al., 2017; Hosseinzadehtalaei et al., 2020) and considering the limited ability of current systems to protect and accommodate flooding. At 3°C GWL, evidence suggests that the magnitudes of severe consequences increase considerably, and such consequences are also more widespread for both economic damage and people affected. Therefore, very high risks are plausible beyond 3°C GWL (e.g., at 3°C GWL, around 600–700 billion EUR in damage in WCE compared with approximately 100 billion EUR in the baseline). We also observe approximate quadrupling in part of Russia (Merz et al., 2021). Furthermore, under SSP5 welfare losses (compared with SSP5 baseline without climate change), WCE is –0.17% and Russia is –0.52%; however, indirect losses are increasing more strongly than direct losses between 1.5°C and 3°C GWL because of the persistence of the damage to the economy (Dottori et al., 2018). For Europe in total, there is a slight increase in the population exposed; the difference between 1.5°C and 3°C GWL is not significant (Dottori et al., 2018). At 4°C GWL, there is an 87% increase in pluvial flood risks (Hosseinzadehtalaei et al., 2020). There is <i>medium confidence</i>, however, given that the evidence finds population exposed to riverine flooding to increase strongly (up to 300%) in NEU, as well as in central and western parts of WCE, but to decrease in eastern parts of WCE. There is also a slight increase in SEU.
Level of risk due to inland flooding with high adaptation			
Undetectable to moderate risk	0.5–1.5	medium	The same references were used as for Figure 13.6 (Table SM13.1); additional evidence from the trends in the hazards (pluvial floods) (Ranasinghe et al., 2021; Seneviratne et al., 2021). The higher transition reflects higher adaptation potential (e.g., 1.5°C GWL under RCP2.6 and SSP1/SSP4), and the population exposed to coastal and riverine flooding decreases by >25% in alpine, northern, continental and southern Europe and increases or stays constant on the Atlantic coast (Papadimitriou et al., 2019). Up to around 1.6°C GWL, increasing flood protection levels in all basins to a minimum of 1 per 100 years would decrease the total expected annual flood losses by 30% (A1B-2050) (Jongman et al., 2014). If insurance penetration increases from 30 to 50%, uninsured losses would drop by 60% (Jongman et al., 2014).
Moderate to high risk	2.5–3	medium	The level of risk is based on quantitative flood risk projections under SSP5 (Alfieri et al., 2017; Arnell et al., 2019; Harrison et al., 2019), considering that flood damage in Europe could be offset by adaptation (Jongman et al., 2014; Alfieri et al., 2016a; Dottori et al., 2020), that city drainage systems are difficult to upgrade and the empirical evidence that city adaptation currently remains slow (Ürge-Vorsatz et al., 2018). Under 3°C GWL, the same number of people would be exposed but with a 12% reduction in damage (i.e., river flood) using building-scale measures (Dottori et al., 2020). If we approximate high adaptation under SSP1 conditions, at 4°C GWL, there would be 14–23% less increase in pluvial flood risks under RCP4.5/SSP1 compared with RCP8.5/SSP5 (where the increases is 87%) (Hosseinzadehtalaei et al., 2020); however, the residual risk still would be considerable. Furthermore, there is persistence of hazard and exposure conditions as well as limitations in drainage systems (Dale et al., 2018; Dale, 2021). Additional evidence is given in the F&E assessment (Table SM13.1).
High to very high risk	n.a.	n.a.	Does not reach this threshold

SM13.11 Supplementary Material Supporting the Development of Adaptation Pathways

This section describes the approach used to derive the illustrative adaptation pathways underpinning Figures 13.29–13.32. Adaptation pathways are established in scientific literature on climate adaptation planning to support decision making under deep uncertainty and break adaptation into manageable steps over time. Adaptation pathways map possible sequencing of adaptation options (or portfolio of options or strategies) as a function of global warming (global surface air temperature) and/or time. The basis for deriving adaptation pathways in Chapter 13 follows the methods described in Haasnoot (2013) and Haasnoot (2019). The adaptation pathways are built on the results of the burning embers developed for the key risks (Section 13.10.2), as well as selection of adaptation options and the assessment of their feasibility and effectiveness in Sections 13.2.2, 13.5.2, 13.6.2 and 13.7.2.

Methodology for Deriving the Illustrative Pathways in Chapter 13

Step 1: Decide on the scope and narratives of the adaptation pathways.

In step 1, the author team developed the pathway narrative based on the key risk aggregation (Section 13.10.2) and the aggregation of the options in the feasibility and effectiveness (F&E) assessments described in Tables SM13.1, SM13.5, SM13.15 and SM13.19. Through discussions, the teams decided on the type of illustrative pathways for the different key risks, for example, for each of the European regions (SEU, NEU, EEU and WCE) or hazard type (e.g., coastal and riverine flood). Based on the burning embers and the F&E assessments, suitable combinations of measures and their sequencing were agreed and translated into narratives supporting the pathways. Only measures with at least *medium confidence* were considered. Each statement has full traceability to the F&E assessment tables in the Supplementary Material.

Step 2: Construct the pathways.

The available information on effectiveness to reduce risk per GWL or time was used when available. In the absence of such information, evidence was used from adaptation options already implemented that have demonstrated medium to high effectiveness in reducing risk. Based on the inventory of adaptation options, the relative effectiveness (e.g., A is better than B in reducing risk) was determined.

Path dependencies were identified during discussions with the team (e.g., if option A is implemented then option B is not possible, difficult to implement or not logical). Storylines and narratives were used to describe the implementation of adaptation options over time (e.g., first option A, then option B, and if risk increases further, then C or D is a long-term option). Changes in GWLs affecting these options, the path dependency of options and feasibility of implementation were considered. Pathways typically started with low-regret options that allowed adding more adaptation options and reducing risk under different levels of warming.

Statements developed in step 1 were aggregated to facilitate graphic visualisation, and to be presented in the text. Confidence levels were assigned to the statements. Several intermediate versions of the pathways were drafted and then discussed with the author team until consensus was reached. When the effectiveness of measures was dependent on the region or the hazard, different pathways maps were designed, for example for Key Risk 1 (heat and human health). All pathways are visually presented using the following logic:

- Measures belonging to a similar strategy (e.g., protect or resist) or similar type of measures (e.g., engineering measures) are presented close to each other in the figure.
- The current situation is indicated in the centre.
- Adaptation measures that reduce risk only at low GWLs (thus low effectiveness) are shown close to the present situation, and options with higher effectiveness at higher GWLs are either on the top or bottom of the figure, as such pathways typically start in the centre of the pathway map (close to the present) and move towards the outer corners.
- When two or more measures are needed to reduce the risk, their respective lines are joined.

References

- Aaheim, A., T. Wei and B. Romstad, 2017: Conflicts of economic interests by limiting global warming to +3°C. *Mitig. Adapt. Strateg. Glob. Chang.*, **22**(8), 1131–1148, doi:10.1007/s11027-016-9718-8.
- Acevedo, P., et al., 2010: A broad assessment of factors determining culicoides imicola abundance: modelling the present and forecasting its future in climate change scenarios. *PLoS ONE*, **5**(12), e14236–e14236.
- Adegoji, T., et al., 2019: Making Birmingham a flood resilient city: challenges and opportunities. *Water*, **11**(8), 1699.
- Adler-Nissen, R., 2016: Towards a practice turn in EU studies: the everyday of European integration. *J. Common Mark. Stud.*, **54**(1), 87–103, doi:10.1111/jcms.12329.
- Adynkiewicz-Piragas, M. and B. Miszuk, 2020: Risk analysis related to impact of climate change on water resources and hydropower production in the Lusatian Neisse river basin. *Sustainability*, **12**(12), doi:10.3390/su12125060.
- Aguiñaldo, M.E.C., T. Daddi, M. Hamza and F. Gasbarro, 2019: Climate change perspectives and adaptation strategies of business enterprises: a case study from Italy. *Int. J. Sustain. Dev. World Ecol.*, **26**(2), 129–140, doi:10.1080/13504509.2018.1528571.
- Akentieva, E.M., G.I. Sidoenko and G.A. Tyusov, 2014: To assess the impact of observed and expected future climate changes on the hydropower potential of the regions of the Russian Federation. *Work. A.i. Voeykov Main Geophys. Obs.*, **570**, 95–105.
- Albouy, C., et al., 2014: From projected species distribution to food-web structure under climate change. *Glob. Change Biol.*, **20**(3), 730–741, doi:10.1111/gcb.12467.
- Alessandrini, J.M., J. Ribéron and D. Da Silva, 2019: Will naturally ventilated dwellings remain safe during heatwaves? *Energy Build.*, **183**, 408–417, doi:10.1016/j.enbuild.2018.10.033.
- Alexander, P., et al., 2018: Adaptation of global land use and management intensity to changes in climate and atmospheric carbon dioxide. *Glob. Change Biol.*, **24**(7), 2791–2809, doi:10.1111/gcb.14110.
- Alfieri, L., et al., 2017: Global projections of river flood risk in a warmer world. *Earth's Future*, **5**(2), 171–182, doi:10.1002/2016EF000485.
- Alfieri, L., P. Burek, L. Feyen and G. Forzieri, 2015a: Global warming increases the frequency of river floods in Europe. *Hydrol. Earth Syst. Sci.*, **19**(5), 2247–2260, doi:10.5194/hess-19-2247-2015.
- Alfieri, L., et al., 2018: Multi-model projections of river flood risk in Europe under global warming. *Climate*, **6**(1), doi:10.3390/cli6010016.
- Alfieri, L., L. Feyen and G. Di Baldassarre, 2016a: Increasing flood risk under climate change: a pan-European assessment of the benefits of four adaptation strategies. *Clim. Chang.*, **136**(3-4), 507–521, doi:10.1007/s10584-016-1641-1.
- Alfieri, L., L. Feyen, F. Dottori and A. Bianchi, 2015b: Ensemble flood risk assessment in Europe under high end climate scenarios. *Glob. Environ. Change Hum. Policy Dimens.*, **35**, 199–212, doi:10.1016/j.gloenvcha.2015.09.004.
- Alfieri, L., et al., 2016b: Modelling the socio-economic impact of river floods in Europe. *Nat. Hazards Earth Syst. Sci.*, **16**(6), 1401–1411, doi:10.5194/nhess-16-1401-2016.
- Alkayal, E., M. Bogurcu, F. Ulutas and G.N. Demirel, 2015: Adaptation to climate change in industry: improving resource efficiency through sustainable production applications. *Water. Environ. Res.*, **87**(1), 14–25, doi:10.2175/106143014x14062131178952.
- Alkishe, A.A., A.T. Peterson and A.M. Samy, 2017: Climate change influences on the potential geographic distribution of the disease vector tick Ixodes ricinus. *PLoS ONE*, **12**(12), doi:10.1371/journal.pone.0189092.
- Allard, C., 2018: The rationale for the duty to consult indigenous peoples: comparative reflections from nordic and Canadian legal contexts. *Arct. Rev. Law Polit.*, **9**(0), doi:10.23865/arctic.v9.729.
- Aldrahlleh, L., et al., 2017: Effects of forest conservation and management on volume growth, harvested amount of timber, carbon stock, and amount of deadwood in Finnish boreal forests under changing climate. *Can. J. For. Res.*, **47**(2), 215–225, doi:10.1139/cjfr-2016-0153.
- Altieri, A.H. and K.B. Gedan, 2015: Climate change and dead zones. *Glob. Change Biol.*, **21**(4), 1395–1406, doi:10.1111/gcb.12754.
- Alvarez, I. and M.N. Lorenzo, 2019: Changes in offshore wind power potential over the Mediterranean Sea using CORDEX projections. *Reg. Environ. Chang.*, **19**(1), 79–88, doi:10.1007/s10113-018-1379-6.
- Amponsah, W., et al., 2018: Integrated high-resolution dataset of high-intensity European and Mediterranean flash floods. *Earth Syst. Sci. Data*, **10**(4), 11, doi:10.5194/essd-10-1783-2018.
- Anaya-Romero, M., et al., 2015: Evaluating soil threats under climate change scenarios in the Andalusia region, southern Spain. *Land Degrad. Dev.*, **26**(5), 441–449, doi:10.1002/ldr.2363.
- Ancillotto, L., et al., 2016: Extraordinary range expansion in a common bat: the potential roles of climate change and urbanisation. *Sci. Nat.*, **103**(3), 15, doi:10.1007/s00114-016-1334-7.
- Andersson-Sköld, Y., et al., 2015: An integrated method for assessing climate-related risks and adaptation alternatives in urban areas. *Clim. Risk Manag.*, **7**, 31–50, doi:10.1016/j.crm.2015.01.003.
- Anghileri, D., et al., 2018: A comparative assessment of the impact of climate change and energy policies on Alpine hydropower. *Water Resour. Res.*, **54**(11), 9144–9161, doi:10.1029/2017wr022289.
- Aparicio, Á., 2017: Transport adaptation policies in Europe: from incremental actions to long-term visions. *Transp. Res. Procedia*, **25**, 3529–3537, doi:10.1016/j.trpro.2017.05.277.
- Arctic Council, 2013: Arctic biodiversity assessment: status and trends in arctic biodiversity. In: *The Conservation of Arctic Flora and Fauna, Akureyri, Iceland* [Barry, T., D. Berteaux and H. Bültmann(eds.)]. ISBN 978-9935431226 674 pp).
- Arnbjerg-Nielsen, K., L. Leonardsen and H. Madsen, 2015: Evaluating adaptation options for urban flooding based on new high-end emission scenario regional climate model simulations. *Clim. Res.*, **64**(1), 73–84, doi:10.3354/cr01299.
- Arnell, N.W., et al., 2019: The global and regional impacts of climate change under representative concentration pathway forcings and shared socioeconomic pathway socioeconomic scenarios. *Environ. Res. Lett.*, **14**(8), 84046, doi:10.1088/1748-9326/ab35a6.
- Arrigo, K.R. and G.L. van Dijken, 2015: Continued increases in Arctic Ocean primary production. *Prog. Oceanogr.*, **136**, 60–70, doi:10.1016/j.pocean.2015.05.002.
- Asse, D., et al., 2018: Warmer winters reduce the advance of tree spring phenology induced by warmer springs in the Alps. *Agric. For. Meteorol.*, **252**, 220–230, doi:10.1016/j.agrformet.2018.01.030.
- Asselman, N.E.M. and F. Klijn, 2016: Making room for rivers: quantification of benefits from a flood risk perspective. *E3S Web Conf.*, **7**, 12001, doi:10.1051/e3sconf/20160712001.
- Assis, J., et al., 2017: Major shifts at the range edge of marine forests: the combined effects of climate changes and limited dispersal. *Sci. Rep.*, **7**(1), 44348–44348, doi:10.1038/srep44348.
- Åström, C., et al., 2017: Vulnerability reduction needed to maintain current burdens of heat-related mortality in a changing climate—magnitude and determinants. *Int. J. Environ. Res. Public Health*, **14**(7), doi:10.3390/ijerph14070741.
- Auerswald, K. and P. Fiener, 2019: Soil organic carbon storage following conversion from cropland to grassland on sites differing in soil drainage and erosion history. *Sci. Total. Environ.*, **661**, 481–491, doi:10.1016/j.scitotenv.2019.01.200.
- Averchenkova, A., et al., 2016: Multinational and large national corporations and climate adaptation: are we asking the right questions? A review of current knowledge and a new research perspective: Multinational and large

- national corporations and climate adaptation. *WIREs. Clim. Change*, **7**(4), 517–536, doi:10.1002/wcc.402.
- Babovic, F. and A. Mijic, 2019: The development of adaptation pathways for the long-term planning of urban drainage systems. *J. Flood Risk Manag.*, **12**(S2), e12538, doi:10.1111/jfr3.12538.
- Babovic, F., A. Mijic and K. Madani, 2018: Decision making under deep uncertainty for adapting urban drainage systems to change. *Urban Water J.*, **15**(6), 552–560, doi:10.1080/1573062X.2018.1529803.
- Bachner, G., 2017: Assessing the economy-wide effects of climate change adaptation options of land transport systems in Austria. *Reg. Environ. Chang.*, **17**(3), 929–940, doi:10.1007/s10113-016-1089-x.
- Bakanev, S.V., 2017: Prospects of snow crab *Chionoecetes opilio* fishery in the Barents Sea. *Fish. Issues*, **18**(3), 268–303.
- Baker-Austin, C., J. Trinanes, N. Gonzalez-Escalona and J. Martinez-Urtaza, 2017: Non-cholera vibrios: the microbial barometer of climate change. *Trends Microbiol.*, **25**(1), 76–84, doi:10.1016/j.tim.2016.09.008.
- Baker-Austin, C., et al., 2013: Emerging *Vibrio* risk at high latitudes in response to ocean warming. *Nat. Clim. Chang.*, **3**(1), 73–77, doi:10.1038/nclimate1628.
- Balint, M., et al., 2011: Cryptic biodiversity loss linked to global climate change. *Nat. Clim. Chang.*, **1**(6), 313–318, doi:10.1038/NCLIMATE1191.
- Ballantyne, A., et al., 2017: Accelerating net terrestrial carbon uptake during the warming hiatus due to reduced respiration. *Nat. Clim. Chang.*, **7**(2), 148–152, doi:10.1038/nclimate3204.
- Bamberg, S., T. Masson, K. Brewitt and N. Nemetschek, 2017: Threat, coping and flood prevention – a meta-analysis. *J. Environ. Psychol.*, **54**, 116–126, doi:10.1016/j.jenvp.2017.08.001.
- Bangash, R.F., et al., 2013: Ecosystem services in Mediterranean river basin: climate change impact on water provisioning and erosion control. *Sci. Total Environ.*, **458–460**, 246–255, doi:10.1016/j.scitotenv.2013.04.025.
- Barbosa, R., R. Vicente and R. Santos, 2015: Climate change and thermal comfort in Southern Europe housing: a case study from Lisbon. *Build. Environ.*, **92**, 440–451, doi:10.1016/j.buildenv.2015.05.019.
- Baron, N. and L.K. Petersen, 2015: Climate change or variable weather: rethinking Danish homeowners' perceptions of floods and climate. *Reg. Environ. Chang.*, **15**(6), 1145–1155, doi:10.1007/s10113-014-0701-1.
- Barredo, J., G. Caudullo and A. Dosio, 2016: Mediterranean habitat loss under future climate conditions: assessing impacts on the Natura 2000 protected area network. *Appl. Geogr.*, **75**, 83–92, doi:10.1016/j.apgeog.2016.08.003.
- Batllo, E., et al., 2020: Forest and woodland replacement patterns following drought-related mortality. *Proc. Natl. Acad. Sci.*, **117**(47), 29720, doi:10.1073/pnas.2002314117.
- Battiston, S., et al., 2017: A climate stress-test of the financial system. *Nat. Clim. Chang.*, **7**(4), 283–288, doi:10.1038/nclimate3255.
- Baudron, A.R., et al., 2020: Changing fish distributions challenge the effective management of European fisheries. *Ecography*, **43**(4), 494–505, doi:10.1111/ecog.04864.
- Bausch, T. and W.C. Gartner, 2020: Winter tourism in the European Alps: is a new paradigm needed? *J. Outdoor Recreat. Tour.*, **31**, 100297, doi:10.1016/j.jort.2020.100297.
- Bausch, T., A. Humpe and S. Gössling, 2019: Does climate change influence guest loyalty at Alpine winter destinations? *Sustainability*, **11**(15), doi:10.3390/su11154233.
- Bedford, J., et al., 2020: Lifeform indicators reveal large-scale shifts in plankton across the North-West European shelf. *Glob. Change Biol.*, **26**(6), 3482–3497, doi:10.1111/gcb.15066.
- Bedia, J., et al., 2018: Seasonal predictions of fire weather index: paving the way for their operational applicability in Mediterranean Europe. *Clim. Serv.*, **9**, 101–110, doi:10.1016/j.cliser.2017.04.001.
- Bedia, J., et al., 2014: Forest fire danger projections in the Mediterranean using ENSEMBLES regional climate change scenarios. *Clim. Chang.*, **122**(1), 185–199, doi:10.1007/s10584-013-1005-z.
- Behrens, P., et al., 2017: Climate change and the vulnerability of electricity generation to water stress in the European Union. *Nat. Energy*, **2**(8), doi:10.1038/nenergy.2017.114.
- Beland Lindahl, K., A. Johansson, A. Zachrisson and R. Viklund, 2018: Competing pathways to sustainability? Exploring conflicts over mine establishments in the Swedish mountain region. *J. Environ. Manag.*, **218**, 402–415, doi:10.1016/j.jenvman.2018.04.063.
- Belonovskaya, E.A., et al., 2016: "Greening" of the Russian arctic and the modern trends of transformation of its biota. *Izv. Akad. Nauk SSSR Ser. Geogr.*, **3**, 28–39.
- Ben-Ari, T., et al., 2018: Causes and implications of the unforeseen 2016 extreme yield loss in the breadbasket of France. *Nat. Commun.*, **9**(1), doi:10.1038/s41467-018-04087-x.
- Benedetti, F., et al., 2019: Climate change may have minor impact on zooplankton functional diversity in the Mediterranean Sea. *Divers. Distrib.*, **25**(4), 568–581, doi:10.1111/ddi.12857.
- Benedetti, F., F. Guilhaumon, F. Adloff and S.-D. Ayata, 2018: Investigating uncertainties in zooplankton composition shifts under climate change scenarios in the Mediterranean Sea. *Ecography*, **41**(2), 345–360, doi:10.1111/ecog.02434.
- Beniston, M., et al., 2018: The European mountain cryosphere: a review of its current state, trends, and future challenges. *Cryosphere*, **12**(2), 759–794, doi:10.5194/tc-12-759-2018.
- Beniston, M. and M. Stoffel, 2016: Rain-on-snow events, floods and climate change in the Alps: events may increase with warming up to 4°C and decrease thereafter. *Sci. Total Environ.*, **571**(May 1999), 228–236, doi:10.1016/j.scitotenv.2016.07.146.
- Berberoglu, S., et al., 2020: Spatial and temporal evaluation of soil erosion in Turkey under climate change scenarios using the Pan-European Soil Erosion Risk Assessment (PESERA) model. *Environ. Monit. Assess.*, **192**(8), 491, doi:10.1007/s10661-020-08429-5.
- Berdnikov, S.V., et al., 2019: Integrated mathematical model of the Barents and White seas large marine ecosystem - a tool for assessing natural risks and efficient use of biological resources. *Доклады Академии Наук*, **487**(5), 566–572, doi:10.31857/s0869-56524875566-572.
- Berghuijs, W.R., S.T. Allen, S. Harrigan and J.W. Kirchner, 2019: Growing spatial scales of synchronous river flooding in Europe. *Geophys. Res. Lett.*, **46**(3), 1423–1428, doi:10.1029/2018GL081883.
- Berlinski, N.A. and Y.I. Popov, 2018: Formation of bottom hypoxia and hydrogen sulfide on the Black Sea shelf. *Visnyk V. N. Karazin Kharkiv Natl. Univ. Ser. "Ecology"*, **18**, 6–13.
- Berrang-Ford, A.R., Siders, A. Lesnikowski, et al., 2021: A global systematic stocktake of evidence on human adaptation to climate change. *Nat. Clim. Chang.*, **11**, 989–1000. doi:10.1038/s41558-021-01170-y
- Berrang-Ford, L., T. Pearce and J.D. Ford, 2015: Systematic review approaches for climate change adaptation research. *Reg. Environ. Chang.*, **15**(5), 755–769, doi:10.1007/s10113-014-0708-7.
- Bertoldi, P., et al., 2020: *Covenant of Mayors: 2019 Assessment*. [JRC (ed.)]. JRC Science for Policy Report, Luxembourg, <https://ec.europa.eu/jrc/en/publication/eur-scientific-and-technical-research-reports/covenant-mayors-2019-assessment>. (63 pp).
- Birchenough, S.N.R., et al., 2015: Climate change and marine benthos: a review of existing research and future directions in the North Atlantic. *WIREs Clim. Chang.*, **6**(2), 203–223, doi:10.1002/wcc.330.
- Bird, D.N., et al., 2016: Modelling climate change impacts on and adaptation strategies for agriculture in Sardinia and Tunisia using AquaCrop and value-at-risk. *Sci. Total Environ.*, **543**, 1019–1027, doi:10.1016/j.scitotenv.2015.07.035.
- Bisbis, M.B., N.S. Gruda and M.M. Blanke, 2019: Securing horticulture in a changing climate – a mini review. *Horticulturae*, **5**(3), doi:10.3390/horticulturae5030056.

- Bisselink, B., et al., 2018: *Impact of a Changing Climate, Land Use, and Water Usage on Europe's Water Resources*, 86–86. Publications Office of the European Union, Luxembourg, ISBN 978-9279802874.
- Bisselink, B., et al., 2020: *Climate Change and Europe's Water Resources*. Publications Office of the European Union, Luxembourg, ISBN 978-9276103981.
- Blagrove, M.S.C., et al., 2020: Potential for Zika virus transmission by mosquitoes in temperate climates. *Proc. R. Soc. B. Biol. Sci.*, **287**(1930), doi:10.1098/rspb.2020.0119.
- Blauhut, V., L. Gudmundsson and K. Stahl, 2015: Towards pan-European drought risk maps: quantifying the link between drought indices and reported drought impacts. *Environ. Res. Lett.*, **10**(1), 14008, doi:10.1088/1748-9326/10/1/014008.
- Blauhut, V., et al., 2016: Estimating drought risk across Europe from reported drought impacts, drought indices, and vulnerability factors. *Hydrol. Earth Syst. Sci.*, **20**(7), 2779–2800, doi:10.5194/hess-20-2779-2016.
- Blöschl, G., et al., 2017: Changing climate shifts timing of European floods. *Science*, **357**(6351), 588–590, doi:10.1126/science.aan2506.
- Blöschl, G., et al., 2019: Changing climate both increases and decreases European river floods. *Nature*, **573**, 108–111.
- Bobretsov, A.V., T.K. Tertitsa and V.P. Teplova, 2019: The impact of climate change on the phenology of plants and animals of the south-eastern part of the Komi Republic (the Pechora-Ilych biosphere reserve). *Probl. Ecol. Monit. Ecosyst. Model.*, **18**(4), 74–93, doi:10.21513/0207-2564-2017-4-74-93.
- Bodoque, J.M., et al., 2019: Enhancing flash flood risk perception and awareness of mitigation actions through risk communication: a pre-post survey design. *J. Hydrol. Reg. Stud.*, **568**, 769–779, doi:10.1016/j.jhydrol.2018.11.007.
- Bogmans, C., G. Dijkema and M. van Vliet, 2017: Adaptation of thermal power plants: the (ir)relevance of climate (change) information. *Energy Econ.*, **62**, 1–18, doi:10.1016/j.eneco.2016.11.012.
- Bokhorst, S., et al., 2016: Changing arctic snow cover: a review of recent developments and assessment of future needs for observations, modelling, and impacts. *Ambio*, **45**(5), 516–537, doi:10.1007/s13280-016-0770-0.
- Bollinger, L.A. and G.P.J. Dijkema, 2016: Evaluating infrastructure resilience to extreme weather – the case of the Dutch electricity transmission network. *Eur. J. Transp. Infrastruct. Res.*, **16**(1), 214–239.
- Bombelli, G.M., A. Soncini, A. Bianchi and D. Bocchiola, 2019: Potentially modified hydropower production under climate change in the Italian Alps. *Hydrol. Process.*, **33**(17), 2355–2372, doi:10.1002/hyp.13473.
- Bondur, R., V. Cloutier, E. Rosa and M. Benzaazoua, 2016: A review and evaluation of the impacts of climate change on geogenic arsenic in groundwater from fractured bedrock aquifers. *Water Air Soil Pollut.*, **227**(9), doi:10.1007/s11270-016-2936-6.
- Bondur, V.G., 2011: Satellite monitoring of wildfires during the anomalous heat wave of 2010 in Russia. *Izv. Atmos. Ocean. Phys.*, **47**(9), 1039–1048, doi:10.1134/S0001433811090040.
- Borrelli, P., et al., 2016: Assessment of the cover changes and the soil loss potential in European forestland: first approach to derive indicators to capture the ecological impacts on soil-related forest ecosystems. *Ecol. Indic.*, **60**, 1208–1220, doi:10.1016/j.ecolind.2015.08.053.
- Borrelli, P., et al., 2020: Land use and climate change impacts on global soil erosion by water (2015–2070). *Proc. Natl. Acad. Sci.*, **117**(36), 21994–22001, doi:10.1073/pnas.2001403117.
- Børshiem, K.Y., 2017: Bacterial and primary production in the Greenland Sea. *J. Mar. Syst.*, **176**, 54–63, doi:10.1016/j.jmarsys.2017.08.003.
- Bosello, F., et al., 2020: *Macroeconomic, Spatially-Resolved Impact Assessment*. CMCC, Venice, Italy.
- Botzen, W.J.W., J.C.J.H. Aerts and J.C.J.M. van den Bergh, 2013: Individual preferences for reducing flood risk to near zero through elevation. *Mitig. Adapt. Strateg. Glob. Chang.*, **18**(2), 229–244, doi:10.1007/s11027-012-9359-5.
- Boucher, M.-A. and M.-H. Ramos, 2018: Ensemble streamflow forecasts for hydropower systems. In: *Handbook of Hydrometeorological Ensemble Forecasting*. Springer, Berlin, pp. 1–19. ISBN 978-3642404573.
- Bouwer, L., et al., 2018: Chapter 4 - Upscaling the impacts of climate change in different sectors and adaptation strategies. In: *Adapting to Climate Change in Europe*. Elsevier, Amsterdam, pp. 173–243. ISBN 978-0128498873.
- Bouzid, M., et al., 2014: Climate change and the emergence of vector-borne diseases in Europe: case study of dengue fever. *BMC Public Health*, **14**, doi:10.1186/1471-2458-14-781.
- Bowler, D.E., et al., 2018: The geography of the Anthropocene differs between the land and the sea. *bioRxiv*, **432880**, doi:10.1101/432880.
- Bowler, D.E., et al., 2015: A cross-taxon analysis of the impact of climate change on abundance trends in central Europe. *Biol. Conserv.*, **187**, 41–50.
- Bowler, D.E., et al., 2019: Long-term declines of European insectivorous bird populations and potential causes. *Conserv. Biol.*, **33**(5), 1120–1130, doi:10.1111/cobi.13307.
- Bowler, D.E., et al., 2017: Cross-realm assessment of climate change impacts on species' abundance trends. *Nat. Ecol. Evol.*, **1**(0067), doi:10.1038/s41559-016-0067.
- Bramanti, L., et al., 2013: Detrimental effects of ocean acidification on the economically important Mediterranean red coral (*Corallium rubrum*). *Glob. Change Biol.*, **19**(6), 1897–1908, doi:10.1111/gcb.12171.
- Brambilla, M., P. Pedrini, A. Rolando and D. Chamberlain, 2016: Climate change will increase the potential conflict between skiing and high-elevation bird species in the Alps. *J. Biogeogr.*, **43**(11), 2299–2309, doi:10.1111/jbi.12796.
- Brännlund, I. and P. Axelsson, 2011: Reindeer management during the colonization of Sami lands: a long-term perspective of vulnerability and adaptation strategies. *Glob. Environ. Chang.*, **21**(3), 1095–1105, doi:10.1016/j.gloenvcha.2011.03.005.
- Brännström, M., 2017: *Skogsbruk och renskötsel på samma mark : en rättsvetenskaplig studie av äganderätten och renskötselrätten*. Umeå University, Umeå.
- Bright, R.M., et al., 2017: Local temperature response to land cover and management change driven by non-radiative processes. *Nat. Clim. Chang.*, **7**, 296, doi:10.1038/nclimate3250. <https://www.nature.com/articles/nclimate3250#supplementary-information>.
- Brodie, J., et al., 2014: The future of the northeast Atlantic benthic flora in a high CO2 world. *Ecol. Evol.*, **4**(13), 2787–2798, doi:10.1002/ecs3.1105.
- Brodrick, T.J., J. Powers, H. Cochard and B. Choat, 2020: Hanging by a thread? Forests and drought. *Science*, **368**(6488), 261–266, doi:10.1126/science.aat7631.
- Brosy, C., K. Zaninovic and A. Matzarakis, 2014: Quantification of climate tourism potential of Croatia based on measured data and regional modeling. *Int. J. Biometeorol.*, **58**(6), 1369–1381, doi:10.1007/s00484-013-0738-8.
- Brugueras, S., et al., 2020: Environmental drivers, climate change and emergent diseases transmitted by mosquitoes and their vectors in southern Europe: a systematic review. *Environ. Res.*, **191**, doi:10.1016/j.envres.2020.110038.
- Brunner, M.I., et al., 2019: Present and future water scarcity in Switzerland: potential for alleviation through reservoirs and lakes. *Sci. Total Environ.*, **666**, 1033–1047, doi:10.1016/j.scitotenv.2019.02.169.
- Bryndum-Buchholz, A., et al., 2020: Climate-change impacts and fisheries management challenges in the North Atlantic Ocean. *Mar. Ecol. Prog. Ser.*, **648**, 1–17.
- Bryndum-Buchholz, A., et al., 2019: Twenty-first-century climate change impacts on marine animal biomass and ecosystem structure across ocean basins. *Glob. Change Biol.*, **25**(2), 459–472, doi:10.1111/gcb.14512.
- Bubeck, P., et al., 2017: Explaining differences in flood management approaches in Europe and in the USA – a comparative analysis. *J. Flood Risk Manag.*, **10**(4), 436–445, doi:10.1111/jfr3.12151.
- Bucak, T., et al., 2017: Future water availability in the largest freshwater Mediterranean lake is at great risk as evidenced from simulations with the SWAT model. *Sci. Total Environ.*, **581–582**, 413–425, doi:10.1016/j.scitotenv.2016.12.149.

- Buelow, F. and N. Cradock-Henry, 2018: What you sow is what you reap? (Dis-)Incentives for adaptation intentions in farming. *Sustainability*, **10**(4), doi:10.3390/su10041133.
- Buonomo, R., et al., 2018: Predicted extinction of unique genetic diversity in marine forests of *Cystoseira* spp. *Mar. Environ. Res.*, **138**, 119–128, doi:10.1016/j.marenvres.2018.04.013.
- Burbidge, R., 2015: Adapting aviation to a changing climate: key priorities for action. *J. Air Transp. Manag.*, **71**, 167–174, doi:10.1016/j.jairtraman.2018.04.004.
- Burke, M., S.M. Hsiang and E. Miguel, 2015: Global non-linear effect of temperature on economic production. *Nature*, **527**(7577), 235–239, doi:10.1038/nature15725.
- Burrows, M.T., et al., 2019: Ocean community warming responses explained by thermal affinities and temperature gradients. *Nat. Clim. Chang.*, **9**(12), 959–963, doi:10.1038/s41558-019-0631-5.
- Buser, M., 2020: Coastal adaptation planning in Fairbourne, Wales: lessons for climate change adaptation. *Plan. Pract. Res.*, **35**(2), 127–147, doi:10.1080/02697459.2019.1696145.
- Buurman, J., M.J.P. Mens and R.J. Dahm, 2017: Strategies for urban drought risk management: a comparison of 10 large cities. *Int. J. Water Resour. Dev.*, **33**(1), 31–50, doi:10.1080/07900627.2016.1138398.
- Byers, E., et al., 2018: Global exposure and vulnerability to multi-sector development and climate change hotspots. *Environ. Res. Lett.*, **13**(5), 55012–55012, doi:10.1088/1748-9326/aabf45.
- Byers, E., et al., 2015: Cooling water for Britain's future electricity supply. *Proc. Inst. Civ. Eng. Energy*, **168**(3), 188–204, doi:10.1680/ener.14.00028.
- Byers, E.A., G. Coxon, J. Freer and J.W. Hall, 2020: Drought and climate change impacts on cooling water shortages and electricity prices in Great Britain. *Nat. Commun.*, **11**(1), 2239, doi:10.1038/s41467-020-16012-2.
- Caffarra, A., et al., 2012: Modelling the impact of climate change on the interaction between grapevine and its pests and pathogens: European grapevine moth and powdery mildew. *Agric. Ecosyst. Environ.*, **148**, 89–101, doi:10.1016/j.agee.2011.11.017.
- Cameron, R.W.F., J.E. Taylor and M.R. Emmett, 2014: What's 'cool' in the world of green façades? How plant choice influences the cooling properties of green walls. *Build. Environ.*, **73**, 198–207, doi:10.1016/j.buildenv.2013.12.005.
- Camia, A., 2017: *Modeling the Impacts of Climate Change on Forest Fire Danger in Europe: Sectorial Results of the PESETA II Project*. Publications Office of the European Union, Luxembourg.
- Camia, A., G. Liberta' and J. San-Miguel-Ayanz, 2017: *Modeling the Impacts of Climate Change on Forest Fire Danger in Europe: Sectorial Results of the PESETA II Project*. Publications Office of the European Union, Luxembourg.
- Caminade, C., et al., 2014: Impact of climate change on global malaria distribution. *Proc. Natl. Acad. Sci. U.S.A.*, **111**(9), 3286–3291, doi:10.1073/pnas.1302089111.
- Caminade, C., et al., 2012: Suitability of European climate for the Asian tiger mosquito *Aedes albopictus*: recent trends and future scenarios. *J. R. Soc. Interface.*, **9**(75), 2708–2717, doi:10.1098/rsif.2012.0138.
- Caminade, C., et al., 2017: Global risk model for vector-borne transmission of Zika virus reveals the role of El Niño 2015. *Proc. Natl. Acad. Sci. U.S.A.*, **114**(1), 119–124, doi:10.1073/pnas.1614303114.
- Cammalleri, C., et al., 2020: *Global Warming and Drought Impacts in the EU*. Publications Office of the European Union, Luxembourg, doi:10.2760/597045.
- Campos, I.S., et al., 2016: Climate adaptation, transitions, and socially innovative action-research approaches. *Ecol. Soc.*, **21**(1), doi:10.5751/ES-08059-210113.
- Campos Rodrigues, L., J. Freire-González, A. González Puig and I. Puig-Ventosa, 2018: Climate change adaptation of Alpine ski tourism in Spain. *Climate*, **6**(2), doi:10.3390/cli6020029.
- Canadell, J. and R.B. Jackson, 2021: *Ecosystem Collapse and Climate Change*. Springer, Heidelberg, New York.
- Capuzzo, E., et al., 2018: A decline in primary production in the North Sea over 25 years, associated with reductions in zooplankton abundance and fish stock recruitment. *Glob. Change Biol.*, **24**(1), e352–e364, doi:10.1111/gcb.13916.
- Carmona, R., et al., 2016: Mortality attributable to extreme temperatures in Spain: a comparative analysis by city. *Environ. Int.*, **91**, 22–28, doi:10.1016/j.envint.2016.02.018.
- Carnicer, J., et al., 2013: Contrasting trait syndromes in angiosperms and conifers are associated with different responses of tree growth to temperature on a large scale. *Front. Plant Sci.*, **4**, 409.
- Carnicer, J., et al., 2011: Widespread crown condition decline, food web disruption, and amplified tree mortality with increased climate change-type drought. *Proc. Natl. Acad. Sci.*, **108**(4), 1474–1478, doi:10.1073/pnas.1010070108.
- Carnicer, J., et al., 2019a: Regime shifts of Mediterranean forest carbon uptake and reduced resilience driven by multidecadal ocean surface temperatures. *Glob. Change Biol.*, **25**(8), 2825–2840, doi:10.1111/gcb.14664.
- Carnicer, J., et al., 2019b: Phenotypic biomarkers of climatic impacts on declining insect populations: a key role for decadal drought, thermal buffering and amplification effects and host plant dynamics. *J. Anim. Ecol.*, **88**(3), 376–391, doi:10.1111/1365-2656.12933.
- Carnicer, J., et al., 2021: Forest resilience to global warming is strongly modulated by local-scale topographic, microclimatic and biotic conditions. *J. Ecol.*, (n/a), doi:10.1111/1365-2745.13752. n/a.
- Carnielo, E. and M. Zinzi, 2013: Optical and thermal characterisation of cool asphalts to mitigate urban temperatures and building cooling demand. *Build. Environ.*, **60**, 56–65, doi:10.1016/j.buildenv.2012.11.004.
- Carozza, D.A., D. Bianchi and E.D. Galbraith, 2019: Metabolic impacts of climate change on marine ecosystems: implications for fish communities and fisheries. *Global Ecol. Biogeogr.*, **28**(2), 158–169, doi:10.1111/geb.12832.
- Carstensen, J., J.H. Andersen, B.G. Gustafsson and D.J. Conley, 2014: Deoxygenation of the Baltic Sea during the last century. *Proc. Natl. Acad. Sci.*, **201323156**, doi:10.1073/pnas.1323156111.
- Carvalho, L.G., et al., 2013: Species richness declines and biotic homogenisation have slowed down for NW-European pollinators and plants. *Ecol. Lett.*, **16**(7), 870–878, doi:10.1111/ele.12121.
- Carvalho-Santos, C., et al., 2016: Assessing the effects of land cover and future climate conditions on the provision of hydrological services in a medium-sized watershed of Portugal: impacts of land cover and future climate on hydrological services. *Hydrol. Process.*, **30**(5), 720–738, doi:10.1002/hyp.10621.
- Carvalho, D., et al., 2017a: Urban resilience to future urban heat waves under a climate change scenario: A case study for Porto urban area (Portugal). *Urban Clim.*, **19**, 1–27, doi:10.1016/j.uclim.2016.11.005.
- Carvalho, D., A. Rocha, M. Gomez-Gesteira and C. Santos, 2017b: Potential impacts of climate change on European wind energy resource under the CMIP5 future climate projections. *Renew. Energy*, **101**, 29–40, doi:10.1016/j.renene.2016.08.036.
- Casado-Amezúa, P., et al., 2019: Distributional shifts of canopy-forming seaweeds from the Atlantic coast of Southern Europe. *Biodivers. Conserv.*, **28**(5), 1151–1172, doi:10.1007/s10531-019-01716-9.
- Casanueva, A., et al., 2019: Overview of existing heat-health warning systems in Europe. *Int. J. Environ. Res. Public Health*, **16**(15), doi:10.3390/ijerph16152657.
- Casanueva, A., et al., 2020: Escalating environmental summer heat exposure—a future threat for the European workforce. *Reg. Environ. Chang.*, **20**(2), 40, doi:10.1007/s10113-020-01625-6.
- Castellanos-Frias, E., D. Garcia De Leon, F. Bastida and J.L. Gonzalez-Andujar, 2016: Predicting global geographical distribution of *Lolium rigidum* (rigid ryegrass) under climate change. *J. Agric. Sci.*, **154**(5), 755–764, doi:10.1017/S0021859615000799.
- Castelle, B., et al., 2018: Spatial and temporal patterns of shoreline change of a 280-km high-energy disrupted sandy coast from 1950 to 2014: SW France. *Estuar. Coast. Shelf Sci.*, **200**, 212–223, doi:10.1016/j.ecss.2017.11.005.

- CDSB, 2020: *The State of EU Environmental Disclosure in 2020*. Climate Disclosure Standards Board (CDSB), London, UK, <https://www.cdsb.net/nfrd2020>. Accessed 2021.
- Ceglar, A., M. Turco, A. Toreti and F.J. Doblas-Reyes, 2017: Linking crop yield anomalies to large-scale atmospheric circulation in Europe. *Agric. For. Meteorol.*, **240–241**, 35–45, doi:10.1016/j.agrformet.2017.03.019.
- Ceglar, A., M. Zampieri, A. Toreti and F. Dentener, 2019: Observed northward migration of agro-climate zones in Europe will further accelerate under climate change. *Earths Future*, **7**(9), 1088–1101, doi:10.1029/2019EF001178.
- Cellura, M., F. Guarino, S. Longo and G. Tumminia, 2018: Climate change and the building sector: modelling and energy implications to an office building in Southern Europe. *Energy Sustain. Dev.*, **45**, 46–65, doi:10.1016/j.esd.2018.05.001.
- Chefaoui, R.M., et al., 2019: Integrating reproductive phenology in ecological niche models changed the predicted future ranges of a marine invader. *Divers. Distrib.*, **25**(5), 688–700, doi:10.1111/ddi.12910.
- Chen, I.C., et al., 2011: Rapid range shifts of species associated with high levels of climate warming. *Science*, **333**(6045), 1024, doi:10.1126/science.1206432.
- Chen, L., et al., 2018: Spring phenology at different altitudes is becoming more uniform under global warming in Europe. *Glob. Change Biol.*, **24**(9), 3969–3975, doi:10.1111/gcb.14288.
- Chen, Y., et al., 2019: Economic losses of carbon emissions from circum-arctic permafrost regions under RCP-SSP scenarios. *Sci. Total Environ.*, **658**, 1064–1068, doi:10.1016/j.scitotenv.2018.12.299.
- Chergui, B., 2019: Socioeconomic factors drive fire-regime variability in the Mediterranean basin. *Ecosystems*, **21**, 1–10, doi:10.1007/s10021-017-0172-6.
- Cheung, W., et al., 2016: Structural uncertainty in projecting global fisheries catches under climate change. *Ecol. Model.*, **325**, 57–66, doi:10.1016/j.ecolmodel.2015.12.018.
- Chivers, W.J., A.W. Walne and G.C. Hays, 2017: Mismatch between marine plankton range movements and the velocity of climate change. *Nat. Commun.*, **8**, 14434, doi:10.1038/ncomms14434.
- Chizhikova, N., 2018: Dynamics of intense rainfalls in the southern half of European Russia for the period 1960–2015. *IOP Conf. Ser. Earth Environ. Sci.*, **107**, 12098–12098, doi:10.1088/1755-1315/107/1/012098.
- Ciais, P., et al., 2005: Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature*, **437**(7058), 529–533, doi:10.1038/nature03972.
- Ciais, P., et al., 2019: Five decades of northern land carbon uptake revealed by the interhemispheric CO₂ gradient. *Nature*, **568**(7751), 221–225, doi:10.1038/s41586-019-1078-6.
- Ciampalini, R., et al., 2020: Modelling soil erosion responses to climate change in three catchments of Great Britain. *Sci. Total Environ.*, **749**, 141657, doi:10.1016/j.scitotenv.2020.141657.
- Ciasca, F., A. Sallati and N. Tolu (eds.), 2017: *Italian National Resilience Plan 2017: For a More Reliable Grid*. 2017 AEIT International Annual Conference, 20–22 Sept., 1–5.
- Cilek, A., et al., 2015: Erosion modelling in a mediterranean subcatchment under climate change scenarios using Pan-European Soil Erosion Risk Assessment (PESERA). *Int. Arch. Photogramm. Remote. Sens. Spatial Inf. Sci. ISR Arch.*, **40**(7W3), 359–365, doi:10.5194/isprsarchives-XL-7-W3-359-2015.
- Ciscar, J.C., et al., 2018: *Climate Impacts in Europe: Final Report of the JRC PESETA III Project*. Joint Research Centre (JRC) Science for Policy report. Joint Research Centre (JRC), Publications Office of the European Union, Luxembourg, 95 pp.
- Ciscar, J.-C., et al., 2014: *Climate Impacts in Europe – The JRC PESETA II Project*. EUR – Scientific and Technical Research, Vol. 26586, 155.
- Clark, N.J., J.T. Kerry and C.I. Fraser, 2020: Rapid winter warming could disrupt coastal marine fish community structure. *Nat. Clim. Chang.*, **10**(9), 862–867, doi:10.1038/s41558-020-0838-5.
- Clark, P.U., et al., 2016: Consequences of twenty-first-century policy for multi-millennial climate and sea-level change. *Nat. Clim. Chang.*, **6**(4), 360–369, doi:10.1038/nclimate2923.
- Cohen, J., K. Moeltner, J. Reichl and M. Schmidhalter, 2018a: Effect of global warming on willingness to pay for uninterrupted electricity supply in European nations. *Nat. Energy*, **3**(1), 37–45, doi:10.1038/s41560-017-0045-4.
- Cohen, J.M., M.J. Lajeunesse and J.R. Rohr, 2018b: A global synthesis of animal phenological responses to climate change. *Nat. Clim. Chang.*, **8**(3), 224–228, doi:10.1038/s41558-018-0067-3.
- Coll, J., et al., 2016: Projected climate change impacts on upland heaths in Ireland. *Clim. Res.*, **69**(2), 177–191.
- Coll, M., et al., 2019: Who is to blame? Plausible pressures on small pelagic fish population changes in the northwestern Mediterranean Sea. *Mar. Ecol. Prog. Ser.*, **2018**, 277–294, doi:10.3354/meps12591.
- Coll, M., et al., 2013: Multivariate effect gradients driving forest demographic responses in the Iberian Peninsula. *For. Ecol. Manag.*, **303**, 195–209, doi:10.1016/j.foreco.2013.04.010.
- Collet, L., et al., 2015: Water supply sustainability and adaptation strategies under anthropogenic and climatic changes of a meso-scale mediterranean catchment. *Sci. Total Environ.*, **536**, 589–602, doi:10.1016/j.scitotenv.2015.07.093.
- Colon-Gonzalez, F.J., et al., 2021: Projecting the risk of mosquito-borne diseases in a warmer and more populated world: a multi-model, multi-scenario intercomparison modelling study. *Lancet Planet. Health*, **5**(7), E404–E414.
- CoM, 2019: *Covenant in Figures, Covenant of Mayors for Climate & Energy*. <Go to WoS>://WOS:000459735100001. Accessed 2021.
- Coma, R., et al., 2009: Global warming-enhanced stratification and mass mortality events in the Mediterranean. *Proc. Natl. Acad. Sci. U.S.A.*, **106**(15), 6176–6181, doi:10.1073/pnas.0805801106.
- Corrales, X., et al., 2018: Future scenarios of marine resources and ecosystem conditions in the Eastern Mediterranean under the impacts of fishing, alien species and sea warming. *Sci. Rep.*, **8**, 1–16, doi:10.1038/s41598-018-32666-x.
- Costa, H., 2020: *European Wildfire Danger and Vulnerability in a Changing Climate: Towards Integrating Risk Dimensions*. JRC PESETA IV Project Task 9 - Forest Fires. Publication Office of the European Union, Luxembourg.
- Costa, J.M., et al., 2019a: Opportunities and limitations of crop phenotyping in southern European countries. *Front. Plant Sci.*, **10**(September), 1–16, doi:10.3389/fpls.2019.01125.
- Costa, J.M., et al., 2019b: Opportunities and limitations of crop phenotyping in southern European countries. *Front. Plant Sci.*, **10**, 1–16, doi:10.3389/fpls.2019.01125.
- Costa, J.M., et al., 2016: Modern viticulture in southern Europe: vulnerabilities and strategies for adaptation to water scarcity. *Agric. Water Manag.*, **164**, 5–18, doi:10.1016/j.agwat.2015.08.021.
- Cox, B., et al., 2016: Mortality related to cold and heat. What do we learn from dairy cattle? *Environ. Res.*, **149**, 231–238, doi:10.1016/j.envres.2016.05.018.
- Cozzi, S., et al., 2019: Flow regime and nutrient-loading trends from the largest south European watersheds: implications for the productivity of mediterranean and black sea's coastal areas. *Water*, **11**(1), doi:10.3390/w11010001.
- Crespo, D., et al., 2019: Tradeoffs between water uses and environmental flows: a hydroeconomic analysis in the Ebro basin. *Water Resour. Manag.*, **33**(7), 2301–2317, doi:10.1007/s11269-019-02254-3.
- Cronin, J., F. Zabel, O. Dessens and G. Anandarajah, 2020: Land suitability for energy crops under scenarios of climate change and land-use. *Glob. Change Biol. Bioenergy*, **12**(8), 648–665, doi:10.1111/gcbb.12697.
- Crossley, M., et al., 2020: No net insect abundance and diversity declines across US long term ecological research sites. *Nat. Ecol. Evol.*, **4**, 1368–1376, doi:10.1038/s41559-020-1269-4.

- Cunze, S., L.K. Koch, J. Kochmann and S. Klimpel, 2016: *Aedes albopictus* and *Aedes japonicus* - two invasive mosquito species with different temperature niches in Europe. *Parasites Vectors*, **9**, doi:10.1186/s13071-016-1853-2.
- D'Amen, M. and E. Azzurro, 2020: Lessepsian fish invasion in Mediterranean marine protected areas: a risk assessment under climate change scenarios. *ICES J. Mar. Sci.*, **77**(1), 388–397, doi:10.1093/icesjms/fsz207.
- D'Orazio, P. and L. Popoyan, 2019: Fostering green investments and tackling climate-related financial risks: which role for macroprudential policies? *Ecol. Econ.*, **160**, 25–37, doi:10.1016/j.ecolecon.2019.01.029.
- Dachary-Bernard, J., H. Rey-Valette and B. Rulleau, 2019: Preferences among coastal and inland residents relating to managed retreat: influence of risk perception in acceptability of relocation strategies. *J. Environ. Manag.*, **232**, 772–780, doi:10.1016/j.jenvman.2018.11.104.
- Dadson, S.J., et al., 2017: A restatement of the natural science evidence concerning catchment-based 'natural' flood management in the UK. *Proc. R. Soc. A. Math. Phys. Eng. Sci.*, **473**(2199), 20160706, doi:10.1098/rspa.2016.0706.
- Dale, M., 2021: Managing the effects of extreme sub-daily rainfall and flash floods; a practitioner's perspective. *Philos. Trans. R. Soc. A. Math. Phys. Eng. Sci.*, **379**(2195), 20190550, doi:10.1098/rsta.2019.0550.
- Dale, M., et al., 2018: Understanding how changing rainfall may impact on urban drainage systems; lessons from projects in the UK and USA. *Water Pract. Technol.*, **13**(3), 654–661, doi:10.2166/wpt.2018.069.
- Damm, A., et al., 2017: Impacts of +2°C global warming on electricity demand in Europe. *Clim. Serv.*, **7**, 12–30, doi:10.1016/j.cliser.2016.07.001.
- Daniel, M., et al., 2018: Increased relative risk of tick-borne encephalitis in warmer weather. *Front. Cell. Infect. Microbiol.*, **8**, doi:10.3389/fcimb.2018.00090.
- Daniel, M., et al., 2003: Shift of the tick *Ixodes ricinus* and tick-borne encephalitis to higher altitudes in Central Europe. *Eur. J. Clin. Microbiol. Infect. Dis.*, **22**(5), 327–328, doi:10.1007/s10096-003-0918-2.
- Dapporto, L., et al., 2017: Rise and fall of island butterfly diversity: understanding genetic differentiation and extinction in a highly diverse archipelago. *Divers. Distrib.*, **23**(10), 1169–1181, doi:10.1111/ddi.12610.
- Darmaraki, S., S. Somot, F. Sevault and P. Nabat, 2019: Past variability of mediterranean sea marine heatwaves. *Geophys. Res. Lett.*, **0**(0), doi:10.1029/2019GL082933.
- Davis, M., K. Abhold, L. Mederake and D. Knoblauch, 2018: *Nature-Based Solutions in European and National Policy Frameworks*. https://ec.europa.eu/futurium/en/system/files/ged/naturvation_nature-based_solutions_in_european_and_national_policy_frameworks.pdf. (50 pp), Accessed 2021.
- Davy, R., N. Gnatiuk, L. Pettersson and L. Bobylev, 2018: Climate change impacts on wind energy potential in the European domain with a focus on the Black Sea. *Renew. Sustain. Energy Rev.*, **81**, 1652–1659, doi:10.1016/j.rser.2017.05.253.
- De'Donato, F., et al., 2018: Temporal variation in the effect of heat and the role of the Italian heat prevention plan. *Public Health*, **161**, 154–162, doi:10.1016/j.puhe.2018.03.030.
- de Bruin, K., et al., 2020: Physical climate risks and the financial sector—synthesis of investors' climate information needs. In: *Handbook of Climate Services* [Leal Filho, W. and D. Jacob(eds.)]. Springer International Publishing, Cham, pp. 135–156. ISBN 9783030368746/9783030368753.
- De Cian, E., et al., 2016: Alleviating inequality in climate policy costs: an integrated perspective on mitigation, damage and adaptation. *Environ. Res. Lett.*, **11**(7), 74015–74015, doi:10.1088/1748-9326/11/7/074015.
- de Coninck, H. et al., 2018: Strengthening and implementing the global response. In: *Global Warming of 1.5°C. An IPCC Special Report on the Impacts of Global Warming of 1.5°C Above Pre-industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*. [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P. R. Shukla, A. Pirani, W. MoufoumaOkia, C. Péan, R. Pidcock, S. Connors, J. B. R. Matthews, Y. Chen, X. Zhou, M. I. Gomis, E. Lonnoy, T. Maycock, M. Tignor and T. Waterfield (eds.)], Cambridge University Press, Cambridge, pp. 313–443.
- De Frenne, P., et al., 2013: Microclimate moderates plant responses to macroclimate warming. *Proc. Natl. Acad. Sci.*, **110**(46), 18561, doi:10.1073/pnas.1311190110.
- de la Hoz, C.F., E. Ramos, A. Puente and J.A. Juanes, 2019: Climate change induced range shifts in seaweeds distributions in Europe. *Mar. Environ. Res.*, **148**, 1–11, doi:10.1016/j.marenvres.2019.04.012.
- De los Santos, C.B., J.A. Godbold and M. Solan, 2017: Short-term growth and biomechanical responses of the temperate seagrass *Cymodocea nodosa* to CO2 enrichment. *Mar. Ecol. Prog. Ser.*, **572**, 91–102, doi:10.3354/meps12153.
- de Munck, C., et al., 2018: Evaluating the impacts of greening scenarios on thermal comfort and energy and water consumptions for adapting Paris city to climate change. *Urban Clim.*, **23**, 260–286, doi:10.1016/j.uclim.2017.01.003.
- de Rigo, D., et al., 2017: *Forest Fire Danger Extremes in Europe Under Climate Change: Variability and Uncertainty*. Publications Office of the European Union, Luxembourg.
- De Roo, A., et al., 2020: *Assessing the Effects of Water Saving Measures on Europe's Water Resources; BLUE2 Project – Freshwater Quantity*. JRC Technical Report. ISBN 978-9276215363.
- De Rosa, M., V. Bianco, F. Scarpa and L.A. Tagliafico, 2015: Historical trends and current state of heating and cooling degree days in Italy. *Energy Convers. Manag.*, **90**, 323–335, doi:10.1016/j.enconman.2014.11.022.
- de Schipper, M.A., et al., 2021: Beach nourishment has complex implications for the future of sandy shores. *Nat. Rev. Earth Environ.*, **2**(1), 70–84, doi:10.1038/s43017-020-00109-9.
- Del Bello, L., 2018: Venice anti-flood gates could wreck lagoon ecosystem. *Nature*, **564**(7734), 16–16, doi:10.1038/d41586-018-07372-3.
- Deléglise, C., et al., 2019: A method for diagnosing summer mountain pastures' vulnerability to climate change, developed in the French alps. *Mt. Res. Dev.*, **39**(2), D27–D41, doi:10.1659/MRD-JOURNAL-D-18-00077.1.
- Delgado, M.D.M., et al., 2020: Differences in spatial versus temporal reaction norms for spring and autumn phenological events. *Proc. Natl. Acad. Sci. U.S.A.*, **117**(49), 31249–31258, doi:10.1073/pnas.2002713117.
- Dellink, R., E. Lanzi and J. Chateau, 2019: The sectoral and regional economic consequences of climate change to 2060. *Environ. Resour. Econ.*, **72**(2), 309–363, doi:10.1007/s10640-017-0197-5.
- Demiroglu, O.C., H. Dannevig and C. Aall, 2018: Climate change acknowledgement and responses of summer (glacier) ski visitors in Norway. *Scand. J. Hosp. Tour.*, **18**(4), 419–438, doi:10.1080/15022250.2018.1522721.
- Denechaud, C., et al., 2020: A century of fish growth in relation to climate change, population dynamics and exploitation. *Glob. Change Biol.*, **26**(10), 5661–5678, doi:10.1111/gcb.15298. PMID - 32741054.
- Dennis, E.B., et al., 2019: Trends and indicators for quantifying moth abundance and occupancy in Scotland. *J. Insect Conserv.*, **23**(2), 369–380, doi:10.1007/s10841-019-00135-z.
- Derkzen, M.L., A.J.A. van Teeffelen and P.H. Verburg, 2017: Green infrastructure for urban climate adaptation: How do residents' views on climate impacts and green infrastructure shape adaptation preferences? *Landsc. Urban Plan.*, **157**, 106–130, doi:10.1016/j.landurbplan.2016.05.027.
- Deryng, D., et al., 2014: Global crop yield response to extreme heat stress under multiple climate change futures. *Environ. Res. Lett.*, **9**(3), doi:10.1088/1748-9326/9/3/034011.
- Desmit, X., et al., 2020: Changes in chlorophyll concentration and phenology in the North Sea in relation to de-eutrophication and sea surface warming. *Limnol. Oceanogr.*, **65**(4), 828–847, doi:10.1002/lno.11351.
- Després, J. and M. Adamovic, 2020: *Seasonal Impacts of Climate Change on Electricity Production*. JRC PESETA IV Project – Task 4 J. R. C. (ed.). Joint Research Centre (JRC), Publications Office of the European Union, Luxembourg. 41 pp.

- Devictor, V., R. Julliard, D. Couvet and F. Jiguet, 2008: Birds are tracking climate warming, but not fast enough. *Proc. R. Soc. B.*, **275**(1652), 2743–2748, doi:10.1098/rspb.2008.0878.
- Devictor, V., et al., 2012: Differences in the climatic debts of birds and butterflies at a continental scale. *Nat. Clim. Chang.*, **2**, 121, doi:10.1038/nclimate1347.
- Devis, A., N.P.M. Van Lipzig and M. Demuzere, 2018: Should future wind speed changes be taken into account in wind farm development? *Environ. Res. Lett.*, **13**(6), 64012, doi:10.1088/1748-9326/aabff7.
- Díaz, J., M. Sáez, R. Carmona, I.J. Mirón, M. A. Barceló, M.Y. Luna and C. Linares, 2019: Mortality attributable to high temperatures over the 2021–2050 and 2051–2100 time horizons in Spain: Adaptation and economic estimate. *Environ. Res.*, **172**, 475–485, doi:10.1016/j.envres.2019.02.041
- Di Baldassarre, G., et al., 2018: Water shortages worsened by reservoir effects. *Nat. Sustain.*, **1**(11), 617–622, doi:10.1038/s41893-018-0159-0.
- Di Giuseppe, F., et al., 2020: Fire weather index: the skill provided by the European centre for medium-range weather forecasts ensemble prediction system. *Nat. Hazards Earth Syst. Sci.*, **20**(8), 2365–2378, doi:10.5194/nhess-20-2365-2020.
- Di Lena, B., O. Silvestroni, V. Lanari and A. Palliotti, 2019: Climate change effects on cv. Montepulciano in some wine-growing areas of the Abruzzi region (Italy). *Theor. Appl. Climatol.*, **136**(3), 1145–1155, doi:10.1007/s00704-018-2545-y.
- Díaz, J., et al., 2019: Mortality attributable to high temperatures over the 2021–2050 and 2051–2100 time horizons in Spain: adaptation and economic estimate. *Environ. Res.*, **172**, 475–485, doi:10.1016/j.envres.2019.02.041.
- Diffenbaugh, N.S. and M. Burke, 2019: Global warming has increased global economic inequality. *Proc. Natl. Acad. Sci.*, **116**(20), 9808–9813, doi:10.1073/pnas.1816020116.
- Ding, Q., X. Chen, R. Hilborn and Y. Chen, 2017: Vulnerability to impacts of climate change on marine fisheries and food security. *Mar. Policy*, **83**, 55–61, doi:10.1016/j.marpol.2017.05.011.
- Dino, I.G. and C. Meral Akgül, 2019: Impact of climate change on the existing residential building stock in Turkey: an analysis on energy use, greenhouse gas emissions and occupant comfort. *Renew. Energy*, **141**, 828–846, doi:10.1016/j.renene.2019.03.150.
- Diogo, V., et al., 2017: Assessing local and regional economic impacts of climatic extremes and feasibility of adaptation measures in Dutch arable farming systems. *Agric. Syst.*, **157**(C), 216–229.
- Dodoo, A. and L. Gustavsson, 2016: Energy use and overheating risk of Swedish multi-storey residential buildings under different climate scenarios. *Energy*, **97**, 534–548, doi:10.1016/j.energy.2015.12.086.
- Doll, C., et al., 2014: Adapting rail and road networks to weather extremes: case studies for southern Germany and Austria. *Nat. Hazards*, **72**(1), 63–85, doi:10.1007/s11069-013-0969-3.
- Dominguez-Amarillo, S., J. Fernández-Agüera, J.J. Sendra and S. Roaf, 2019: The performance of Mediterranean low-income housing in scenarios involving climate change. *Energy Build.*, **202**, 109374, doi:10.1016/j.enbuild.2019.109374.
- Donatelli, M., et al., 2015: Climate change impact and potential adaptation strategies under alternate realizations of climate scenarios for three major crops in Europe. *Environ. Res. Lett.*, **10**(7), 75005, doi:10.1088/1748-9326/10/7/075005.
- Donnelly, A., L. Liu, X. Zhang and A. Wingler, 2018: Autumn leaf phenology: discrepancies between in situ observations and satellite data at urban and rural sites. *Int. J. Remote. Sens.*, **39**(22), 8129–8150, doi:10.1080/01431161.2018.1482021.
- Donner, J., J.M. Müller and J. Köppel, 2015: Urban heat: towards adapted German cities? *J. Environ. Assess. Policy Manag.*, **17**(02), 1550020, doi:10.1142/S1464333215500209.
- Dono, G., et al., 2013: Adapting to uncertainty associated with short-term climate variability changes in irrigated mediterranean farming systems. *Agric. Syst.*, **117**, 1–12, doi:10.1016/j.agry.2013.01.005.
- Dórea, F.C., et al., 2016: Vector-borne disease surveillance in livestock populations: a critical review of literature recommendations and implemented surveillance (BTV-8) in five European countries. *Prev. Vet. Med.*, **125**, 1–9, doi:10.1016/j.prevetmed.2016.01.005.
- Dornelas, M., et al., 2014: Assemblage time series reveal biodiversity change but not systematic loss. *Science*, **344**(6181), 296, doi:10.1126/science.1248484.
- Dottori, F., et al., 2020: *Adapting to rising river flood risk in the EU under climate change*. Publications Office of the European Union, Luxembourg, ISBN 978-9276129462.
- Dottori, F., et al., 2018: Increased human and economic losses from river flooding with anthropogenic warming. *Nat. Clim. Chang.*, **8**(9), 781–786, doi:10.1038/s41558-018-0257-z.
- Drobyshev, I., et al., 2015: A 700-year record of large fire years in northern Scandinavia shows large variability and increased frequency during the 1800 s. *J. Quat. Sci.*, **30**(3), 211–221, doi:10.1002/jqs.2765.
- Drobyshev, I., et al., 2016: Atlantic SSTs control regime shifts in forest fire activity of Northern Scandinavia. *Sci. Rep.*, **6**(1), 22532, doi:10.1038/srep22532.
- Dullinger, I., et al., 2020: A socio-ecological model for predicting impacts of land-use and climate change on regional plant diversity in the Austrian Alps. *Glob. Change Biol.*, **26**(4), 2336–2352, doi:10.1111/gcb.14977.
- Dunford, R.W., A.C. Smith, P.A. Harrison and D. Hanganu, 2015: Ecosystem service provision in a changing Europe: adapting to the impacts of combined climate and socio-economic change. *Landsc. Ecol.*, **30**(3), 443–461, doi:10.1007/s10980-014-0148-2.
- Dupuy, J., et al., 2020: Climate change impact on future wildfire danger and activity in southern Europe: a review. *Ann. For. Sci.*, **77**(2), 35, doi:10.1007/s13595-020-00933-5.
- Durant, J.M. and D. Hjermann, 2017: Age-structure, harvesting and climate effects on population growth of Arcto-boreal fish stocks. *Mar. Ecol. Prog. Ser.*, **577**, 177–188, doi:10.3354/meps12210.
- Durant, J.M., et al., 2019: Contrasting effects of rising temperatures on trophic interactions in marine ecosystems. *Sci. Rep.*, **9**(1), 1–9, doi:10.1038/s41598-019-51607-w.
- Dury, M., et al., 2011: Responses of European forest ecosystems to 21st century climate: assessing changes in interannual variability and fire intensity. *Forest - Biogeosciences For.*, **4**(2), 82–99, doi:10.3832/for0572-004.
- Duvillard, P.-A., L. Ravel, M. Marcer and P. Schoeneich, 2019: Recent evolution of damage to infrastructure on permafrost in the French Alps. *Reg. Environ. Chang.*, **19**(5), 1281–1293, doi:10.1007/s10113-019-01465-z.
- Dyderski, M.K., S. Paž, L.E. Frelich and A.M. Jagodziński, 2018: How much does climate change threaten European forest tree species distributions? *Glob. Change Biol.*, **24**(3), 1150–1163, doi:10.1111/gcb.13925.
- EASAC, 2017: *Multi-functionality and Sustainability in the European Union's Forests*. Sciences, EASAC, Halle, www.easac.eu. Accessed 2021.
- EASAC, 2019: *Forest Bioenergy, Carbon Capture and Storage, and Carbon Dioxide Removal: An Update*. EASAC, Halle, 12 pp.
- Ebi, K.L., et al., 2021: Burning embers: synthesis of the health risks of climate change. *Environ. Res. Lett.*, **16**(4), 44042, doi:10.1088/1748-9326/abeadd.
- ECB, 2021a: *Climate-related Risk and Financial Stability*. ECB/ESRB Project Team on climate risk monitoring, Frankfurt, Germany, https://data.europa.eu/doi/10.2866/913118. Accessed 2021.
- ECB, 2021b: *Shining a Light on Climate Risks: the ECB's Economy-wide Climate Stress Test*. European Central Bank, Frankfurt am Main.
- Edelgeriev, R.S.-H. (ed.), 2019: *National Report "Global Climate and Soil Cover of Russia: Desertification and Land Degradation, Institutional, Infrastructure, Technological Adaptation Measures (Agriculture and Forestry)"*. Vol. 2. LLC "Publishing House MBA", Moscow, ISBN 978-5604322567. 476 pp.
- EEA, 2014: *Adaptation of Transport to Climate Change in Europe - Challenges and Options Across Transport Modes and Stakeholders* [EEA (ed.)]. Publications Office of the European Union, Luxembourg, 58 pp.
- EEA, 2016a: *Climate-ADAPT*. https://climate-adapt.eea.europa.eu/metadata/case-studies/urban-storm-water-management-in-augustenburg-

- malmo/#source. Publications Office of the European Union, Luxembourg, Accessed 2021.
- EEA, 2016b: *Urban Adaptation to Climate Change in Europe 2016 - Transforming Cities in a Changing Climate*. Publications Office of the European Union, Luxembourg. 135 pp.
- EEA, 2017a: *Climate Change Adaptation and Disaster Risk Reduction in Europe Enhancing Coherence of the Knowledge Base, Policies and Practices*. Publications Office of the European Union, Luxembourg.
- EEA, 2017b: *Climate Change, Impacts and Vulnerability in Europe 2016 - An Indicator-based Report* [Agency, E. E. (ed.)]. EEA report, Vol. 1/2017. Publications Office of the European Union, Luxembourg. 424 pp.
- EEA, 2018: *Climate-Adapt, 2019, Climate-Adapt. Sharing Adaptation Information Across Europe*. EEA Report, Vol. 3/2018. Publications Office of the European Union, Luxembourg.
- EEA, 2019a: *Adaptation Challenges and Opportunities for the European Energy System - Building a Climate-resilient Low-carbon Energy System*. Publications Office of the European Union, Luxembourg. 122 pp.
- EEA, 2019b: *Climate Change Adaptation in the Agriculture Sector in Europe*. EEA Report Vol. 04/2019, Publications Office of the European Union, Luxembourg, 112 pp.
- EEA, 2019c: *Flood Defence Framework for National Grid Substations in United Kingdom — Climate-ADAPT*.
- EEA, 2020a: *Monitoring and Evaluation of National Adaptation Policies Throughout the Policy Cycle*. Publications Office of the European Union, Luxembourg, <https://data.europa.eu/doi/10.2800/83221>. Accessed 2020.
- EEA, 2020b: *Urban Adaptation in Europe - How Cities and Towns Respond to Climate Change*. Publications Office of the European Union, Luxembourg. 192 pp.
- EEA, 2020c: *Urban Adaptation in Europe: How Cities and Towns Respond to Climate Change*. EEA Report. Publications Office of the European Union, Luxembourg. 192 pp.
- Eftestøl, S., K. Flydal, D. Tsegaye and J.E. Colman, 2019: Mining activity disturbs habitat use of reindeer in Finnmark, Northern Norway. *Polar Biol.*, **42**(10), 1849–1858, doi:10.1007/s00300-019-02563-8.
- Eisenack, K., 2016: Institutional adaptation to cooling water scarcity for thermoelectric power generation under global warming. *Ecol. Econ.*, **124**, 153–163, doi:10.1016/j.ecolecon.2016.01.016.
- Elmendorf, S.C., et al., 2012: Plot-scale evidence of tundra vegetation change and links to recent summer warming. *Nat. Clim. Chang.*, **2**(6), 453–457, doi:10.1038/nclimate1465.
- Elmhagen, B., J. Kindberg, P. Hellström and A. Angerbjörn, 2015: A boreal invasion in response to climate change? Range shifts and community effects in the borderland between forest and tundra. *Ambio*, **44**(1), 39–50, doi:10.1007/s13280-014-0606-8.
- Emmanuel, R. and A. Loconsole, 2015: Green infrastructure as an adaptation approach to tackling urban overheating in the Glasgow Clyde Valley Region, UK. *Landsc. Urban Plan.*, **138**, 71–86, doi:10.1016/j.landurbplan.2015.02.012.
- Enríquez-de-Salamanca, Á., 2019: Environmental impacts of climate change adaptation of road pavements and mitigation options. *Int. J. Pavement Eng.*, **20**(6), 691–696, doi:10.1080/10298436.2017.1326236.
- Erauskin-Extramiana, M., et al., 2019: Historical trends and future distribution of anchovy spawning in the Bay of Biscay. *Deep. Res. Part II Top. Stud. Oceanogr.*, **159**, 169–182, doi:10.1016/j.dsr2.2018.07.007.
- Erjavec, E., et al., 2017: Interactions between European agricultural policy and climate change: a Slovenian case study. *Clim. Policy*, **17**(8), 1014–1030, doi:10.1080/14693062.2016.1222259.
- Escobar, L.E., et al., 2015: A global map of suitability for coastal *Vibrio cholerae* under current and future climate conditions. *Acta Trop.*, **149**, 202–211, doi:10.1016/j.actatropica.2015.05.028.
- Esteve, P., C. Varela-Ortega, I. Blanco-Gutiérrez and T.E. Downing, 2015: A hydro-economic model for the assessment of climate change impacts and adaptation in irrigated agriculture. *Ecol. Econ.*, **120**, 49–58, doi:10.1016/j.ecolecon.2015.09.017.
- Estrada-Pena, A., N. Ayllon and J. de la Fuente, 2012: Impact of climate trends on tick-borne pathogen transmission. *Front. Physiol.*, **3**(64), doi:10.3389/fphys.2012.00064.
- Estrada-Pena, A. and N. Fernandez-Ruiz, 2020: A retrospective assessment of temperature trends in northern Europe reveals a deep impact on the life cycle of *Ixodes ricinus* (Acari: Ixodidae). *Pathogens*, **9**(5), doi:10.3390/pathogens9050345.
- Ettinger, A.K., et al., 2020: Winter temperatures predominate in spring phenological responses to warming. *Nat. Clim. Chang.*, **10**(12), 1137–1142.
- European Commission, 2015: Our life insurance, our natural capital: an EU biodiversity strategy to 2020. *Landsc. Ecol. Manag.*, **20**(1), 37–40, doi:10.5738/jale.20.37.
- European Commission, 2020: *Nature-Based Solutions for Flood Mitigation and Coastal Resilience. Analysis of EU-funded Projects*. Publications Office of the European Union, Luxembourg. 56 pp.
- Fader, M., et al., 2016: Mediterranean irrigation under climate change: more efficient irrigation needed to compensate for increases in irrigation water requirements. *Hydrol. Earth Syst. Sci.*, **20**(2), 953–973, doi:10.5194/hess-20-953-2016.
- Falk, M., 2015: The demand for winter sports: empirical evidence for the largest French ski-lift operator. *Tour. Econ.*, **21**(3), 561–580, doi:10.5367/te.2013.0366.
- Falk, M. and X. Lin, 2018: Sensitivity of winter tourism to temperature increases over the last decades. *Econ. Model.*, **71**, 174–183, doi:10.1016/j.econmod.2017.12.011.
- Falk, M. and M. Scaglione, 2018: Effects of ski lift ticket discounts on local tourism demand. *Tour. Rev.*, **73**(4), 480–491, doi:10.1108/TR-08-2017-0133.
- Falk, M. and L. Vanat, 2016: Gains from investments in snowmaking facilities. *Ecol. Econ.*, **130**, 339–349, doi:10.1016/j.ecolecon.2016.08.003.
- Fallmann, J., S. Emeis and P. Suppan, 2013: Mitigation of urban heat stress - a modelling case study for the area of Stuttgart. *Erde*, **144**(3–4), 202–216.
- Fantappiè, M., G. L'Abate and E.A.C. Costantini, 2011: The influence of climate change on the soil organic carbon content in Italy from 1961 to 2008. *Geomorphology*, **135**(3–4), 343–352, doi:10.1016/j.geomorph.2011.02.006.
- Faria, N. and M.B. Morales, 2020: Farmland management regulates ecosystem services in Mediterranean drylands: Assessing the sustainability of agri-environmental payments for bird conservation. *J. Nat. Conserv.*, **58**, 125913–125913, doi:10.1016/j.jnc.2020.125913.
- Faust, A.-K., C. Gonseth and M. Vielle, 2015: The economic impact of climate-driven changes in water availability in Switzerland. *Water Policy*, **17**(5), 848–864, doi:10.2166/wp.2015.064.
- Feodoroff, P., 2021: Indigenous Female Bodies as Indicators of Change. In: *2021 Compendium of Indigenous Knowledge and Local Knowledge: Towards Inclusion of Indigenous Knowledge and Local Knowledge in Global Reports on Climate Change*. [Mustonen, T., S. Harper, M. Rivera-Ferre, J. C. Postigo, A. Ayansina, T. Benjaminsen, R. Morgan and A. Okem (eds.)]. Snowchange Cooperative, Kontiolahti, Finland.
- Feridun, M. and H. Güngör, 2020: Climate-related prudential risks in the banking sector: a review of the emerging regulatory and supervisory practices. *Sustainability*, **12**(13), 5325, doi:10.3390/su12135325.
- Fernandes, J.A., et al., 2017: Estimating the ecological, economic and social impacts of ocean acidification and warming on UK fisheries. *Fish Fish.*, **18**(3), 389–411, doi:10.1111/faf.12183.
- Fernández-Giménez, M.E. and J. Ritten, 2020: An economic analysis of transhumance in the Central Spanish Pyrenees. *Pastoralism*, **10**(1), doi:10.1186/s13570-020-00163-4.
- Fernández-Martínez, M., et al., 2019: Global trends in carbon sinks and their relationships with CO₂ and temperature. *Nat. Clim. Chang.*, **9**(1), 73–79, doi:10.1038/s41558-018-0367-7.
- Fernández-Montblanc, T., M.I. Voudoukas, L. Mentaschi and P. Ciavola, 2020: A Pan-European high resolution storm surge hindcast. *Environ. Int.*, **135**, 105367, doi:10.1016/j.envint.2019.105367.

- Ferranti, E., et al., 2018: The hottest July day on the railway network: insights and thoughts for the future. *Meteorol. Appl.*, **25**(2), 195–208, doi:10.1002/met.1681.
- Ferranti, E., et al., 2016: Heat-related failures on southeast England's railway network: insights and implications for heat risk management. *Weather Clim. Soc.*, **8**(2), 177–191, doi:10.1175/wcas-d-15-0068.1.
- Ferrara, M. and E. Fabrizio, 2017: Cost optimal nZEBs in future climate scenarios. *Energy Procedia*, **122**, 877–882, doi:10.1016/j.egypro.2017.07.377.
- Ferretto, A., et al., 2019: Potential carbon loss from Scottish peatlands under climate change. *Reg. Environ. Chang.*, **19**(7), 2101–2111, doi:10.1007/s10113-019-01550-3.
- Feyen, L., et al., 2020: *Climate change impacts and adaptation in Europe*, JRC PESETA IV Final Report. Publications Office of the European Union, Luxembourg, ISBN 978-9276181231.
- Filatov, N., L. Nazarova and P. Druzhinin, 2019: Influence of climatic and anthropogenic factors on the white sea – catchment system. *Proc. Karelian Res. Centre Russ. Acad. Sci.*, **9**, 30–30, doi:10.17076/lim1117.
- Filipchuk, A., B. Moiseev, N. Malysheva and V. Strakhov, 2018: Russian forests: a new approach to the assessment of carbon stocks and sequestration capacity. *Environ. Dev.*, **26**, 68–75, doi:10.1016/j.envdev.2018.03.002.
- Filipe, A., et al., 2013: Forecasting fish distribution along stream networks: brown trout (*Salmo trutta*) in Europe. *Divers. Distrib.*, **19**(8), 1059–1071, doi:10.1111/ddi.12086.
- Fischer, K., T. Stenius and S. Holmgren, 2020a: Swedish forests in the bioeconomy: stories from the national forest program. *Soc. Nat. Resour.*, **33**(7), 896–913, doi:10.1080/08941920.2020.1725202.
- Fischer, L., et al., 2020b: Rising temperature and its impact on receptivity to malaria transmission in Europe: a systematic review. *Travel Med. Infect. Dis.*, **36**, doi:10.1016/j.tmaid.2020.101815.
- Flekens, L., et al., 2013: Regional consequences of the way land users respond to future water availability in Murcia, Spain. *Reg. Environ. Chang.*, **13**(3), 615–632, doi:10.1007/s10113-012-0283-8.
- Forbes, B.C., et al., 2016: Sea ice, rain-on-snow and tundra reindeer nomadism in Arctic Russia. *Biol. Lett.*, **12**(11), 20160466, doi:10.1098/rsbl.2016.0466.
- Ford, H., et al., 2021: Hedgerow effects on CO2 emissions are regulated by soil type and season: implications for carbon flux dynamics in livestock-grazed pasture. *Geoderma*, **382**, 114697–114697, doi:10.1016/j.geoderma.2020.114697.
- Fornara, D., R. Olave and A. Higgins, 2019: Evidence of low response of soil carbon stocks to grassland intensification. *Agric. Ecosyst. Environ.*, **287**, doi:10.1016/j.agee.2019.106705.
- Fortibuoni, T., et al., 2015: Climate impact on Italian fisheries (Mediterranean Sea). *Reg. Environ. Chang.*, **15**(5), 931–937, doi:10.1007/s10113-015-0781-6.
- Forzieri, G., et al., 2018: Escalating impacts of climate extremes on critical infrastructures in Europe. *Glob. Environ. Chang.*, **48**(November 2017), 97–107, doi:10.1016/j.gloenvcha.2017.11.007.
- Forzieri, G., A. Cescatti, F.B. e Silva and L. Feyen, 2017: Increasing risk over time of weather-related hazards to the European population: a data-driven prognostic study. *Lancet Planet. Health*, **1**(5), e200–e208, doi:10.1016/S2542-5196(17)30082-7.
- Fossheim, M., et al., 2015: Recent warming leads to a rapid borealization of fish communities in the Arctic. *Nat. Clim. Chang.*, **5**(7), 673–677, doi:10.1038/nclimate2647.
- Fourcade, Y., S. Åström and E. Öckinger, 2019: Climate and land-cover change alter bumblebee species richness and community composition in subalpine areas. *Biodivers. Conserv.*, **28**(3), 639–653.
- Fox-Kemper, B., et al., 2021: Ocean, Cryosphere and Sea Level Change. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press, Cambridge.
- Fraga, H., J.G. Pinto and J.A. Santos, 2019: Climate change projections for chilling and heat forcing conditions in European vineyards and olive orchards: a multi-model assessment. *Clim. Chang.*, **152**(1), 179–193, doi:10.1007/s10584-018-2337-5.
- Frainer, A., et al., 2020: Opinion: cultural and linguistic diversities are underappreciated pillars of biodiversity. *Proc. Natl. Acad. Sci.*, **117**(43), 26539–26543, doi:10.1073/pnas.2019469117.
- Frainer, A., et al., 2017: Climate-driven changes in functional biogeography of Arctic marine fish communities. *Proc. Natl. Acad. Sci. U.S.A.*, **114**(46), 12202–12207, doi:10.1073/pnas.1706080114.
- Franz, M., et al., 2019: Long-term records of hard-bottom communities in the southwestern Baltic Sea reveal the decline of a foundation species. *Estuar. Coast. Shelf Sci.*, **219**, 242–251, doi:10.1016/j.ecss.2019.02.029.
- Franzén, M. and E. Öckinger, 2012: Climate-driven changes in pollinator assemblages during the last 60 years in an Arctic mountain region in Northern Scandinavia. *J. Insect Conserv.*, **16**(2), 227–238, doi:10.1007/s10841-011-9410-y.
- Free, C.M., et al., 2019: Impacts of historical warming on marine fisheries production. *Science*, **363**(6430), 979–983, doi:10.1126/science.aau1758.
- Fréjaville, T., 2017: Seasonal changes in the human alteration of fire regimes beyond the climate forcing. *Environ. Res. Lett.*, **12**, 35006.
- Frölicher, T.L., E.M. Fischer and N. Gruber, 2018: Marine heatwaves under global warming. *Nature*, **560**(7718), 360–364, doi:10.1038/s41586-018-0383-9.
- Frolov, A.V., et al., 2014: *Second Roshydromet Assessment Report on Climate Change and its Consequences in Russian Federation* [Yasukevich, V. V., V. A. Govorkova, I. A. Korneva, T. V. Pavlova and E. N. Popova (eds.)]. Roshydromet, Obninsk, Russia. 1004 pp. http://downloads.igce.ru/publications/OD_2_2014/v2014/htm/1.htm.
- Fu, G., et al., 2018: Integrated approach to assess the resilience of future electricity infrastructure networks to climate hazards. *IEEE Syst. J.*, **12**(4), 3169–3180, doi:10.1109/JSYST.2017.2700791.
- Fu, Y.H., et al., 2015: Declining global warming effects on the phenology of spring leaf unfolding. *Nature*, **526**(7571), 104–107, doi:10.1038/nature15402.
- Furberg, M., B. Evengård and M. Nilsson, 2011: Facing the limit of resilience: perceptions of climate change among reindeer herding Sami in Sweden. *Glob. Health Action*, **4**(1), 8417, doi:10.3402/gha.v4i0.8417.
- Gabalton-Leal, C., et al., 2015: Strategies for adapting maize to climate change and extreme temperatures in Andalusia, Spain. *Clim. Res.*, **65**, 159–173, doi:10.3354/cr01311.
- Gain, A.K., C. Giupponi and Y. Wada, 2016: Measuring global water security towards sustainable development goals. *Environ. Res. Lett.*, **11**(12), 124015, doi:10.1088/1748-9326/11/12/124015.
- Gallego-Sala, A.V., et al., 2010: Bioclimatic envelope model of climate change impacts on blanket peatland distribution in Great Britain. *Clim. Res.*, **45**, 151–162.
- Galli, G., C. Solidoro and T. Lovato, 2017: Marine heat waves hazard 3D maps and the risk for low motility organisms in a warming Mediterranean Sea. *Front. Mar. Sci.*, **4**(136), doi:10.3389/fmars.2017.00136.
- Galofré, J., J.A. Jiménez and H.I. Valdemoro, 2016: Beach Restoration in the Tarragona Coast (Spain); Sand Management During the Last 25 years and Future Plans. *Coast. Eng. Proc.*, **1**(36), risk.20. doi:10.9753/icce.v36.risk.20.
- Gamito, R., C.M. Teixeira, M.J. Costa and H.N. Cabral, 2015: Are regional fisheries' catches changing with climate? *Fish. Res.*, **161**, 207–216, doi:10.1016/j.fishres.2014.07.014.
- Gampe, D., G. Nikulin and R. Ludwig, 2016: Using an ensemble of regional climate models to assess climate change impacts on water scarcity in European river basins. *Sci. Total. Environ.*, **573**, 1503–1518, doi:10.1016/j.scitotenv.2016.08.053.
- Ganguli, P. and B. Merz, 2019: Trends in compound flooding in northwestern Europe during 1901–2014. *Geophys. Res. Lett.*, **46**(19), 10810–10820, doi:10.1029/2019GL084220.

- Gao, G., A.S. Clare, C. Rose and G.S. Caldwell, 2018: Ulva rigida in the future ocean: potential for carbon capture, bioremediation and biomethane production. *Glob. Change Biol. Bioenergy*, **10**(1), 39–51, doi:10.1111/gcbb.12465.
- García-Herrera, R., et al., 2019: The European 2016/2017 drought. *J. Climate*, **0**(0), doi:10.1175/jcli-d-18-0331.1.
- García-León, D., et al., 2021: Current and projected regional economic impacts of heatwaves in Europe. *Nat. Commun.*, **12**, 5807. doi:10.1038/s41467-021-26050-z.
- García-Molinos, J., M.T. Burrows and E.S. Poloczanska, 2017: Ocean currents modify the coupling between climate change and biogeographical shifts. *Sci. Rep.*, **7**(1), 1332, doi:10.1038/s41598-017-01309-y.
- García-Mozo, H., J. Oteros and C. Galán, 2015: Phenological changes in olive (*Ola europaea* L.) reproductive cycle in southern Spain due to climate change. *Ann. Agric. Environ. Med.*, **22**(3), 421–428, doi:10.5604/12321966.1167706.
- García Sánchez, F., W.D. Solecki and C. Ribalaygua Batalla, 2018: Climate change adaptation in Europe and the United States: a comparative approach to urban green spaces in Bilbao and New York City. *Land Use Policy*, **79**, 164–173, doi:10.1016/j.landusepol.2018.08.010.
- García Molinos, J., et al., 2016: Climate velocity and the future global redistribution of marine biodiversity. *Nat. Clim. Chang.*, **6**(1), 83–88, doi:10.1038/nclimate2769.
- Garnier, E., et al., 2018: Historical analysis of storm events: case studies in France, England, Portugal and Italy. *Coast. Eng.*, **134**, 10–23, doi:10.1016/j.coastaleng.2017.06.014.
- Garnier, M. and I. Holman, 2019: Critical review of adaptation measures to reduce the vulnerability of European drinking water resources to the pressures of climate change. *Environ. Manag.*, **64**(2), 138–153, doi:10.1007/s00267-019-01184-5.
- Garonna, I., et al., 2014: Strong contribution of autumn phenology to changes in satellite-derived growing season length estimates across Europe (1982–2011). *Glob. Change Biol.*, **20**(11), 3457–3470, doi:10.1111/gcb.12625.
- Garrabou, J., et al., 2009: Mass mortality in Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave. *Glob. Change Biol.*, **15**(5), 1090–1103, doi:10.1111/j.1365-2486.2008.01823.x.
- Garrabou, J., et al., 2019: Collaborative database to track mass mortality events in the Mediterranean sea. *Front. Mar. Sci.*, **6**, 2775, doi:10.3389/fmars.2019.00707.
- Garrote, L., et al., 2015: Quantitative assessment of climate change vulnerability of irrigation demands in Mediterranean Europe. *Water Resour. Manag.*, **29**(2), 325–338, doi:10.1007/s11269-014-0736-6.
- Gasbarro, F., F. Rizzi and M. Frey, 2016: Adaptation measures of energy and utility companies to cope with water scarcity induced by climate change. *Bus. Strat. Env.*, **25**(1), 54–72, doi:10.1002/bse.1857.
- Gasho, E. G. (ed.) 2019: *The Priorities of the Megalopolis Climate Change Adaptation: People, Nature, Technology. Algorithm, Strategy, and Action Plan. Scientific and Methodical Edition.* The Moscow Government, Moscow, ISBN 978-5990923041.
- Gasparrini, A., et al., 2015: Temporal variation in heat–mortality associations: a multicountry study. *Environ. Health Perspect.*, **123**(11), 1200–1207, doi:10.1289/ehp.1409070.
- Gasparrini, A., et al., 2017: Projections of temperature-related excess mortality under climate change scenarios. *Lancet Planet. Health*, **1**(9), e360–e367, doi:10.1016/S2542-5196(17)30156-0.
- Gaudard, L., M. Gilli and F. Romero, 2013: Climate change impacts on hydropower management. *Water Resour. Manag.*, **27**(15), 5143–5156, doi:10.1007/s11269-013-0458-1.
- Gaudin, F., et al., 2018: Marine sublittoral benthos fails to track temperature in response to climate change in a biogeographical transition zone. *ICES J. Mar. Sci.*, **75**(6), 1894–1907, doi:10.1093/icesjms/fsy095.
- Gauzere, J., et al., 2017: Integrating interactive effects of chilling and photoperiod in phenological process-based models. A case study with two European tree species: *Fagus sylvatica* and *Quercus petraea*. *Agric. For. Meteorol.*, **244–245**, 9–20, doi:10.1016/j.agrformet.2017.05.011.
- Gauzère, P., F. Jiguet and V. Devictor, 2016: Can protected areas mitigate the impacts of climate change on bird's species and communities? *Divers. Distrib.*, **22**(6), 625–637, doi:10.1111/ddi.12426.
- Gazeau, F., et al., 2014: Impact of ocean acidification and warming on the Mediterranean mussel (*Mytilus galloprovincialis*). *Front. Mar. Sci.*, **1**(NOV), 62–62, doi:10.3389/fmars.2014.00062.
- Gazol, A., et al., 2015: Distinct effects of climate warming on populations of silver fir (*Abies alba*) across Europe. *J. Biogeogr.*, **42**(6), 1150–1162, doi:10.1111/jbi.12512.
- Gazol, A., et al., 2018: Forest resilience to drought varies across biomes. *Glob. Change Biol.*, **24**(5), 2143–2158.
- Gedikli, B. and O. Balaban, 2018: An evaluation of local policies and actions that address climate change in Turkish metropolitan cities. *Eur. Plan. Stud.*, **26**(3), 458–479, doi:10.1080/09654313.2017.1397107.
- Geneletti, D. and L. Zardo, 2016: Ecosystem-based adaptation in cities: an analysis of European urban climate adaptation plans. *Land Use Policy*, **50**, 38–47, doi:10.1016/j.landusepol.2015.09.003.
- Ghanem, D., S. Mander and C. Gough, 2016: “I think we need to get a better generator”: Household resilience to disruption to power supply during storm events. *Energy Policy*, **92**, 171–180, doi:10.1016/j.enpol.2016.02.003.
- Gianinetto, M., et al., 2020: Future scenarios of soil erosion in the Alps under climate change and land cover transformations simulated with automatic machine learning. *Climate*, **8**(2), 28.
- Gill, A.L., et al., 2015: Changes in autumn senescence in northern hemisphere deciduous trees: a meta-analysis of autumn phenology studies. *Ann. Bot.*, **116**(6), 875–888, doi:10.1093/aob/mcv055.
- Giuntoli, I., B. Renard, J.P. Vidal and A. Bard, 2013: Low flows in France and their relationship to large-scale climate indices. *J. Hydrol.*, **482**, 105–118, doi:10.1016/j.jhydrol.2012.12.038.
- Glushenkov, O.V., 2017: The extending of ranges of some bird species at the north-eastern border of their distribution due to intra-century climate changes. *Nat. Conserv. Res.*, **2**(3), 23–39, doi:10.24189/ncr.2017.047.
- Gocht, M. and G. Meon, 2016: Modelling and assessment of the combined impacts of climatic and demographic change on a multipurpose reservoir system in the Harz mountains. *Environ. Earth Sci.*, **75**(21), 1395, doi:10.1007/s12665-016-6099-y.
- Goldberg, D.S., I. v. Rijn, M. Kiflawi and J. Belmaker, 2019: Decreases in length at maturation of Mediterranean fishes associated with higher sea temperatures. *ICES J. Mar. Sci.*, **76**(4), 946–959, doi:10.1093/icesjms/fsz011.
- Gómez-Martín, M.B., X.A. Armesto-López, M. Cors-Iglesias and J. Muñoz-Negrete, 2014: Adaptation strategies to climate change in the tourist sector: the case of coastal tourism in Spain. *Tour. Int. Interdiscip. J.*, **62**(3), 15.
- Gómez, J., 2019: The silent extinction: climate change and the potential hybridization-mediated extinction of endemic high-mountain plants. *Biodivers. Conserv.*, **24**, 1843–1857, doi:10.1007/s10531-015-0909-5. SpringerLink.
- Green, J.K., et al., 2019: Large influence of soil moisture on long-term terrestrial carbon uptake. *Nature*, **565**(7740), 476–479, doi:10.1038/s41586-018-0848-x.
- Greve, P., et al., 2018: Global assessment of water challenges under uncertainty in water scarcity projections. *Nat. Sustain.*, **1**(9), 486–494, doi:10.1038/s41893-018-0134-9.
- Grillakis, M., A. Koutroulis and I. Tsanis, 2016: The 2°C global warming effect on summer European tourism through different indices. *Int. J. Biometeorol.*, **60**(8), 1205–1215, doi:10.1007/s00484-015-1115-6.
- Grillakis, M.G., 2019: Increase in severe and extreme soil moisture droughts for Europe under climate change. *Sci. Total Environ.*, **660**, 1245–1255, doi:10.1016/j.scitotenv.2019.01.001.
- Grossel, A., et al., 2016: The effect of tile-drainage on nitrous oxide emissions from soils and drainage streams in a cropped landscape in Central France. *Agric. Ecosyst. Environ.*, **230**, 251–260.

- Grüneis, H., et al., 2018: Why do we not pick the low-hanging fruit? Governing adaptation to climate change and resilience in Tyrolean mountain agriculture. *Land Use Policy*, **79**, 386–396, doi:10.1016/j.landusepol.2018.08.025.
- Guerra, C.A., M.J. Metzger, J. Maes and T. Pinto-Correia, 2016: Policy impacts on regulating ecosystem services: looking at the implications of 60 years of landscape change on soil erosion prevention in a Mediterranean silvo-pastoral system. *Landsc. Ecol.*, **31**(2), 271–290, doi:10.1007/s10980-015-0241-1.
- Gulev, S.K., P.W. Thorne, J. Ahn, F.J. Dentener, C.M. Domingues, S. Gerland, D. Gong, D.S. Kaufman, H.C. Nnamchi, J. Quaas, J.A. Rivera, S. Sathyendranath, S.L. Smith, B. Trewin, K. von Schuckmann, and R.S. Vose, 2021: Changing State of the Climate System. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. [Masson- Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge. In Press.
- Guo, Y., et al., 2018: Quantifying excess deaths related to heatwaves under climate change scenarios: a multicountry time series modelling study. *PLoS Med.*, **15**(7), e1002629.
- Gusarov, A.V., 2020: The response of water flow, suspended sediment yield and erosion intensity to contemporary long-term changes in climate and land use/cover in river basins of the Middle Volga Region, European Russia. *Sci. Total Environ.*, **719**, 134770–134770, doi:10.1016/j.scitotenv.2019.134770.
- Gusarov, A.V., V.N. Golosov and A.G. Sharifullin, 2018: Contribution of climate and land cover changes to reduction in soil erosion rates within small cultivated catchments in the eastern part of the Russian Plain during the last 60 years. *Environ. Res.*, **167**, 21–33, doi:10.1016/j.envres.2018.06.046.
- Güsewell, S., R. Furrer, R. Gehrig and B. Pietragalla, 2017: Changes in temperature sensitivity of spring phenology with recent climate warming in Switzerland are related to shifts of the pre-season. *Glob. Change Biol.*, **23**(12), 5189–5202, doi:10.1111/gcb.13781.
- Gutiérrez, C., et al., 2020: Future evolution of surface solar radiation and photovoltaic potential in Europe: investigating the role of aerosols. *Environ. Res. Lett.*, **15**, 34035, doi:10.1088/1748-9326/ab6666.
- Haanpää, S., S. Juhola and M. Landauer, 2015: Adapting to climate change: perceptions of vulnerability of down-hill ski area operators in Southern and Middle Finland. *Curr. Issues Tour.*, **18**(10), 966–978, doi:10.1080/13683500.2014.892917.
- Haasnoot, M., J. Kwakkel, W. Walker and J. ter Maat, 2013: Dynamic adaptive policy pathways: a method for crafting robust decisions for a deeply uncertain world. *Glob. Environ. Change Policy Dimens.*, **23**(2), 485–498, doi:10.1016/j.gloenvcha.2012.12.006.
- Haasnoot, M., A. Warren and J.H. Kwakkel, 2019: Dynamic Adaptive Policy Pathways (DAPP). In: *Decision Making under Deep Uncertainty: From Theory to Practice* [Marchau, V.A.W.J., W.E. Walker, P.J.T.M. Bloemen and S.W. Popper(eds.)]. Springer International Publishing, Cham, pp. 71–92. ISBN 978-3030052522.
- Haasnoot, M., et al., 2021: Long-term sea-level rise necessitates a commitment to adaptation: a first order assessment. *Clim. Risk Manag.*, **34**, 100355, doi:10.1016/j.crm.2021.100355.
- Habel, J., M. Samways and T. Schmitt, 2019: Mitigating the precipitous decline of terrestrial European insects: requirements for a new strategy. *Biodivers. Conserv.*, **28**(6), 1343–1360, doi:10.1007/s10531-019-01741-8.
- Haer, T., W.J.W. Botzen and J.C.J.H. Aerts, 2019: Advancing disaster policies by integrating dynamic adaptive behaviour in risk assessments using an agent-based modelling approach. *Environ. Res. Lett.*, **14**(4), 44022–44022, doi:10.1088/1748-9326/ab0770.
- Haigh, I., R. Nicholls and N. Wells, 2011: Rising sea levels in the English Channel 1900 to 2100. *Proc. Inst. Civ. Eng. Eng.*, **164**(2), 81–92, doi:10.1680/maen.2011.164.2.81.
- Halkos, G., A. Skouloudis, C. Malesios and K. Evangelinos, 2018: Bouncing back from extreme weather events: some preliminary findings on resilience barriers facing small and medium-sized enterprises. *Bus. Strateg. Environ.*, **27**(4), 547–559, doi:10.1002/bse.2019.
- Halupka, L. and K. Halupka, 2017: The effect of climate change on the duration of avian breeding seasons: a meta-analysis. *Proc. R. Soc. B.*, **284**(1867), 20171710, doi:10.1098/rspb.2017.1710.
- Hamdy, M., S. Carlucci, P.-J. Hoes and J.L.M. Hensen, 2017: The impact of climate change on the overheating risk in dwellings—a Dutch case study. *Build. Environ.*, **122**, 307–323, doi:10.1016/j.buildenv.2017.06.031.
- Hamidov, A., et al., 2018: Impacts of climate change adaptation options on soil functions: a review of European case-studies. *Land Degrad. Dev.*, **29**(8), 2378–2389, doi:10.1002/ldr.3006.
- Handisyde, N., T.C. Telfer and L.G. Ross, 2017: Vulnerability of aquaculture-related livelihoods to changing climate at the global scale. *Fish Fish.*, **18**(3), 466–488, doi:10.1111/faf.12186.
- Hanger, S., et al., 2018: Insurance, public assistance, and household flood risk reduction: a comparative study of Austria, England, and Romania. *Risk Anal.*, **38**(4), 680–693, doi:10.1111/risa.12881.
- Hannah, L., et al., 2013: Climate change, wine, and conservation. *Proc. Natl. Acad. Sci.*, **110**(17), 6907–6912, doi:10.1073/pnas.1210127110.
- Hansen, B.B., et al., 2014: Warmer and wetter winters: characteristics and implications of an extreme weather event in the High Arctic. *Environ. Res. Lett.*, **9**(11), 114021, doi:10.1088/1748-9326/9/11/114021.
- Hanski, J., T. Rosqvist and D. Crawford-Brown, 2018: Assessing climate change adaptation strategies—the case of drought and heat wave in the French nuclear sector. *Reg. Environ. Chang.*, **18**(6), 1801–1813, doi:10.1007/s10113-018-1312-z.
- Harman, B.P., S. Heyenga, B.M. Taylor and C.S. Fletcher, 2015: Global lessons for adapting coastal communities to protect against storm surge inundation. *J. Coast. Res.*, **31**(4), 790–801, doi:10.2112/JCOASTRES-D-13-00095.1.
- Harmanny, K.S. and Ž. Malek, 2019: Adaptations in irrigated agriculture in the Mediterranean region: an overview and spatial analysis of implemented strategies. *Reg. Environ. Chang.*, **19**(5), 1401–1416, doi:10.1007/s10113-019-01494-8.
- Harris, J., N. Rodenhouse and R. Holmes, 2019: Decline in beetle abundance and diversity in an intact temperate forest linked to climate warming. *Biol. Conserv.*, **240**, doi:10.1016/j.biocon.2019.108219.
- Harrison, P.A., et al., 2019: Differences between low-end and high-end climate change impacts in Europe across multiple sectors. *Reg. Environ. Chang.*, **16**, 695–709, doi:10.1007/s10113-018-1352-4.
- Harrus, S. and G. Baneth, 2005: Drivers for the emergence and re-emergence of vector-borne protozoal and bacterial diseases. *Int. J. Parasitol.*, **35**(11–12), 1309–1318, doi:10.1016/j.ijpara.2005.06.005.
- Hartl, L., A. Fischer and M. Olefs, 2018: Analysis of past changes in wet bulb temperature in relation to snow making conditions based on long term observations Austria and Germany. *Glob. Planet. Chang.*, **167**, 123–136, doi:10.1016/j.gloplacha.2018.05.011.
- Hassall, C., D.J. Thompson, G.C. French and I.F. Harvey, 2007: Historical changes in the phenology of British Odonata are related to climate. *Glob. Change Biol.*, **13**(5), 933–941, doi:10.1111/j.1365-2486.2007.01318.x.
- Haussig, J., et al., 2018: Early start of the West Nile fever transmission season 2018 in Europe. *Euro Surveill.*, **23**(32), 7–12, doi:10.2807/1560-7917.ES.2018.23.32.1800428.
- Hayashi, N., 2017: The human dimension of climate change research in Greenland: towards a new form of knowledge generation. *Low Temp. Sci.*, **75**, 131–141, doi:10.14943/lowtemsci.75.131.
- Hayashi, N. and M. Walls, 2019: Endogenous community development in Greenland: a perspective on creative transformation and the perception of future. *Polar Sci.*, **21**, 52–57, doi:10.1016/j.polar.2019.06.002.
- Hellmann, F., R. Alkemade and O. Knol, 2016: Dispersal based climate change sensitivity scores for European species. *Ecol. Indic.*, **71**, 41–46, doi:10.1016/j.ecolind.2016.06.013.

- Hendel, M., K. Azos-Diaz and B. Treameac, 2017: Behavioral adaptation to heat-related health risks in cities. *Energy Build.*, **152**, 823–829, doi:10.1016/j.enbuild.2016.11.063.
- Hendel, M., M. Colombert, Y. Diab and L. Royon, 2015: An analysis of pavement heat flux to optimize the water efficiency of a pavement-watering method. *Appl. Therm. Eng.*, **78**, 658–669, doi:10.1016/j.applthermaleng.2014.11.060.
- Hendel, M., et al., 2016: Measuring the effects of urban heat island mitigation techniques in the field: application to the case of pavement-watering in Paris. *Urban Clim.*, **16**, 43–58, doi:10.1016/j.uclim.2016.02.003.
- Hendriks, M., A. van Hinsberg, P. Janssen and B. de Knecht (eds.), 2016: *A species-by-species model to assess anthropogenic impacts on terrestrial biodiversity in Europe*. PBL Netherlands Environmental Assessment Agency, with the cooperation of Wageningen University & Research, Wageningen.
- Hennige, S.J., et al., 2015: Hidden impacts of ocean acidification to live and dead coral framework. *Proc. R. Soc. B.*, **282**(1813), 20150990, doi:10.1098/rspb.2015.0990.
- Heracleous, C. and A. Michael, 2018: Assessment of overheating risk and the impact of natural ventilation in educational buildings of Southern Europe under current and future climatic conditions. *Energy*, **165**, 1228–1239, doi:10.1016/j.energy.2018.10.051.
- Hernández-Morcillo, M., et al., 2018: Scanning agroforestry-based solutions for climate change mitigation and adaptation in Europe. *Environ. Sci. Policy*, **80**(November 2017), 44–52, doi:10.1016/j.envsci.2017.11.013.
- Herrando, S., et al., 2019: Contrasting impacts of precipitation on Mediterranean birds and butterflies. *Sci. Rep.*, **9**(1), 5680, doi:10.1038/s41598-019-42171-4.
- Herrera, M., W. Ds, E. Vázquez and G. Macho, 2019: Climate change implications for reproductive success: temperature effect on penis development in the barnacle *Semibalanus balanoides*. *Mar. Ecol. Prog. Ser.*, **610**, 109–123.
- Herrmann, J. and E. Guenther, 2017: Exploring a scale of organizational barriers for enterprises' climate change adaptation strategies. *J. Clean. Prod.*, **160**, 38–49, doi:10.1016/j.jclepro.2017.03.009.
- Herrmann, M., C. Estournel, F. Adloff and F. Diaz, 2014a: Impact of climate change on the northwestern Mediterranean Sea pelagic planktonic ecosystem and associated carbon cycle. *J. Geophys. Res. Oceans*, **119**(9), 5815–5836, doi:10.1002/2014JC010016.
- Herrmann, T.M., et al., 2014b: Effects of mining on reindeer/caribou populations and indigenous livelihoods: community-based monitoring by Sami reindeer herders in Sweden and First Nations in Canada. *Polar J.*, **4**(1), 28–51, doi:10.1080/2154896X.2014.913917.
- Hertig, E., 2019: Distribution of Anopheles vectors and potential malaria transmission stability in Europe and the Mediterranean area under future climate change. *Parasites Vectors*, **12**, doi:10.1186/s13071-018-3278-6.
- Hidalgo-Galvez, M.D., et al., 2018: Phenological behaviour of early spring flowering trees in Spain in response to recent climate changes. *Theor. Appl. Climatol.*, **132**(1), 263–273, doi:10.1007/s00704-017-2089-6.
- Hidalgo, M., et al., 2019: Accounting for ocean connectivity and hydroclimate in fish recruitment fluctuations within transboundary metapopulations. *Ecol. Appl.*, **29**(5), doi:10.1002/eap.1913. PMID - 31144784.
- Hiddink, J.G., M.T. Burrows and J. García Molinos, 2015: Temperature tracking by North Sea benthic invertebrates in response to climate change. *Glob. Change Biol.*, **21**(1), 117–129, doi:10.1111/gcb.12726.
- Hill, M.O. and C.D. Preston, 2015: Disappearance of boreal plants in southern Britain: habitat loss or climate change? *Biol. J. Linn. Soc.*, **115**(3), 598–610, doi:10.1111/bij.12500.
- Hillebrand, H., et al., 2018: Biodiversity change is uncoupled from species richness trends: consequences for conservation and monitoring. *J. Appl. Ecol.*, **55**(1), 169–184, doi:10.1111/1365-2664.12959.
- Himanen, S.J., H. Mäkinen, K. Rimhanen and R. Savikko, 2016: Engaging farmers in climate change adaptation planning: assessing intercropping as a means to support farm adaptive capacity. *Agriculture*, **6**(3), doi:10.3390/agriculture6030034.
- Hinojosa, J.C., et al., 2019: *Erebia epiphron* and *Erebia orientalis*: sibling butterfly species with contrasting histories. *Biol. J. Linn. Soc.*, **126**(2), 338–348, doi:10.1093/biolinnean/bly182.
- Hintz, M.J., C. Luederitz, D.J. Lang and H. von Wehrden, 2018: Facing the heat: a systematic literature review exploring the transferability of solutions to cope with urban heat waves. *Urban Clim.*, **24**, 714–727, doi:10.1016/j.uclim.2017.08.011.
- Hjerne, O., et al., 2019: Climate driven changes in timing, composition and magnitude of the Baltic Sea phytoplankton spring bloom. *Front. Mar. Sci.*, **6**, 482, doi:10.3389/fmars.2019.00482.
- Hodd, R.L., D. Bourke and M.S. Skeffington, 2014: Projected range contractions of European protected oceanic montane plant communities: focus on climate change impacts is essential for their future conservation. *PLoS ONE*, **9**(4), e95147.
- Hoffmann, I., 2013: Adaptation to climate change – exploring the potential of locally adapted breeds. *Animal*, **7**, 346–362, doi:10.1017/S1757131113000815.
- Hofstede, J.L.A., 2019: On the feasibility of managed retreat in the Wadden Sea of Schleswig-Holstein. *J. Coast. Conserv.*, **23**(6), 1069–1079, doi:10.1007/s11852-019-00714-x.
- Holbrook, N.J., et al., 2019: A global assessment of marine heatwaves and their drivers. *Nat. Commun.*, **10**(1), 2624, doi:10.1038/s41467-019-10206-z.
- Holman, I.P., C. Brown, V. Janes and D. Sanders, 2017: Can we be certain about future land use change in Europe? A multi-scenario, integrated-assessment analysis. *Agric. Syst.*, **151**, 126–135, doi:10.1016/j.agry.2016.12.001.
- Holman, I.P., P.A. Harrison and M.J. Metzger, 2016: *Cross-sectoral Impacts of Climate and Socio-economic Change in Scotland: Implications for Adaptation Policy*. Springer, Berlin, Heidelberg.
- Holt, J., et al., 2018: Climate-driven change in the North Atlantic and Arctic oceans can greatly reduce the circulation of the North Sea. *Geophys. Res. Lett.*, **45**(21), 11,827–11,836, doi:10.1029/2018GL078878.
- Holt, J., et al., 2016: Potential impacts of climate change on the primary production of regional seas: a comparative analysis of five European seas. *Prog. Oceanogr.*, **140**, 91–115, doi:10.1016/j.pocean.2015.11.004.
- Holzämper, A., 2020: Varietal adaptations matter for agricultural water use – a simulation study on grain maize in Western Switzerland. *Agric. Water Manag.*, **237**, 106202–106202, doi:10.1016/j.agwat.2020.106202.
- Hosseinizadehtalaei, P., H. Tabari and P. Willems, 2020: Satellite-based data driven quantification of pluvial floods over Europe under future climatic and socioeconomic changes. *Sci. Total Environ.*, **721**, 137688–137688, doi:10.1016/j.scitotenv.2020.137688.
- Hubble, D.S., 2014: *A Review of the Scarce and Threatened Beetles of Great Britain: The Leaf Beetles and their Allies Chrysomelidae, Megalopodidae and Orsodacnidae Species Status, No.19*. Natural England Commissioned Report, Vol. NECR161, Natural England, Worcester.
- Hudson, P., 2018: A comparison of definitions of affordability for flood risk adaption measures: a case study of current and future risk-based flood insurance premiums in Europe. *Mitig. Adapt. Strateg. Glob. Chang.*, **23**(7), 1019–1038, doi:10.1007/s11027-017-9769-5.
- Hudson, P., W. Botzen, L. Feyen and J. Aerts, 2016: Incentivising flood risk adaptation through risk based insurance premiums: trade-offs between affordability and risk reduction. *Ecol. Econ.*, **125**, 1–13, doi:10.1016/j.ecolecon.2016.01.015.
- Hudson, P., et al., 2014: Evaluating the effectiveness of flood damage mitigation measures by the application of propensity score matching. *Nat. Hazards Earth Syst. Sci.*, **14**(7), 1731–1747, doi:10.5194/nhess-14-1731-2014.
- Hudson, P., W.J.W. Botzen, J. Poussin and J.C.J.H. Aerts, 2019: Impacts of flooding and flood preparedness on subjective well-being: a monetisation of the tangible and intangible impacts. *J. Happiness Stud.*, **20**(2), 665–682, doi:10.1007/s10902-017-9916-4.
- Huete-Stauffer, C., et al., 2011: *Paramuricea clavata* (Anthozoa, Octocorallia) loss in the marine protected area of Tavolara (Sardinia, Italy) due to a

- mass mortality event. *Mar. Ecol.*, **32**(Suppl), 107–116, doi:10.1111/j.1439-0485.2011.00429.x.
- Humphrey, V., et al., 2018: Sensitivity of atmospheric CO₂ growth rate to observed changes in terrestrial water storage. *Nature*, **560**(7720), 628–631, doi:10.1038/s41586-018-0424-4.
- Hunt, A., et al., 2017: Climate and weather service provision: economic appraisal of adaptation to health impacts. *Clim. Serv.*, **7**, 78–86, doi:10.1016/j.cliser.2016.10.004.
- Huntington, H.P., A. Begossi, S. Fox Gearheard, B. Kersey, P.A. Loring, T. Mustonen, P.K. Paudel, R.A.M. Silvano, and R. Vave, R., 2017: How small communities respond to environmental change: patterns from tropical to polar ecosystems. *Ecol. Soc.*, **22**(3), 9, doi:10.5751/ES-09171-220309.
- Huttunen, I., et al., 2021: Agricultural nutrient loading under alternative climate, societal and manure recycling scenarios. *Sci. Total Environ.*, **783**, 146871, doi:10.1016/j.scitotenv.2021.146871.
- IAEA, 2019: *Adapting the Energy Sector to Climate Change*. INTERNATIONAL ATOMIC ENERGY AGENCY, Vienna.
- Ibrahim, A. and S.L.J. Pelsmakers, 2018: Low-energy housing retrofit in North England: overheating risks and possible mitigation strategies. *Build. Serv. Eng. Res. Technol.*, **39**(2), 161–172, doi:10.1177/0143624418754386.
- Iglesias, A. and L. Garrote, 2015: Adaptation strategies for agricultural water management under climate change in Europe. *Agric. Water Manag.*, **155**, 113–124, doi:10.1016/j.agwat.2015.03.014.
- Iglesias, A., D. Santillán and L. Garrote, 2018: On the barriers to adaption to less water under climate change: policy choices in mediterranean countries. *Water Resour. Manag.*, **32**(15), 4819–4832, doi:10.1007/s11269-018-2043-0.
- Ikpewe, I.E., A.R. Baudron, A. Ponchon and P.G. Fernandes, 2021: Bigger juveniles and smaller adults: changes in fish size correlate with warming seas. *J. Appl. Ecol.*, **58**(4), 847–856, doi:10.1111/1365-2664.13807.
- Inuit Circumpolar Council, 2020: *Food Sovereignty and Self-governance: Inuit Role in Managing Arctic Marine Resources*. Anchorage, AK, https://www.culturalsurvival.org/sites/default/files/FSSG%20Report_%20LR%20%281%29.pdf. Accessed 2021.
- Ipbcs, 2016: *The Assessment Report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on Pollinators, Pollination and Food Production*, 552–552. ISBN 978-9280735673. S.G. Potts, V. L. Imperatriz-Fonseca, and H. T. Ngo (eds).
- IPBES, 2018: *The Regional Assessment Report on Biodiversity and Ecosystem Services for Europe and Central Asia* [Rounsevell, M., M. Fischer, A. Torre-Marín Rando and A. Mader (eds.)]. IPBES Secretariat, Bonn, Germany. 892 pp.
- IPCC, 2019: *Special Report: The Ocean and Cryosphere in a Changing Climate* [Pörtner, H.-O., D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, B. Rama and N. Weyer (eds.)]. Cambridge University Press, Cambridge, 1170 pp.
- IRIS LIFE project, 2019: *Guidelines for a Resilient Industry*. <http://www.lifeiris.eu/wp-content/uploads/2019/05/Guidelines.pdf>. Accessed 2021 (24 pp).
- Ito, A., et al., 2020: Pronounced and unavoidable impacts of low-end global warming on northern high-latitude land ecosystems. *Environ. Res. Lett.*, **15**(4), 44006, doi:10.1088/1748-9326/ab702b.
- Iwamura, T., A. Guzman-Holst and K. A. Murray, 2020: Accelerating invasion potential of disease vector *Aedes aegypti* under climate change. *Nat. Commun.*, **11**(1), doi:10.1038/s41467-020-16010-4.
- Jaenson, T., et al., 2012: Changes in the geographical distribution and abundance of the tick *Ixodes ricinus* during the past 30 years in Sweden. *Parasites Vectors*, **5**, doi:10.1186/1756-3305-5-8.
- Jaenson, T. and E. Lindgren, 2011: The range of *Ixodes ricinus* and the risk of contracting Lyme borreliosis will increase northwards when the vegetation period becomes longer. *Ticks Tick Borne Dis.*, **2**(1), 44–49, doi:10.1016/j.ttbdis.2010.10.006.
- Jäger, H., G. Peratoner, U. Tappeiner and E. Tasser, 2020: Grassland biomass balance in the European Alps: current and future ecosystem service perspectives. *Ecosyst. Serv.*, **45**(July), 101163–101163, doi:10.1016/j.ecoser.2020.101163.
- Jaime, L., et al., 2019: Scots pine (*Pinus sylvestris* L.) mortality is explained by the climatic suitability of both host tree and bark beetle populations. *For. Ecol. Manag.*, **448**, 119–129, doi:10.1016/j.foreco.2019.05.070.
- Jaime, R., J.M. Alcántara, A.J. Manzaneda and P.J. Rey, 2018: Climate change decreases suitable areas for rapeseed cultivation in Europe but provides new opportunities for white mustard as an alternative oilseed for biofuel production. *PLoS ONE*, **13**, e207124, doi:10.1371/journal.pone.0207124.
- Jakoby, O., H. Lischke and B. Wermelinger, 2019a: Climate change alters elevational phenology patterns of the European spruce bark beetle (*Ips typographus*). *Glob. Change Biol.*, **25**(12), 4048–4063, doi:10.1111/gcb.14766.
- Jakoby, O., H. Lischke and B. Wermelinger, 2019b: Climate change alters elevational phenology patterns of the European spruce bark beetle (*Ips typographus*). *Glob. Change Biol.*, **25**(11), 4048–4063, doi:10.1111/gcb.14766.
- Jarić, I., et al., 2019: Susceptibility of European freshwater fish to climate change: species profiling based on life-history and environmental characteristics. *Glob. Change Biol.*, **25**(2), 448–458, doi:10.1111/gcb.14518.
- Jenkins, K., et al., 2014a: Implications of climate change for thermal discomfort on underground railways. *Transp. Res. Part D Transp. Environ.*, **30**, 1–9, doi:10.1016/j.trd.2014.05.002.
- Jenkins, K., et al., 2014b: Probabilistic spatial risk assessment of heat impacts and adaptations for London. *Clim. Chang.*, **124**(1), 105–117, doi:10.1007/s10584-014-1105-4.
- Jerez, S., et al., 2015: The impact of climate change on photovoltaic power generation in Europe. *Nat. Commun.*, **6**, doi:10.1038/ncomms10014.
- Jessen, G.L., et al., 2017: Hypoxia causes preservation of labile organic matter and changes seafloor microbial community composition (Black Sea). *Sci. Adv.*, **3**(2), e1601897–e1601897, doi:10.1126/sciadv.1601897.
- Jiang, L., et al., 2020: Effects of sea-level rise on tides and sediment dynamics in a Dutch tidal bay. *Ocean Sci.*, **16**, 307–321.
- Jiguet, F., V. Devictor, R. Ottvall, C. Van Turnhout, H. Van der Jeugd and Å. Lindström, 2010: Bird population trends are linearly affected by climate change along species thermal ranges. *Proc. R. Soc. B*, **277**(1700), 3601–3608, doi:10.1098/rspb.2010.0796.
- Jiménez, J.A. and H.I. Valdemoro, 2019: Shoreline evolution and its management implications in beaches along the Catalan coast. In: *The Spanish Coastal Systems: Dynamic Processes, Sediments and Management*. Springer International Publishing, Cham, pp. 745–764. ISBN 978-3319931692.
- Jiménez, J.A., et al., 2017: Impacts of sea-level rise-induced erosion on the Catalan coast. *Reg. Environ. Chang.*, **17**(2), 593–603, doi:10.1007/s10113-016-1052-x.
- Johansson, C., V.A. Pohjola, C. Jonasson and T.V. Callaghan, 2011: Multi-decadal changes in snow characteristics in sub-arctic Sweden. *Ambio*, **40**(6), 566, doi:10.1007/s13280-011-0164-2.
- Jokinen, S., J.J. Virtasalo, T.S. Jilbert and J. Kaiser, 2018: A 1500-year multiproxy record of coastal hypoxia from the northern Baltic Sea indicates unprecedented deoxygenation over the 20th century. *Biogeosciences*, **15**, 3975–4001, doi:10.1016/S0016-7037(00)00539-1.
- Jolly, W.M., et al., 2015: Climate-induced variations in global wildfire danger from 1979 to 2013. *Nat. Commun.*, **6**(1), 7537, doi:10.1038/ncomms8537.
- Jones, K., et al., 2017: Built asset management climate change adaptation model. *IJRBE*, **8**(3), 263–274, doi:10.1108/IJRBE-07-2016-0032.
- Jongman, B., et al., 2014: Increasing stress on disaster-risk finance due to large floods. *Nat. Clim. Chang.*, **4**(4), 264–268, doi:10.1038/NCLIMATE2124.
- Jonsson, P.R., et al., 2018: High climate velocity and population fragmentation may constrain climate-driven range shift of the key habitat former *Fucus vesiculosus*. *Divers. Distrib.*, **24**(7), 892–905, doi:10.1111/ddi.12733.
- Jore, S., et al., 2014: Climate and environmental change drives *Ixodes ricinus* geographical expansion at the northern range margin. *Parasites Vectors*, **7**, doi:10.1186/1756-3305-7-11.

- Jørgensen, P.S., et al., 2016: Continent-scale global change attribution in European birds - combining annual and decadal time scales. *Glob. Change Biol.*, **22**(2), 530–543, doi:10.1111/gcb.13097.
- Jørgensen, S.L., M. Termansen and U. Pascual, 2020: Natural insurance as condition for market insurance: climate change adaptation in agriculture. *Ecol. Econ.*, **169**, 106489–106489, doi:10.1016/j.ecolecon.2019.106489.
- Joye, J.-F., 2018: Tourism development and adaptation to climate change through legal constraint. *Worldw. Hosp. Tour. Themes*, **10**(2), 244–252, doi:10.1108/WHATT-12-2017-0074.
- Jurt, C., et al., 2015a: Cultural values of glaciers. In: *The High-Mountain Cryosphere: Environmental Changes and Human Risks* [Huggel, C., M. Carey, J.J. Clague and A. Kaab(eds.)]. Cambridge University Press, Cambridge, pp. 90–106. ISBN 978-1107588653.
- Jurt, C., et al., 2015b: Local perceptions in climate change debates: insights from case studies in the Alps and the Andes. *Clim. Chang.*, **133**(3), 511–523, doi:10.1007/s10584-015-1529-5.
- Juschten, M., et al., 2019a: Out of the city heat—way to less or more sustainable futures? *Sustainability*, **11**(1), 214.
- Juschten, M., A. Jiricka-Pürner, W. Unbehaun and R. Hössinger, 2019b: The mountains are calling! An extended TPB model for understanding metropolitan residents' intentions to visit nearby alpine destinations in summer. *Tour. Manag.*, **75**, 293–306, doi:10.1016/j.tourman.2019.05.014.
- Kahil, M.T., A. Dinar and J. Albiac, 2015: Modeling water scarcity and droughts for policy adaptation to climate change in arid and semiarid regions. *J. Hydrol. Reg. Stud.*, **522**, 95–109, doi:10.1016/j.jhydrol.2014.12.042.
- Kaiser, N., et al., 2010: Depression and anxiety in the reindeer-herding Sami population of Sweden. *Int. J. Circumpolar Health*, **69**(4), 383–393, doi:10.3402/ijch.v69i4.17674.
- Kaloveloni, A., et al., 2015: Winners and losers of climate change for the genus *Merodon* (Diptera: Syrphidae) across the Balkan Peninsula. *Ecol. Model.*, **313**, 201–211, doi:10.1016/j.ecolmodel.2015.06.032.
- Kärcher, O., D. Hering, K. Frank and D. Markovic, 2019: Freshwater species distributions along thermal gradients. *Ecol. Evol.*, **9**(1), 111–124, doi:10.1002/ece3.4659.
- Karkanis, A., et al., 2018: Interference of weeds in vegetable crop cultivation, in the changing climate of Southern Europe with emphasis on drought and elevated temperatures: a review. *J. Agric. Sci.*, **156**(10), 1175–1185, doi:10.1017/S0021859619000108.
- Karlsson, B., 2014: Extended season for northern butterflies. *Int. J. Biometeorol.*, **58**, doi:10.1007/s00484-013-0649-8.
- Karypidou, M.C., et al., 2020: Projected shifts in the distribution of malaria vectors due to climate change. *Clim. Chang.*, **163**(4), 2117–2133, doi:10.1007/s10584-020-02926-9.
- Katopodis, T., et al., 2019: Assessment of climate change impacts on wind resource characteristics and wind energy potential in Greece. *J. Renew. Sustain. Energy*, **11**(6), 66502, doi:10.1063/1.5118878.
- Kayaga, S. and I. Smout, 2014: Tariff structures and incentives for water demand management. *Proc. Inst. Civ. Eng. Water Manag.*, **167**(8), 448–456, doi:10.1680/wama.12.00120.
- Kebede, A.S., et al., 2015: Direct and indirect impacts of climate and socio-economic change in Europe: a sensitivity analysis for key land- and water-based sectors. *Clim. Chang.*, **128**(3), 261–277, doi:10.1007/s10584-014-1313-y.
- Kebede, A.S., et al., 2021: Integrated assessment of the food-water-land-ecosystems nexus in Europe: implications for sustainability. *Sci. Total Environ.*, **768**, 144461–144461, doi:10.1016/j.scitotenv.2020.144461.
- Keenan, T.F., et al., 2016: Recent pause in the growth rate of atmospheric CO₂ due to enhanced terrestrial carbon uptake. *Nat. Commun.*, **7**, 1–9, doi:10.1038/ncomms13428.
- Kendrovski, V., et al., 2017: Quantifying projected heat mortality impacts under 21st-century warming conditions for selected European countries. *Int. J. Environ. Res. Public Health*, **14**(7), 729, doi:10.3390/ijerph14070729.
- Keogan, K., et al., 2021: No evidence for fitness signatures consistent with increasing trophic mismatch over 30 years in a population of European shag *Phalacrocorax aristotelis*. *J. Anim. Ecol.*, **90**(2), 432–446, doi:10.1111/1365-2656.13376.
- Kerr, J.T., et al., 2015: Climate change impacts on bumblebees converge across continents. *Science*, **349**(6244), 177–180, doi:10.1126/science.aaa7031.
- Kersting, D.K., N. Bensoussan and C. Linares, 2013: Long-term responses of the endemic reef-builder *Cladocora caespitosa* to Mediterranean warming. *PLoS ONE*, **8**(8), doi:10.1371/journal.pone.0070820.
- Keskitalo, E., G. Vulturius and P. Scholten, 2014: Adaptation to climate change in the insurance sector: examples from the UK, Germany and the Netherlands. *Nat. Hazards*, **71**(1), 315–334, doi:10.1007/s11069-013-0912-7.
- Khabarov, N., et al., 2016: Forest fires and adaptation options in Europe. *Reg. Environ. Chang.*, **16**(1), 21–30, doi:10.1007/s10113-014-0621-0.
- Khan, Z., P. Linares and J. García-González, 2016: Adaptation to climate-induced regional water constraints in the Spanish energy sector: an integrated assessment. *Energy Policy*, **97**, 123–135, doi:10.1016/j.enpol.2016.06.046.
- Khare, S., et al., 2015: Heat protection behaviour in the UK: results of an online survey after the 2013 heatwave. *BMC Public Health*, **15**(1), 878, doi:10.1186/s12889-015-2181-8.
- Kingsborough, A., E. Borgomeo and J.W. Hall, 2016: Adaptation pathways in practice: mapping options and trade-offs for London's water resources. *Sustain. Cities Soc.*, **27**, 386–397, doi:10.1016/j.scs.2016.08.013.
- Kingsborough, A., K. Jenkins and J.W. Hall, 2017: Development and appraisal of long-term adaptation pathways for managing heat-risk in London. *Clim. Risk Manag.*, **16**, 73–92, doi:10.1016/j.crm.2017.01.001.
- Kivinen, S., et al., 2012: Forest fragmentation and landscape transformation in a reindeer husbandry area in Sweden. *Environ. Manag.*, **49**(2), 295–304, doi:10.1007/s00267-011-9788-z.
- Kjesbu, O.S., et al., 2014: Synergies between climate and management for Atlantic cod fisheries at high latitudes. *Proc. Natl. Acad. Sci.*, **111**(9), 3478–3483, doi:10.1073/pnas.1316342111.
- Klein, G., et al., 2016: Shorter snow cover duration since 1970 in the Swiss Alps due to earlier snowmelt more than to later snow onset. *Clim. Chang.*, **139**(3), 637–649, doi:10.1007/s10584-016-1806-y.
- Klimenko, V.V., E. V. Fedotova and A.G. Tereshin, 2018: Vulnerability of the Russian power industry to the climate change. *Energy*, **142**, 1010–1022, doi:10.1016/j.energy.2017.10.069.
- Knittel, N., et al., 2020: A global analysis of heat-related labour productivity losses under climate change—implications for Germany's foreign trade. *Clim. Chang.*, **160**(2), 251–269, doi:10.1007/s10584-020-02661-1.
- Knox, J., A. Daccache, T. Hess and D. Haro, 2016: Meta-analysis of climate impacts and uncertainty on crop yields in Europe. *Environ. Res. Lett.*, **11**(11), 113004, doi:10.1088/1748-9326/11/11/113004.
- Koch, H., et al., 2014: Security of water supply and electricity production: aspects of integrated management. *Water Resour. Manag.*, **28**(6), 1767–1780, doi:10.1007/s11269-014-0589-z.
- Koerth, J., et al., 2013: Household adaptation and intention to adapt to coastal flooding in the Axios – Loudias – Aliakmonas National Park, Greece. *Ocean. Coast. Manag.*, **82**, 43–50, doi:10.1016/j.ocecoaman.2013.05.008.
- Koks, E.E., et al., 2019: The macroeconomic impacts of future river flooding in Europe. *Environ. Res. Lett.*, **14**(8), 84042, doi:10.1088/1748-9326/ab3306.
- Koletsis, I., V. Kotroni, K. Lagouvardos and T. Soukissian, 2016: Assessment of offshore wind speed and power potential over the Mediterranean and the Black Seas under future climate changes. *Renew. Sustain. Energy Rev.*, **60**, 234–245, doi:10.1016/j.rser.2016.01.080.
- Koopman, J.F.L., O. Kuik, R.S.J. Tol and R. Brouwer, 2017: The potential of water markets to allocate water between industry, agriculture, and public water utilities as an adaptation mechanism to climate change. *Mitig. Adapt. Strateg. Glob. Chang.*, **22**(2), 325–347, doi:10.1007/s11027-015-9662-z.
- Kortsch, S., et al., 2015: Climate change alters the structure of arctic marine food webs due to poleward shifts of boreal generalists. *Proc. R. Soc. B.*, **282**(1814), 20151546–20151546, doi:10.1098/rspb.2015.1546.

- Kotenev, B.N., et al., 2017: Development prospects of world fisheries in relation to climate change. *Sci. Notes Russ. State Hydrometeorol. Univ.*, **48**, 167–185.
- Kotta, J., et al., 2018: Novel crab predator causes marine ecosystem regime shift. *Sci. Rep.*, **8**(1), 4956–4956, doi:10.1038/s41598-018-23282-w.
- Kougioumoutzis, K., et al., 2020: Plant diversity patterns and conservation implications under climate-change scenarios in the Mediterranean: the case of crete (Aegean, Greece). *Diversity*, **12**(7), doi:10.3390/d12070270.
- Koutroulis, A.G., et al., 2019: Global water availability under high-end climate change: a vulnerability based assessment. *Glob. Planet. Chang.*, **175**, 52–63, doi:10.1016/j.gloplacha.2019.01.013.
- Kovats, R.S., R. Valentini, L.M. Bouwer, E. Georgopoulou, D. Jacob, E. Martin, M. Rounsevell, M. and J.F. Soussana, 2014: Europe. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*. [Barros, V. R., C. B. Field, D. J. Dokken, M. D. Mastrandrea, K. J. Mach, T. E. Bilir, M. Chatterjee, K. L. Ebi, Y. O. Estrada, R. C. Genova, B. Girma, E. S. Kissel, A. N. Levy, S. MacCracken, P. R. Mastrandrea and L. L. White (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 1267–1326.
- Kraemer, M., et al., 2015: The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. *Trop. Med. Int. Health*, **20**, 38–38.
- Krause, A., T. Knoke and A. Rammig, 2020: A regional assessment of land-based carbon mitigation potentials: bioenergy, BECCS, reforestation, and forest management. *Glob. Change Biol. Bioenergy*, **12**(5), 346–360, doi:10.1111/gcbb.12675.
- Kreibich, H., P. Bubeck, M. Van Vliet and H. De Moel, 2015: A review of damage-reducing measures to manage fluvial flood risks in a changing climate. *Mitig. Adapt. Strateg. Glob. Change*, **20**(6), 967–989, doi:10.1007/s11027-014-9629-5.
- Kreibich, H., et al., 2017: Adaptation to flood risk: Results of international paired flood event studies: Adaptation to flood risk. *Earth's Future*, **5**(10), 953–965, doi:10.1002/2017EF000606.
- Kreibich, H., P. Hudson and B. Merz, 2021: Knowing what to do substantially improves the effectiveness of flood early warning. *Bull. Am. Meteorol. Soc.*, **102**(7), E1450–E1463, doi:10.1175/BAMS-D-20-0262.1.
- Kröncke, I., et al., 2019: Comparison of biological and ecological long-term trends related to northern hemisphere climate in different marine ecosystems. *Nat. Conserv.*, **34**, 311–341, doi:10.3897/natureconservation.34.30209.
- Krovnin, A.S., S.P. Melnikov, D.V. Artemenkov and G.P. Moury, 2019: Climate change impact on fish communities in the North Atlantic region. In: *Modern Problems of Hydrometeorology and Sustainable Development of the Russian Federation* [Mikheev, V.L., I.I. Musket, A.A. Ershova and A.A. Fokicheva (eds.)]. Russian State Hydrometeorological University, Saint-Petersburg, pp. 382–383.
- Krumhansl, K.A., et al., 2016: Global patterns of kelp forest change over the past half-century. *Proc. Natl. Acad. Sci.*, **113**(48), 13785–13790, doi:10.1073/pnas.1606102113.
- Kundzewicz, Z.W., I. Pińskwar and G.R. Brakenridge, 2017: Changes in river flood hazard in Europe: a review. *Hydrol. Res.*, **49**(2), 294–302, doi:10.2166/nh.2017.016.
- Kwiatkowski, L., O. Aumont and L. Bopp, 2019: Consistent trophic amplification of marine biomass declines under climate change. *Glob. Change Biol.*, **25**(1), 218–229, doi:10.1111/gcb.14468.
- Lacoue-Labarthe, T., et al., 2016: Impacts of ocean acidification in a warming Mediterranean sea: an overview. *Reg. Stud. Mar. Sci.*, **5**, 1–11, doi:10.1016/j.rsma.2015.12.005.
- Lahaye, S., et al., 2018: What are the drivers of dangerous fires in Mediterranean France? *Int. J. Wildland Fire*, **27**(3), 155–163.
- Lam, V.W.Y., W.W.L. Cheung, G. Reygondeau and U.R. Sumaila, 2016: Projected change in global fisheries revenues under climate change. *Sci. Rep.*, **6**(1), 32607, doi:10.1038/srep32607.
- Lamichhane, J.R., J. Constantin, J.N. Aubertot and C. Dürr, 2019: Will climate change affect sugar beet establishment of the 21st century? Insights from a simulation study using a crop emergence model. *Field Crop. Res.*, **238**, 64–73, doi:10.1016/j.fcr.2019.04.022.
- Landauer, M., M. E. Goodsite and S. Juhola, 2018: Nordic national climate adaptation and tourism strategies – (how) are they interlinked? *Scand. J. Hosp. Tour.*, **18**(sup1), S75–S86, doi:10.1080/15022250.2017.1340540.
- Landauer, M., W. Haider and U. Pröbstl-Haider, 2013: The influence of culture on climate change adaptation strategies: preferences of cross-country skiers in Austria and Finland. *J. Travel Res.*, **53**(1), 96–110, doi:10.1177/0047287513481276.
- Lange, S., et al., 2020: Projecting exposure to extreme climate impact events across six event categories and three spatial scales. *Earths Future*, **8**(12), e2020EF001616–e002020EF001616, doi:10.1029/2020EF001616.
- Larsen, R.K. and K. Raitio, 2019: Implementing the state duty to consult in land and resource decisions: perspectives from Sami communities and Swedish state officials. *Arct. Rev. Law Polit.*, **10**(0), 4, doi:10.23865/arctic.v10.1323.
- Latchinsky, A.V., 2017: Climate changes and locusts: what to expect? *Sci. Notes Russ. State Hydrometeorol. Univ.*, **46**, 134–143.
- Laufkotter, C., et al., 2015: Drivers and uncertainties of future global marine primary production in marine ecosystem models. *Biogeosciences*, **12**(23), 6955–6984, doi:10.5194/bg-12-6955-2015.
- Lavorel, S., et al., 2019: Mustering the power of ecosystems for adaptation to climate change. *Environ. Sci. Policy*, **92**, 87–97, doi:10.1016/j.envsci.2018.11.010.
- Lavorel, S., B. Locatelli, M.J. Colloff and E. Bruley, 2020: Co-producing ecosystem services for adapting to climate change. *Philos. Trans. R. Soc. B Biol. Sci.*, **375**(1794), 20190119, doi:10.1098/rstb.2019.0119.
- Lawrence, R., 2014: Internal Colonisation and indigenous resource sovereignty: wind power developments on traditional Saami lands. *Environ Plan D*, **32**(6), 1036–1053, doi:10.1068/d9012.
- Lawrence, R. and R. Kløcker Larsen, 2017: The politics of planning: assessing the impacts of mining on Sami lands. *Third World Q.*, **38**(5), 1164–1180, doi:10.1080/01436597.2016.1257909.
- Lawrence, R. and R. Kløcker Larsen, 2019: *Fighting to Be Herd: Impacts of the Proposed Boliden Copper Mine in Laver, Älvsbyn, Sweden for the Semisjaur Njarg Sami Reindeer Herding Community*. Stockholm Environment Institute, Stockholm, <https://www.sei.org/wp-content/uploads/2019/04/sei-report-fighting-to-be-herd-300419.pdf>. (96 pp).
- Leclère, D., P.-A. Jayet and N. de Noblet-Ducoudré, 2013: Farm-level autonomous adaptation of European agricultural supply to climate change. *Ecol. Econ.*, **87**, 1–14, doi:10.1016/j.ecolecon.2012.11.010.
- Lee, H., et al., 2019: Implementing land-based mitigation to achieve the Paris Agreement in Europe requires food system transformation. *Environ. Res. Lett.*, **14**(10), 104009, doi:10.1088/1748-9326/ab3744.
- Lee, W., et al., 2020: Projections of excess mortality related to diurnal temperature range under climate change scenarios: a multi-country modelling study. *Lancet Planet. Health*, **4**(11), e512–e521, doi:10.1016/S2542-5196(20)30222-9.
- Lehikoinen, A., et al., 2019a: Declining population trends of European mountain birds. *Glob. Change Biol.*, **25**(2), 577–588, doi:10.1111/gcb.14522.
- Lehikoinen, A., et al., 2019b: Phenology of the avian spring migratory passage in Europe and North America: Asymmetric advancement in time and increase in duration. *Ecol. Indic.*, **101**, 985–991.
- Leitinger, G., et al., 2015: Impact of droughts on water provision in managed alpine grasslands in two climatically different regions of the Alps. *Ecophysiology*, **8**(8), 1600–1613, doi:10.1002/eco.1607.
- Lenderink, G., et al., 2019: Systematic increases in the thermodynamic response of hourly precipitation extremes in an idealized warming experiment with a convection-permitting climate model. *Environ. Res. Lett.*, **14**(7), 74012, doi:10.1088/1748-9326/ab214a.
- Lenoir, J., et al., 2008: A significant upward shift in plant species optimum elevation during the 20th century. *Science*, **320**(5884), 1768, doi:10.1126/science.1156831.

- Lenoir, J., et al., 2013: Local temperatures inferred from plant communities suggest strong spatial buffering of climate warming across Northern Europe. *Glob. Change Biol.*, **19**(5), 1470–1481, doi:10.1111/gcb.12129.
- Lewis, K.M., G.L. van Dijken and K.R. Arrigo, 2020: Changes in phytoplankton concentration now drive increased Arctic Ocean primary production. *Science*, **369**(6500), 198–202, doi:10.1126/science.aay8380.
- Li, S., et al., 2017: Relating farmer's perceptions of climate change risk to adaptation behaviour in Hungary. *J. Environ. Manag.*, **185**, 21–30, doi:10.1016/j.jenvman.2016.10.051.
- Li, S., et al., 2018: Modelling regional cropping patterns under scenarios of climate and socio-economic change in Hungary. *Sci. Total. Environ.*, **622–623**, 1611–1620, doi:10.1016/j.scitotenv.2017.10.038.
- Lian, X., et al., 2020: Summer soil drying exacerbated by earlier spring greening of northern vegetation. *Sci. Adv.*, **6**(1), eaax255, doi:10.1126/sciadv.aax255.
- Lincke, D., et al., 2020: The effectiveness of setback zones for adapting to sea-level rise in Croatia. *Reg. Environ. Chang.*, **20**(2), doi:10.1007/s10113-020-01628-3.
- Indegren, M., et al., 2018: Productivity and recovery of forage fish under climate change and fishing: North Sea sandeel as a case study. *Fish. Oceanogr.*, **27**(3), 212–221, doi:10.1111/fog.12246.
- Litskas, V.D., et al., 2019: Impacts of climate change on tomato, a notorious pest and its natural enemy: small scale agriculture at higher risk. *Environ. Res. Lett.*, **14**(8), 84041, doi:10.1088/1748-9326/ab3313.
- Litvin, L.F., Z.P. Kiryukhina, S.F. Krasnov and N.G. Dobrovolskaya, 2017: Dynamics of agricultural soil erosion in European Russia. *Eurasian Soil Sci.*, **50**(11), 1344–1353, doi:10.1134/S1064229317110084.
- Liu-Helmersson, J., et al., 2016: Climate change and Aedes vectors: 21st century projections for dengue transmission in Europe. *EBioMedicine*, **7**, 267–277, doi:10.1016/j.ebiom.2016.03.046.
- Liu-Helmersson, J., J. Rocklöv, M. Sewe and A. Brannström, 2019: Climate change may enable Aedes aegypti infestation in major European cities by 2100. *Environ. Res.*, **172**, 693–699, doi:10.1016/j.envres.2019.02.026.
- Liu, J., et al., 2018: Managing the energy-water-food nexus for sustainable development. *Appl. Energy*, **210**, 377–381, doi:10.1016/j.apenergy.2017.10.064.
- Liu, L. and M.B. Jensen, 2018: Green infrastructure for sustainable urban water management: practices of five forerunner cities. *Cities*, **74**, 126–133, doi:10.1016/j.cities.2017.11.013.
- Liu, Y., et al., 2020: Reviewing estimates of the basic reproduction number for dengue, Zika and chikungunya across global climate zones. *Environ. Res.*, **182**, doi:10.1016/j.envres.2020.109114.
- Liverpool, W.M., 2016: *The Status of the Snowdon Beetle Chrysolina Cerealis on Yr Wyddfa in 2015. NRW Evidence Report*.
- Ljungqvist, F.C., et al., 2016: Northern Hemisphere hydroclimate variability over the past twelve centuries. *Nature*, **532**(7597), 94–98, doi:10.1038/nature17418.
- Lloret, J., et al., 2018: Small-scale coastal fisheries in European Seas are not what they were: ecological, social and economic changes. *Mar. Policy*, **98**, 176–186, doi:10.1016/j.marpol.2016.11.007.
- Lobanova, A., et al., 2016: Impacts of changing climate on the hydrology and hydropower production of the Tagus River basin. *Hydrol. Process.*, **30**(26), 5039–5052, doi:10.1002/hyp.10966.
- Löf, A., 2013: Examining limits and barriers to climate change adaptation in an Indigenous reindeer herding community. *Clim. Dev.*, **5**(4), 328–339, doi:10.1080/17565529.2013.831338.
- Löf, A., 2014: *Challenging Adaptability: Analysing the Governance of Reindeer Husbandry in Sweden*. Umeå University, Faculty of Social Sciences, Department of Political Science, Umeå.
- Long, T.B., V. Blok and I. Coninx, 2016: Barriers to the adoption and diffusion of technological innovations for climate-smart agriculture in Europe: evidence from the Netherlands, France, Switzerland and Italy. *J. Clean. Prod.*, **112**, 9–21, doi:10.1016/j.jclepro.2015.06.044.
- López-i-Gelats, F., E.D.G. Fraser, J.F. Morton and M.G. Rivera-Ferre, 2016: What drives the vulnerability of pastoralists to global environmental change? A qualitative meta-analysis. *Glob. Environ. Chang.*, **39**, 258–274, doi:10.1016/j.gloenvcha.2016.05.011.
- Lotze, H.K., et al., 2019a: Global ensemble projections reveal trophic amplification of ocean biomass declines with climate change. *Proc. Natl. Acad. Sci. U.S.A.*, **116**(26), 12907–12912, doi:10.1073/pnas.1900194116.
- Lotze, H.K., et al., 2019b: Global ensemble projections reveal trophic amplification of ocean biomass declines with climate change. *Proc. Natl. Acad. Sci. U.S.A.*, **116**(26), 12907–12912, doi:10.1073/pnas.1900194116.
- Luetzenburg, G., et al., 2020: Climate and land use change effects on soil erosion in two small agricultural catchment systems Fugnitz – Austria, Can Revull – Spain. *Sci. Total. Environ.*, **704**, 135389–135389, doi:10.1016/j.scitotenv.2019.135389.
- Lugato, E., et al., 2018: Soil erosion is unlikely to drive a future carbon sink in Europe. *Sci. Adv.*, **4**(11), eaau3523, doi:10.1126/sciadv.aau3523.
- Lujala, P., H. Lein and J.K. Rød, 2015: Climate change, natural hazards, and risk perception: the role of proximity and personal experience. *Local Environ.*, **20**(4), 489–509, doi:10.1080/13549839.2014.887666.
- Lüscher, A., et al., 2014: Potential of legume-based grassland–livestock systems in Europe: a review. *Grass Forage Sci.*, **69**(2), 206–228, doi:10.1111/gfs.12124.
- Luysaert, S., et al., 2018: Trade-offs in using European forests to meet climate objectives. *Nature*, **562**(7726), 259–262, doi:10.1038/s41586-018-0577-1.
- Ma, Q., J.G. Huang, H. Hänninen and F. Berninger, 2019: Divergent trends in the risk of spring frost damage to trees in Europe with recent warming. *Glob. Change Biol.*, **25**(1), 351–360, doi:10.1111/gcb.14479.
- Macgregor, C.J., et al., 2019: Climate-induced phenology shifts linked to range expansions in species with multiple reproductive cycles per year. *Nat. Commun.*, **10**(1), 4455, doi:10.1038/s41467-019-12479-w.
- Macholdt, J. and B. Honermeier, 2017: Importance of variety choice: adapting to climate change in organic and conventional farming systems in Germany. *Outlook Agric.*, **46**(3), 178–184, doi:10.1177/0030727017722420.
- Macintyre, H. and C. Heaviside, 2019: Potential benefits of cool roofs in reducing heat-related mortality during heatwaves in a European city. *Environ Int.*, **127**, 430–441, doi:10.1016/j.envint.2019.02.065.
- MacLachlan, N.J. and A.J. Guthrie, 2010: Re-emergence of bluetongue, African horse sickness, and other orbivirus diseases. *Vet. Res.*, **41**(6), 35–35, doi:10.1051/vetres/2010007.
- Madsen, H.M., P.S. Mikkelsen and A. Blok, 2019: Framing professional climate risk knowledge: Extreme weather events as drivers of adaptation innovation in Copenhagen, Denmark. *Environ. Sci. Policy*, **98**, 30–38, doi:10.1016/j.envsci.2019.04.004.
- Maier, C., et al., 2013: End of the century pCO₂ levels do not impact calcification in Mediterranean cold-water corals. *PLoS ONE*, **8**(4), e62655–e62655, doi:10.1371/journal.pone.0062655.
- Malagon Santos, V., I.D. Haigh and T. Wahl, 2017: Spatial and temporal clustering analysis of extreme wave events around the UK coastline. *J. Mar. Sci. Eng.*, **5**(3), doi:10.3390/jmse5030028.
- Malcolm, I.A., C.P. Millar and K.J. Millidine, 2015: Spatio-temporal variability in Scottish smolt emigration times and sizes. *Scott. Mar. Freshw. Sci.*, **6**(2), 15.
- Mallory, C.D. and M.S. Boyce, 2018: Observed and predicted effects of climate change on Arctic caribou and reindeer. *Environ. Rev.*, **26**(1), 13–25, doi:10.1139/er-2017-0032.
- Maltby, K.M., et al., 2020: Projected impacts of warming seas on commercially fished species at a biogeographic boundary of the European continental shelf. *J. Appl. Ecol.*, **57**(11), 2222–2233, doi:10.1111/1365-2664.13724.
- Mandel, A., et al., 2021: Risks on global financial stability induced by climate change. *Clim. Change*, **166**, 4, doi:10.1007/s10584-021-03092-2.
- Mandryk, M., et al., 2015: Institutional constraints for adaptive capacity to climate change in Flevoland's agriculture. *Environ. Sci. Policy*, **48**, 147–162, doi:10.1016/j.envsci.2015.01.001.

- Mangi, S.C., et al., 2018: The economic impacts of ocean acidification on shellfish fisheries and aquaculture in the United Kingdom. *Environ. Sci. Policy*, **86**, 95–105, doi:10.1016/j.envsci.2018.05.008.
- Manoussel, D., B. Anderson and M. Nagarajan, 2018: Domestic water demand during droughts in temperate climates: synthesising evidence for an integrated framework. *Water Resour. Manag.*, **32**(2), 433–447, doi:10.1007/s11269-017-1818-z.
- Maragno, D., et al., 2018: Fine-scale analysis of urban flooding reduction from green infrastructure: an ecosystem services approach for the management of water flows. *Ecol. Model.*, **386**, 1–10, doi:10.1016/j.ecolmodel.2018.08.002.
- Margalef-Marrase, J., M.Á. Pérez-Navarro and F. Lloret, 2020: Relationship between heatwave-induced forest die-off and climatic suitability in multiple tree species. *Glob. Change Biol.*, **26**(5), 3134–3146, doi:10.1111/gcb.15042.
- Marini, G., et al., 2021: Spring temperature shapes West Nile virus transmission in Europe. *Acta Trop.*, **215**, doi:10.1016/j.actatropica.2020.105796.
- Marini, G., et al., 2016: The role of climatic and density dependent factors in shaping mosquito population dynamics: the case of *Culex pipiens* in northwestern Italy. *PLoS ONE*, **11**(4), doi:10.1371/journal.pone.0154018.
- Markovic, D., et al., 2014: Europe's freshwater biodiversity under climate change: distribution shifts and conservation needs. *Divers. Distrib.*, **20**(9), 1097–1107, doi:10.1111/ddi.12232.
- Marqués, L., et al., 2018: Last-century forest productivity in a managed dry-edge Scots pine population: the two sides of climate warming. *Ecol. Appl.*, **28**(1), 95–105, doi:10.1002/eap.1631.
- Marshall, L., et al., 2018: The interplay of climate and land use change affects the distribution of EU bumblebees. *Glob. Change Biol.*, **24**(1), 101–116, doi:10.1111/gcb.13867.
- Martin, A.C., et al., 2017: Shrub growth and expansion in the Arctic tundra: an assessment of controlling factors using an evidence-based approach. *Environ. Res. Lett.*, **12**(8), 85007–85007, doi:10.1088/1748-9326/aa7989.
- Martínez-López, B., et al., 2014: Farm-level risk factors for the occurrence, new infection or persistence of tuberculosis in cattle herds from South-Central Spain. *Prev. Vet. Med.*, **116**(3), 268–278, doi:10.1016/j.prevetmed.2013.11.002.
- Martynova, D.V., et al., 2020: The White Sea Pelagic Study: Dynamics of the Main Indicators (Hydrology, Hydrochemistry, Zooplankton) Based on the Results of Marine Expeditions and its Connection with Climate Change. In: *Results of Expeditionary Research in 2019 in the Oceans, Inland Waters*, pp. 26–38.
- Marzeion, B. and A. Levermann, 2014: Loss of cultural world heritage and currently inhabited places to sea-level rise. *Environ. Res. Lett.*, **9**(3), doi:10.1088/1748-9326/9/3/034001.
- Mascarenhas, M., et al., 2018: A scoping review of published literature on chikungunya virus. *PLoS ONE*, **13**(11), doi:10.1371/journal.pone.0207554.
- Massimino, D., et al., 2017: Projected reductions in climatic suitability for vulnerable British birds. *Clim. Chang.*, **145**(1), 117–130, doi:10.1007/s10584-017-2081-2.
- Matzarakis, A., C. Endler and P.T. Nastos, 2014: Quantification of climate-tourism potential for Athens, Greece—recent and future climate simulations. *Glob. NEST J.*, **16**(1), 43–51.
- Matzarakis, A., J. Rammelberg and J. Junk, 2013: Assessment of thermal bioclimate and tourism climate potential for central Europe... the example of Luxembourg. *Theor. Appl. Climatol.*, **114**, 193–202.
- Maugendre, L., et al., 2014: Effect of ocean warming and acidification on a plankton community in the NW Mediterranean Sea. *ICES J. Mar. Sci.*, **72**(6), 1744–1755, doi:10.1093/icesjms/fsu161.
- Maynou, F., A. Sabatés and V. Raya, 2020: Changes in the spawning habitat of two small pelagic fish in the Northwestern Mediterranean. *Fish. Oceanogr.*, **29**(2), 201–213, doi:10.1111/fog.12464.
- Mayor, S.J., et al., 2017: Increasing phenological asynchrony between spring green-up and arrival of migratory birds. *Sci. Rep.*, **7**(1), 1902, doi:10.1038/s41598-017-02045-z.
- Mayr, B., T. Thaler and J. Hübl, 2020: Successful small-scale household relocation after a millennial flood event in Simbach, Germany 2016. *Water (Switzerland)*, **12**(1), doi:10.3390/w12010156.
- McGill, B.J., M. Dornelas, N.J. Gotelli and A.E. Magurran, 2015: Fifteen forms of biodiversity trend in the Anthropocene. *Trends Ecol. Evol.*, **30**(2), 104–113, doi:10.1016/j.tree.2014.11.006.
- McIntyre, K.M., et al., 2017: Systematic assessment of the climate sensitivity of important human and domestic animals pathogens in Europe. *Sci. Rep.*, **7**(1), 7134, doi:10.1038/s41598-017-06948-9.
- McVittie, A., et al., 2018: Ecosystem-based solutions for disaster risk reduction: Lessons from European applications of ecosystem-based adaptation measures. *Int. J. Disaster Risk Reduct.*, **32**, 42–54, doi:10.1016/j.ijdr.2017.12.014.
- Medlock, J.M., et al., 2013: Driving forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. *Parasites Vectors*, **6**(1), 1, doi:10.1186/1756-3305-6-1.
- Medlock, J.M., et al., 2018: Assessment of the public health threats posed by vector-borne disease in the United Kingdom (UK). *Int J Environ Res Public Health*, **15**(10), doi:10.3390/ijerph15102145.
- Medlock, J.M., et al., 2015: An entomological review of invasive mosquitoes in Europe. *Bull. Entomol. Res.*, **105**(6), 637–663, doi:10.1017/s0007485315000103.
- Mekonnen, Z.A., et al., 2021: Arctic tundra shrubification: a review of mechanisms and impacts on ecosystem carbon balance. *Environ. Res. Lett.*, **16**(5), 53001.
- Melero, Y., C. Stefanescu and J. Pino, 2016: General declines in Mediterranean butterflies over the last two decades are modulated by species traits. *Biol. Conserv.*, **201**, 336–342, doi:10.1016/j.biocon.2016.07.029.
- Menéndez, R., et al., 2006: Species richness changes lag behind climate change. *Proc. Biol. Sci.*, **273**(1593), 1465–1470, doi:10.1098/rspb.2006.3484.
- Mentaschi, L., et al., 2018: Global long-term observations of coastal erosion and accretion. *Sci. Rep.*, **8**(1), 12876, doi:10.1038/s41598-018-30904-w.
- Menzel, A., et al., 2020: Climate change fingerprints in recent European plant phenology. *Glob. Change Biol.*, **26**(4), 2599–2612, doi:10.1111/gcb.15000.
- Meredith, M., M. Sommerkorn, S. Cassotta, C. Derksen, A. Ekaykin, A. Hollowed, G. Kofinas, A. Mackintosh, J. Melbourne-Thomas, M.M.C. Muelbert, G. Ottersen, H. Pritchard, and E.A.G. Schuur, 2019: Polar Regions. In: *PCC Special Report on the Ocean and Cryosphere in a Changing Climate*. [Pörtner, H.-O., D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegria, M. Nicolai, A. Okem, J. Petzold, B. Rama and N. M. Weyer (eds.)]. Cambridge University Press, Cambridge, pp. 203–320.
- Merz, B., et al., 2021: Causes, impacts and patterns of disastrous river floods. *Nat. Rev. Earth Environ.*, **2**(9), 592–609, doi:10.1038/s43017-021-00195-3.
- Merz, B., et al., 2020: Impact forecasting to support emergency management of natural hazards. *Rev. Geophys.*, **58**(4), doi:10.1029/2020RG000704.
- Messina, J.P., et al., 2019: The current and future global distribution and population at risk of dengue. *Nat. Microbiol.*, **4**(9), 1508–1515, doi:10.1038/s41564-019-0476-8.
- Metelmann, S., et al., 2019: The UK's suitability for *Aedes albopictus* in current and future climates. *J. R. Soc. Interface.*, **16**(152), doi:10.1098/rsif.2018.0761.
- Metin, A.D., et al., 2018: How do changes along the risk chain affect flood risk? *Nat. Hazards Earth Syst. Sci.*, **18**(11), 3089–3108, doi:10.5194/nhess-18-3089-2018.
- Mezősi, G., et al., 2013: Assessment of regional climate change impacts on Hungarian landscapes. *Reg. Environ. Chang.*, **13**(4), 797–811, doi:10.1007/s10113-012-0326-1.
- Michailidou, A.V., C. Vlachokostas and N. Moussiopoulos, 2016: Interactions between climate change and the tourism sector: multiple-criteria decision analysis to assess mitigation and adaptation options in tourism areas. *Tour. Manag.*, **55**, 1–12, doi:10.1016/j.tourman.2016.01.010.
- Michetti, M. and M. Pinar, 2019: Forest fires across Italian regions and implications for climate change: a panel data analysis. *Environ. Resour. Econ.*, **72**(1), 207–246, doi:10.1007/s10640-018-0279-z.

- Mikhaylova, G., 2018: The Arctic society under the environmental and climate change (based on survey results). *Arct. North*, **32**, 95–106, doi:10.17238/issn2221-2698.2018.32.95.
- Miles, W.T.S., et al., 2017: Quantifying full phenological event distributions reveals simultaneous advances, temporal stability and delays in spring and autumn migration timing in long-distance migratory birds. *Glob. Change Biol.*, **23**(4), 1400–1414, doi:10.1111/gcb.13486.
- Mills, S.C., et al., 2017: European butterfly populations vary in sensitivity to weather across their geographical ranges. *Glob. Ecol. Biogeogr.*, **26**(12), 1374–1385, doi:10.1111/geb.12659.
- Mina, M., et al., 2017: Future ecosystem services from European mountain forests under climate change. *J. Appl. Ecol.*, **54**(2), 389–401, doi:10.1111/1365-2664.12772.
- Minicheva, G.G., et al., 2018: Assessment of the response of the Black Sea ecosystems to climate factors. *Algology*, **28**(2), 121–135.
- Mitchell, I., F. Daunt, M. Frederiksen and K. Wade, 2020: Impacts of climate change on seabirds, relevant to the coastal and marine environment around the UK. *MCIP Sci. Rev.*, **2020**, 382–399, doi:10.14465/2020.arc17.sbi.
- Mitter, H., et al., 2020: Shared socio-economic pathways for European agriculture and food systems: the Eur-Agri-SSPs. *Glob. Environ. Chang.*, **65**, 102159–102159, doi:10.1016/j.gloenvcha.2020.102159.
- Moemken, J., M. Meyers, H. Feldmann and J. Pinto, 2018: Future changes of wind speed and wind energy potentials in EURO-CORDEX ensemble simulations. *J. Geophys. Res.*, **123**(12), 6373–6389, doi:10.1029/2018JD028473.
- Monchamp, M.E., et al., 2018: Homogenization of lake cyanobacterial communities over a century of climate change and eutrophication. *Nat. Ecol. Evol.*, **2**(2), 317, doi:10.1038/s41559-017-0407-0.
- Montazeri, H., Y. Toparlar, B. Blocken and J.L.M. Hensen, 2017: Simulating the cooling effects of water spray systems in urban landscapes: a computational fluid dynamics study in Rotterdam, The Netherlands. *Landsc. Urban Plan.*, **159**, 85–100, doi:10.1016/j.landurbplan.2016.10.001.
- Montero Serra, I., M. Edwards and M.J. Genner, 2015: Warming shelf seas drive the subtropicalization of European pelagic fish communities. *Glob. Change Biol.*, **21**(1), 144–153, doi:10.1111/gcb.12747.
- Mora, C., et al., 2017: Global risk of deadly heat. *Nat. Clim. Chang.*, **7**, 501, doi:10.1038/nclimate3322.
- Morabito, M., et al., 2017: Increasing heatwave hazards in the southeastern European Union capitals. *Atmosphere*, **8**(7), doi:10.3390/atmos8070115.
- Morabito, M., et al., 2019: An occupational heat-health warning system for Europe: the HEAT-SHIELD platform. *Int J Environ Res Public Health*, **16**(16), doi:10.3390/ijerph16162890.
- Moraine, M., et al., 2014: Farming system design for innovative crop-livestock integration in Europe. *Animal*, **8**(8), 1204–1217, doi:10.1017/S1751731114001189.
- Morán-Ordóñez, A., et al., 2020: Future impact of climate extremes in the Mediterranean: soil erosion projections when fire and extreme rainfall meet. *Land Degrad. Dev.*, **31**(18), 3040–3054, doi:10.1002/ldr.3694.
- Moreno, M.V., M. Conedera, E. Chuvieco and G.B. Pezzatti, 2014: Fire regime changes and major driving forces in Spain from 1968 to 2010. *Environ. Sci. Policy*, **37**, 11–22, doi:10.1016/j.envsci.2013.08.005.
- Moretti, A., M. Pascale and A.F. Logrieco, 2019: Mycotoxin risks under a climate change scenario in Europe. *Trends Food Sci. Technol.*, **84**, 38–40, doi:10.1016/j.tifs.2018.03.008.
- Morgan, E.R. and J. van Dijk, 2012: Climate and the epidemiology of gastrointestinal nematode infections of sheep in Europe. *Vet. Parasitol.*, **189**(1), 8–14, doi:10.1016/j.vetpar.2012.03.028.
- Mori, E., A. Sforzi, G. Bogliani and P. Milanese, 2018: Range expansion and redefinition of a crop-raiding rodent associated with global warming and temperature increase. *Clim. Chang.*, **150**(3), 319–331, doi:10.1007/s10584-018-2261-8.
- Morignat, E., et al., 2014: Assessment of the impact of the 2003 and 2006 heat waves on cattle mortality in France. *PLoS ONE*, **9**(3), e93176–e93176.
- Moriondo, M., et al., 2006: Potential impact of climate change on fire risk in the Mediterranean area. *Clim. Res.*, **31**(1), 85–95, doi:10.3354/cr031085.
- Morote, Á.-F., J. Olcina and M. Hernández, 2019: The use of non-conventional water resources as a means of adaptation to drought and climate change in semi-arid regions: south-eastern Spain. *Water*, **11**(1), doi:10.3390/w11010093.
- Moullec, F., et al., 2019: An end-to-end model reveals losers and winners in a warming Mediterranean Sea. *Front. Mar. Sci.*, **6**, 1–19, doi:10.3389/fmars.2019.00345.
- Mourey, J., C. Perrin-Malterre and L. Ravanel, 2020: Strategies used by French Alpine guides to adapt to the effects of climate change. *J. Outdoor Recreat. Tour.*, **29**, 100278, doi:10.1016/j.jort.2020.100278.
- Muenzel, D. and S. Martino, 2018: Assessing the feasibility of carbon payments and payments for ecosystem services to reduce livestock grazing pressure on saltmarshes. *J. Environ. Manag.*, **225**, 46–61, doi:10.1016/j.jenvman.2018.07.060.
- Mullan, D., et al., 2019: Climate impacts on soil erosion and muddy flooding at 1.5 versus 2°C warming. *Land Degrad. Dev.*, **30**(1), 94–108, doi:10.1002/ldr.3214.
- Muller, J., D. Folini, M. Wild and S. Pfenniger, 2019: CMIP-5 models project photovoltaics are a no-regrets investment in Europe irrespective of climate change. *Energy*, **171**, 135–148, doi:10.1016/j.energy.2018.12.139.
- Mulville, M. and S. Stravaravdis, 2016: The impact of regulations on overheating risk in dwellings. *Build. Res. Inf.*, **44**(5-6), 520–534, doi:10.1080/09613218.2016.1153355.
- Munari, C., 2011: Effects of the 2003 European heatwave on the benthic community of a severe transitional ecosystem (Comacchio Saltworks, Italy). *Mar. Pollut. Bull.*, **62**(12), 2761–2770, doi:10.1016/j.marpolbul.2011.09.011.
- Murdock, C.C., E.D. Sternberg and M.B. Thomas, 2016: Malaria transmission potential could be reduced with current and future climate change. *Sci. Rep.*, **6**, doi:10.1038/srep27771.
- Murtagh, N., B. Gatersleben and C. Fife-Schaw, 2019: Occupants' motivation to protect residential building stock from climate-related overheating: a study in southern England. *J. Clean. Prod.*, **226**, 186–194, doi:10.1016/j.jclepro.2019.04.080.
- Mustonen, K., T. Mustonen, J. Kirillov and S. Council, 2018: *Traditional Knowledge of Northern Waters*. Snowchange Cooperative, Kontiolahti, Finland. 39 pp.
- Mustonen, T., 2014: Endemic time-spaces of Finland: aquatic regimes. *Fennia*, **192**(2), 120–139, doi:10.11143/40845.
- Mustonen, T., 2018: Meaningful engagement and oral histories of the indigenous peoples of the north. *Nord. Geogr. Publ.*, **47**(5), 21–38.
- Mustonen, T. and N. Huusari, 2020: How to know about waters? Finnish traditional knowledge related to waters and implications for management reforms. *Rev. Fish Biol. Fish.*, **30**, 699–718, doi:10.1007/s11160-020-09619-7.
- Mustonen, T., et al., 2021a: Community-based monitoring in the Ponoy River, Kola Peninsula (Russia): reflections on Atlantic salmon, pink salmon, Northern pike and weather/climate change. *Polar Biol.*, **44**(1), 173–194, doi:10.1007/s00300-020-02790-4.
- Mustonen, T., et al., 2021b: Ponoi river affected by pink salmon expansion and severe weather change. *Polar Biol.*, **44**, 173–194.
- Myers-Smith, I.H., et al., 2020: Complexity revealed in the greening of the Arctic. *Nat. Clim. Chang.*, **10**(2), 106–117, doi:10.1038/s41558-019-0688-1.
- Myers, B.J.E., et al., 2017: Global synthesis of the documented and projected effects of climate change on inland fishes. *Rev. Fish Biol. Fish.*, **27**(2), 339–361, doi:10.1007/s11160-017-9476-z.
- Nabuurs, G.-J., E.J.M.M. Arets and M.-J. Schelhaas, 2017: European forests show no carbon debt, only a long parity effect. *For. Policy Econ.*, **75**, 120–125, doi:10.1016/j.forpol.2016.10.009.
- Nabuurs, G.-J., M.-J. Schelhaas, G.M.J. Mohren and C.B. Field, 2003: Temporal evolution of the European forest sector carbon sink from 1950 to 1999. *Glob. Change Biol.*, **9**(2), 152–160, doi:10.1046/j.1365-2486.2003.00570.x.

- Nabuurs, G., 2018: Future scenarios of European forests. *J. Landsc. Ecol.*, **11**, 175–184, doi:10.2478/jlecol-2018-0020.
- Nah, K., A. Bede-Fazekas, A.J. Trajer and J.H. Wu, 2020: The potential impact of climate change on the transmission risk of tick-borne encephalitis in Hungary. *BMC Infect. Dis.*, **20**(1), doi:10.1186/s12879-019-4734-4.
- Narayan, S., et al., 2016: The effectiveness, costs and coastal protection benefits of natural and nature-based defences. *PLoS ONE*, **11**(5), e154735, doi:10.1371/journal.pone.0154735.
- Narita, D. and K. Rehdanz, 2017: Economic impact of ocean acidification on shellfish production in Europe. *J. Environ. Plan. Manag.*, **60**(3), 500–518, doi:10.1080/09640568.2016.1162705.
- Natali, S.M., et al., 2019: Large loss of CO₂ in winter observed across the northern permafrost region. *Nat. Clim. Chang.*, **9**(11), 852–857, doi:10.1038/s41558-019-0592-8.
- Naudts, K., et al., 2016: Europe's forest management did not mitigate climate warming. *Science*, **351**(6273), 597–600.
- Naueis, A., et al., 2019: Attributing long-term sea-level rise to Paris Agreement emission pledges. *Proc. Natl. Acad. Sci.*, **116**(47), 23487, doi:10.1073/pnas.1907461116.
- Naumann, G., et al., 2018: Global changes in drought conditions under different levels of warming. *Geophys. Res. Lett.*, **45**(7), 3285–3296, doi:10.1002/2017GL076521.
- Naumann, G., C. Cammalleri, L. Mentaschi and L. Feyen, 2021: Increased economic drought impacts in Europe with anthropogenic warming. *Nat. Clim. Chang.*, **11**(June), doi:10.1038/s41558-021-01044-3.
- Naumann, G., et al., 2020a: *Global warming and human impacts of heat and cold extremes in the EU*. Publications Office of the European Union, Luxembourg.
- Naumann, G., et al., 2020b: *Global warming and human impacts of heat and cold extremes in the EU*. Publications Office of the European Union, Luxembourg. JRC118540.
- Nellemann, C., P. Jordhøy, O.-G. Støen and O. Strand, 2000: Cumulative impacts of tourist resorts on wild reindeer (*Rangifer tarandus tarandus*) during winter. *Arctic*, **53**(1), 9–17, doi:10.14430/arctic829.
- Neset, T.-S., T. Asplund, J. Käyhkö and S. Juhola, 2019: Making sense of maladaptation: nordic agriculture stakeholders' perspectives. *Clim. Chang.*, **153**(4), 1–15.
- Newson, S.E., et al., 2016: Long-term changes in the migration phenology of UK breeding birds detected by large-scale citizen science recording schemes. *Ibis*, **158**(3), 481–495, doi:10.1111/ibi.12367.
- Nguyen, T.P.L., G. Seddaiu and P.P. Roggero, 2019: Declarative or procedural knowledge? Knowledge for enhancing farmers' mitigation and adaptation behaviour to climate change. *J. Rural Stud.*, **67**, 46–56, doi:10.1016/j.jrurstud.2019.02.005.
- Nielsen, A., T. Reitan, A.W. Rinvoll and A.K. Brysting, 2017: Effects of competition and climate on a crop pollinator community. *Agric. Ecosyst. Environ.*, **246**, 253–260, doi:10.1016/j.agee.2017.06.006.
- Nohe, A., et al., 2020: Marked changes in diatom and dinoflagellate biomass, composition and seasonality in the Belgian Part of the North Sea between the 1970s and 2000s. *Sci. Total. Environ.*, **716**, 136316–136316, doi:10.1016/j.scitotenv.2019.136316.
- Novick, K.A., et al., 2016: The increasing importance of atmospheric demand for ecosystem water and carbon fluxes. *Nat. Clim. Chang.*, **6**(11), 1023, doi:10.1038/nclimate3114.
- Nsoesie, E., et al., 2016: Global distribution and environmental suitability for chikungunya virus, 1952 to 2015. *Euro Surveill.*, **21**(20), 7–18, doi:10.2807/1560-7917.ES.2016.21.20.30234.
- O'Hare, P., I. White and A. Connelly, 2016: Insurance as maladaptation: resilience and the "business as usual" paradox. *Environ. Plan. C Gov. Policy*, **34**(6), 1175–1193, doi:10.1177/0263774X15602022.
- Oggioni, S.D., R. Ochoa-Hueso and B. Peco, 2020: Livestock grazing abandonment reduces soil microbial activity and carbon storage in a Mediterranean Dehesa. *Appl. Soil Ecol.*, **153**, 103588–103588, doi:10.1016/j.apsoil.2020.103588.
- Ojea, E., I. Pearlman, S.D. Gaines and S.E. Lester, 2017: Fisheries regulatory regimes and resilience to climate change. *Ambio*, **46**(4), 399–412, doi:10.1007/s13280-016-0850-1.
- Oliveira, S., J. Rocha, C.A. Sousa and C. Capinha, 2021: Wide and increasing suitability for *Aedes albopictus* in Europe is congruent across distribution models. *Sci. Rep.*, **11**(1), doi:10.1038/s41598-021-89096-5.
- Oliver, E.C.J., et al., 2018: Longer and more frequent marine heatwaves over the past century. *Nat. Commun.*, **9**(1), 12, doi:10.1038/s41467-018-03732-9.
- Oliver, T.H., et al., 2015: Interacting effects of climate change and habitat fragmentation on drought-sensitive butterflies. *Nat. Clim. Chang.*, **5**, 941, doi:10.1038/nclimate2746.
- Oliver, T.H., et al., 2014: Latitudinal gradients in butterfly population variability are influenced by landscape heterogeneity. *Ecography*, **37**(9), 863–871, doi:10.1111/ecog.00608.
- Öllerer, K., et al., 2019: Beyond the obvious impact of domestic livestock grazing on temperate forest vegetation – A global review. *Biol. Conserv.*, **237**, 209–219, doi:10.1016/j.biocon.2019.07.007.
- Olsen, L.S., 2016: Sami tourism in destination development: conflict and collaboration. *Polar Geogr.*, **39**(3), 179–195, doi:10.1080/1088937X.2016.1201870.
- Omazic, A., et al., 2019: Identifying climate-sensitive infectious diseases in animals and humans in Northern regions. *Acta Vet. Scand.*, **61**(1), 53, doi:10.1186/s13028-019-0490-0.
- Orehova, N.A., 2017: Components of the carbon cycle of the ecosystem of Sevastopol Bay (the Black Sea) according to 2017. *Ecol. Saf. Coast. Shelf Zones Sea*, **4**, 39–46.
- Orellana-Macias, J.M., et al., 2020: Shifts in crane migration phenology associated with climate change in Southwestern Europe. *Avian Conserv. Ecol.*, **15**(1), 1–13, doi:10.5751/ACE-01565-150116.
- Orlov, A., et al., 2019: Economic losses of heat-induced reductions in outdoor worker productivity: a case study of Europe. *EconDisCliCha*, **3**(3), 191–211, doi:10.1007/s41885-019-00044-0.
- Osberghaus, D., 2015: The determinants of private flood mitigation measures in Germany — Evidence from a nationwide survey. *Ecol. Econ.*, **110**, 36–50, doi:10.1016/j.ecolecon.2014.12.010.
- Osberghaus, D., 2017: The effect of flood experience on household mitigation—Evidence from longitudinal and insurance data. *Glob. Environ. Chang.*, **43**, 126–136, doi:10.1016/j.gloenvcha.2017.02.003.
- Össbo, Å., 2018: Recurring colonial ignorance: a genealogy of the Swedish energy system. *J. North. Stud.*, **12**(2), 63–80.
- Össbo, Å. and P. Lantto, 2011: Colonial tutelage and industrial colonialism: reindeer husbandry and early 20th-century hydroelectric development in Sweden. *Scand. J. Hist.*, **36**(3), 324–348, doi:10.1080/03468755.2011.580077.
- Österlin, C. and K. Raitio, 2020: Fragmented landscapes and planscapes—the double pressure of increasing natural resource exploitation on indigenous Sámi lands in Northern Sweden. *Resources*, **9**(9), 104, doi:10.3390/resources909104.
- Outhwaite, C., et al., 2020: Complex long-term biodiversity change among invertebrates, bryophytes and lichens. *Nat. Ecol. Evol.*, **4**(3), 384, doi:10.1038/s41559-020-1111-z.
- Ovaskainen, O., et al., 2013: Community-level phenological response to climate change. *Proc. Natl. Acad. Sci. U.S.A.*, **110**(33), 13434–13439, doi:10.1073/pnas.1305533110.
- Palkowski, C., S. von Schwarzenberg and A. Simo, 2019: Seasonal cooling performance of air conditioners: the importance of independent test procedures used for MEPS and labels. *Int. J. Refrig.*, **104**, 417–425, doi:10.1016/j.jrefrig.2019.05.021.
- Panagos, P., et al., 2017: Towards estimates of future rainfall erosivity in Europe based on REDES and WorldClim datasets. *J. Hydrol.*, **548**, 251–262, doi:10.1016/j.jhydrol.2017.03.006.

- Panagos, P., et al., 2015: The new assessment of soil loss by water erosion in Europe. *Environ. Sci. Policy*, **54**, 438–447, doi:10.1016/j.envsci.2015.08.012.
- Papadaskalopoulou, C., et al., 2015a: Challenges for water resources and their management in light of climate change: the case of Cyprus. *Desalin. Water Treat.*, **53**(12), 3224–3233, doi:10.1080/19443994.2014.933619.
- Papadaskalopoulou, C., et al., 2016: Review of the current EU framework on adaptation to climate change and assessment of the relative adaptation framework in Cyprus. *Desalin. Water Treat.*, **57**(5), 2219–2231, doi:10.1080/19443994.2015.1107179.
- Papadaskalopoulou, C., et al., 2015b: Review and assessment of the adaptive capacity of the water sector in Cyprus against climate change impacts on water availability. *Resour. Conserv. Recycl.*, **105**, 95–112, doi:10.1016/j.resconrec.2015.10.017.
- Papadimitriou, L., I.P. Holman, R. Dunford and P.A. Harrison, 2019: Trade-offs are unavoidable in multi-objective adaptation even in a post-Paris Agreement world. *Sci. Total Environ.*, **696**, 134027–134027, doi:10.1016/j.scitotenv.2019.134027.
- Pappenberger, F., et al., 2015: The monetary benefit of early flood warnings in Europe. *Environ. Sci. Policy*, **51**, 278–291, doi:10.1016/j.envsci.2015.04.016.
- Paprotny, D., A. Sebastian, O. Morales-Nápoles and S.N. Jonkman, 2018: Trends in flood losses in Europe over the past 150 years. *Nat. Commun.*, **9**(1), 1985, doi:10.1038/s41467-018-04253-1.
- Parent, B., et al., 2018: Maize yields over Europe may increase in spite of climate change, with an appropriate use of the genetic variability of flowering time. *Proc. Natl. Acad. Sci.*, **115**(42), 10642–10647, doi:10.1073/pnas.1720716115.
- Parkinson, R.W. and D.E. Ogurcak, 2018: Beach nourishment is not a sustainable strategy to mitigate climate change. *Estuar. Coast. Shelf Sci.*, **212**, 203–209, doi:10.1016/j.ecss.2018.07.011.
- Parmesan, C., et al., 1999: Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature*, **399**(6736), 579–583, doi:10.1038/21181.
- Parrado, R., et al., 2020: Fiscal effects and the potential implications on economic growth of sea-level rise impacts and coastal zone protection. *Clim. Chang.*, **160**(2), 283–302, doi:10.1007/s10584-020-02664-y.
- Pastor, A.V., et al., 2019a: Projecting future impacts of global change including fires on soil erosion to anticipate better land management in the forests of NW Portugal. *Water (Switzerland)*, **11**(12), doi:10.3390/w11122617.
- Pastor, A.V., et al., 2019b: Projecting future impacts of global change including fires on soil erosion to anticipate better land management in the forests of NW Portugal. *Water*, **11**(12), 2617, doi:10.3390/w11122617.
- Patro, E.R., C. De Michele and F. Avanzi, 2018: Future perspectives of run-of-the-river hydropower and the impact of glaciers' shrinkage: The case of Italian Alps. *Appl. Energy*, **231**, 699–713, doi:10.1016/j.apenergy.2018.09.063.
- Paudel, Y., W. Botzen and J. Aerts, 2015: Influence of climate change and socio-economic development on catastrophe insurance: a case study of flood risk scenarios in the Netherlands. *Reg. Environ. Chang.*, **15**(8), 1717–1729, doi:10.1007/s10113-014-0736-3.
- Pausas, J. and J.E. Keeley, 2019: Abrupt climate-independent fire regime changes. *Ecosystems*, **17**, 1109–1120, doi:10.1007/s10021-014-9773-5. SpringerLink.
- Pavlova, L.V., 2020: Main areas and results of the benthos research in the Barents Sea by the Murmansk Marine Biological Institute in 2015–2019. *Work. Kola Sci. Cent. Russ. Acad. Sci.*, **11**, 108–134, doi:10.37614/2307-5252.2020.11.4.005.
- Pavón-Jordán, D., et al., 2020: Positive impacts of important bird and biodiversity areas on wintering waterbirds under changing temperatures throughout Europe and North Africa. *Biol. Conserv.*, **246**, 108549, doi:10.1016/j.biocon.2020.108549.
- Payet-Burin, R., F. Bertoni, C. Davidsen and P. Bauer-Gottwein, 2018: Optimization of regional water - power systems under cooling constraints and climate change. *Energy*, **155**, 484–494, doi:10.1016/j.energy.2018.05.043.
- Paz, S., 2015: Climate change impacts on West Nile virus transmission in a global context. *Philos. Trans. R.Soc. B Biol. Sci.*, **370**(1665), 1–11, doi:10.1098/rstb.2013.0561.
- Pearce-Higgins, J.W., et al., 2017: A national-scale assessment of climate change impacts on species: assessing the balance of risks and opportunities for multiple taxa. *Biol. Conserv.*, **213**, 124–134, doi:10.1016/j.biocon.2017.06.035.
- Peaucelle, M., et al., 2019: Spatial variance of spring phenology in temperate deciduous forests is constrained by background climatic conditions. *Nat. Commun.*, **10**(1), 5388, doi:10.1038/s41467-019-13365-1.
- Pecl, G.T., et al., 2017: Biodiversity redistribution under climate change: impacts on ecosystems and human well-being. *Science*, **355**(6332), eaai9214, doi:10.1126/science.aai9214.
- Pecuchet, L., et al., 2020: Spatio-temporal dynamics of multi-trophic communities reveal ecosystem-wide functional reorganization. *Ecography*, **43**(2), 197–208, doi:10.1111/ecog.04643.
- Pedde, S., et al., 2021: Enriching the shared socioeconomic pathways to co-create consistent multi-sector scenarios for the UK. *Sci. Total Environ.*, **756**, 143172, doi:10.1016/j.scitotenv.2020.143172.
- Peltonen-Sainio, P., et al., 2016: Harmfulness of weather events and the adaptive capacity of farmers at high latitudes of Europe. *Clim. Res.*, **67**(3), 221–240.
- Penning-Rowsell, E.C. and S.J. Priest, 2015: Sharing the burden of increasing flood risk: who pays for flood insurance and flood risk management in the United Kingdom. *Mitig. Adapt. Strateg. Glob. Chang.*, **20**(6), 991–1009, doi:10.1007/s11027-014-9622-z.
- Pennino, M.G., et al., 2020: Current and future influence of environmental factors on small pelagic fish distributions in the northwestern Mediterranean sea. *Front. Mar. Sci.*, **7**, 622–622, doi:10.3389/fmars.2020.00622.
- Peñuelas, J., et al., 2017a: Shifting from a fertilization-dominated to a warming-dominated period. *Nat. Ecol. Evol.*, **1**(10), 1438–1445, doi:10.1038/s41559-017-0274-8.
- Peñuelas, J., et al., 2013: Evidence of current impact of climate change on life: a walk from genes to the biosphere. *Glob. Change Biol.*, **19**(8), 2303–2338, doi:10.1111/gcb.12143.
- Peñuelas, J., et al., 2017b: Impacts of global change on Mediterranean forests and their services. *Forests*, **8**(12), doi:10.3390/f8120463.
- Perch-Nielsen, S.L., B. Amelung and R. Knutti, 2010: Future climate resources for tourism in Europe based on the daily Tourism Climatic Index. *Clim. Chang.*, **103**(3), 363–381, doi:10.1007/s10584-009-9772-2.
- Pérez-Domínguez, I. and T. Fellmann, 2018: *PESETA III: Agro-economic analysis of climate change impacts in Europe*. Publications Office of the European Union, Luxembourg. JRC Technical Reports, EUR 29431 EN.
- Pérez-Morales, A., S. Gil-Guirado and J. Olcina-Cantos, 2018: Housing bubbles and the increase of flood exposure. Failures in flood risk management on the Spanish south-eastern coast (1975–2013). *J. Flood Risk Manag.*, **11**(S1), 302–S313, doi:10.1111/jfr3.12207.
- Pérez Navarro, M., et al., 2019: Climatic suitability derived from species distribution models captures community responses to an extreme drought episode. *Ecosystems*, **22**(1), 77–90, doi:10.1007/s10021-018-0254-0.
- Persson, S., D. Harnesk and M. Islar, 2017: What local people? Examining the Gállok mining conflict and the rights of the Sámi population in terms of justice and power. *Geoforum*, **86**, 20–29, doi:10.1016/j.geoforum.2017.08.009.
- Petrik, C.M., et al., 2019: Bottom-up drivers of global patterns of demersal, forage, and pelagic fishes. *Prog. Oceanogr.*, **176**, 102124, doi:10.1016/j.pocean.2019.102124.
- Petz, K., et al., 2016: *Indicators and modelling of land use, land management and ecosystem services. Methodological documentation*. PBL Netherlands Environmental Assessment Agency, The Hague. 109 pp.
- Piao, S., et al., 2019: Plant phenology and global climate change: current progresses and challenges. *Glob. Change Biol.*, **25**(6), 1922–1940, doi:10.1111/gcb.14619.

- Pietrapertosa, F., et al., 2019: Urban climate change mitigation and adaptation planning: Are Italian cities ready? *Cities*, **91**, 93–105, doi:10.1016/j.cities.2018.11.009.
- Pinkse, J. and F. Gasbarro, 2019: Managing physical impacts of climate change: an attentional perspective on corporate adaptation. *Bus. Soc.*, **58**(2), 333–368, doi:10.1177/0007650316648688.
- Piperaki, E. and G. Daikos, 2016: Malaria in Europe: emerging threat or minor nuisance? *Clin. Microbiol. Infect.*, **22**(6), 487–493, doi:10.1016/j.cmi.2016.04.023.
- Pirlone, F., I. Spadaro and S. Candia, 2020: More resilient cities to face higher risks. The case of genoa. *Sustainability*, **12**(12), 4825, doi:10.3390/su12124825.
- Plard, F., et al., 2014: Mismatch between birth date and vegetation phenology slows the demography of roe deer. *PLoS Biol.*, **12**(4), e1001828–e1001828, doi:10.1371/journal.pbio.1001828.
- Polce, C., et al., 2014: Climate-driven spatial mismatches between British orchards and their pollinators: increased risks of pollination deficits. *Glob. Change Biol.*, **20**(9), 2815–2828, doi:10.1111/gcb.12577.
- Polce, C., et al., 2016: Global change impacts on ecosystem services: a spatially explicit assessment for Europe. *One Ecosyst.*, **1**, e9990.
- Polemio, M. and T. Lonigro, 2015: Trends in climate, short-duration rainfall, and damaging hydrogeological events (Apulia, Southern Italy). *Nat. Hazards*, **75**(1), 515–540, doi:10.1007/s11069-014-1333-y.
- Polte, P., et al., 2021: Reduced reproductive success of Western Baltic herring (*Clupea harengus*) as a response to warming winters. *Front. Mar. Sci.*, **8**, 589242, doi:10.3389/fmars.2021.589242.
- Popp, A., et al., 2017: Land-use futures in the shared socio-economic pathways. *Glob. Environ. Chang.*, **42**, 331–345, doi:10.1016/j.gloenvcha.2016.10.002.
- Porfiriev, B., et al., 2017: Climate change impact on economic growth and specific sectors' development of the Russian Arctic. *Arct. Ecol. Econ.*, **4**(28), 13, doi:10.25283/2223-4594-2017-4-4-17.
- Posledovich, D., et al., 2018: Phenological synchrony between a butterfly and its host plants: experimental test of effects of spring temperature. *J. Anim. Ecol.*, **87**(1), 150–161, doi:10.1111/1365-2656.12770.
- Post, E., et al., 2019: The polar regions in a 2°C warmer world. *Sci. Adv.*, **5**(12), eaaw9883, doi:10.1126/sciadv.aaw9883.
- Potopová, V., et al., 2017: The impacts of key adverse weather events on the field-grown vegetable yield variability in the Czech Republic from 1961 to 2014. *Int. J. Climatol.*, **37**(3), 1648–1664, doi:10.1002/joc.4807.
- Poussin, J.K., W.J.W. Botzen and J.C.J.H. Aerts, 2013: Stimulating flood damage mitigation through insurance: an assessment of the French CatNat system. *Environ. Hazards*, **12**(3–4), 258–277, doi:10.1080/17477891.2013.832650.
- Poussin, J.K., W.J. Wouter Botzen and J.C.J.H. Aerts, 2015: Effectiveness of flood damage mitigation measures: empirical evidence from French flood disasters. *Glob. Environ. Chang.*, **31**, 74–84, doi:10.1016/j.gloenvcha.2014.12.007.
- Powney, G.D., et al., 2019: Widespread losses of pollinating insects in Britain. *Nat. Commun.*, **10**(1), 1018–1018, doi:10.1038/s41467-019-08974-9.
- Pranzini, E., L. Wetzel and A.T. Williams, 2015: Aspects of coastal erosion and protection in Europe. *J. Coast. Conserv.*, **19**(4), 445–459, doi:10.1007/s11852-015-0399-3.
- Pregnoiato, M., et al., 2017: Impact of climate change on disruption to urban transport networks from pluvial flooding. *J. Infrastruct. Syst.*, **23**(4), 4017015, doi:10.1061/(ASCE)IS.1943-555X.0000372.
- Prem, E.M., C. Reitschuler and P. Illmer, 2014: Livestock grazing on alpine soils causes changes in abiotic and biotic soil properties and thus in abundance and activity of microorganisms engaged in the methane cycle. *Eur. J. Soil Biol.*, **62**, 22–29, doi:10.1016/j.ejsobi.2014.02.014.
- Pretzsch, H., et al., 2014: Forest stand growth dynamics in Central Europe have accelerated since 1870. *Nat. Commun.*, **5**(1), doi:10.1038/ncomms5967.
- Prins, A.G., et al., 2017: *Perspectives on the Future of Nature in Europe: Impacts and Combinations*. PBL Netherlands Environmental Assessment Agency, The Hague, 117 pp.
- Prislan, P., et al., 2019: Growing season and radial growth predicted for *Fagus sylvatica* under climate change. *Clim. Chang.*, **153**(1), 181–197, doi:10.1007/s10584-019-02374-0.
- Pritchard, O.G., S.H. Hallett and T.S. Farewell, 2015: Probabilistic soil moisture projections to assess Great Britain's future clay-related subsidence hazard. *Clim. Chang.*, **133**(4), 635–650, doi:10.1007/s10584-015-1486-z.
- Proestos, Y., et al., 2015: Present and future projections of habitat suitability of the Asian tiger mosquito, a vector of viral pathogens, from global climate simulation. *Philos. Trans. R. Soc. B Biol. Sci.*, **370**(1665), doi:10.1098/rstb.2013.0554.
- Prokosheva, I.V., 2017: Dynamics of phenology process in mountain taiga zone of Vishera nature reserve (northern Urals) under the influence of climatic changes. *Probl. Ecol. Monit. Ecosyst. Model.*, **XXVIII**(2), 40–55, doi:10.21513/0207-2564-2017-2-40-55.
- Pushnya, M.V. and Z.A. Shirinyan, 2015: A new harmful pest of soyabean in Krasnodar Territory. *Zashchita Karantin Rast.*, **10**, 27–29.
- Pyatinsky, M.M., V.A. Shlyakhov and O.V. Shlyakhova, 2020: Dynamics of sprat reserves in the Black Sea and perspectives for its development. *Fish. Issues*, **21**(4), 937 032036, doi:10.1088/1755-1315/937/3/032036.
- Pycroft, J., J. Abrell and J.-C. Ciscar, 2016: The global impacts of extreme sea-level rise: a comprehensive economic assessment. *Environ. Resour. Econ.*, **64**(2), 225–253, doi:10.1007/s10640-014-9866-9.
- Qiu, C., et al., 2020: The role of northern peatlands in the global carbon cycle for the 21st century. *Glob. Ecol. Biogeogr.*, **29**(5), 956–973, doi:10.1111/geb.13081.
- Queirós, A.M., J. Fernandes, L. Genevier and C.P. Lynam, 2018: Climate change alters fish community size-structure, requiring adaptive policy targets. *Fish Fish.*, **19**(4), 613–621, doi:10.1111/faf.12278.
- Rabin, S.S., et al., 2020: Impacts of future agricultural change on ecosystem service indicators. *Earth Syst. Dyn.*, **11**(2), 357–376, doi:10.5194/esd-11-357-2020.
- Radenković, S., et al., 2017: Living on the edge: Forecasting the trends in abundance and distribution of the largest hoverfly genus (Diptera: Syrphidae) on the Balkan Peninsula under future climate change. *Biol. Conserv.*, **212**, 216–229, doi:10.1016/j.biocon.2017.06.026.
- Ragazzola, F., et al., 2016: Impact of high CO₂ on the geochemistry of the coralline algae *Lithothamnion glaciale*. *Sci. Rep.*, **6**, 20572, doi:10.1038/srep20572.
- Ranasinghe, R., A.C. Ruane, R. Vautard, N. Arnell, E. Coppola, F.A. Cruz, S. Dessai, A.S. Islam, M. Rahimi, D. Ruiz
- Carrascal, J. Sillmann, M.B. Sylla, C. Tebaldi, W. Wang, and R. Zaaboul, 2021: Climate Change Information for Regional Impact and for Risk Assessment. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press, Cambridge. In press.
- Ranzani, A., et al., 2018: Hydropower future: between climate change, renewable deployment, carbon and fuel prices. *Water*, **10**(9), doi:10.3390/w10091197.
- Rasmont, P., et al., 2015: Climatic risk and distribution atlas of European bumblebees. *BioRisk*, **10**, 1–236.
- Rasmus, S., S. Kivinen and M. Irannezhad, 2018: Basal ice formation in snow cover in Northern Finland between 1948 and 2016. *Environ. Res. Lett.*, **13**(11), 114009, doi:10.1088/1748-9326/aae541.
- Rasmussen, K., J. Thyrring, R. Muscarella and F. Borchsenius, 2017: Climate-change-induced range shifts of three allergenic ragweeds (*Ambrosia* L.) in Europe and their potential impact on human health. *PeerJ*, **2017**(3), e3104–e3104, doi:10.7717/peerj.3104.
- Ratcliffe, S., et al., 2017: Biodiversity and ecosystem functioning relations in European forests depend on environmental context. *Ecol. Lett.*, **20**(11), 1414–1426, doi:10.1111/ele.12849.

- Ravanel, L., F. Magnin and P. Deline, 2017: Impacts of the 2003 and 2015 summer heatwaves on permafrost-affected rock-walls in the Mont Blanc massif. *Sci. Total. Environ.*, **609**, 132–143, doi:10.1016/j.scitotenv.2017.07.055.
- Raybaud, V., M. Bacha, R. Amara and G. Beaugrand, 2017: Forecasting climate-driven changes in the geographical range of the European anchovy (*Engraulis encrasicolus*). *ICES J. Mar. Sci.*, **74**(5), 1288–1299, doi:10.1093/icesjms/fsx003.
- Reckien, D., J. Flacke, M. Olazabal and O. Heidrich, 2015: The influence of drivers and barriers on urban adaptation and mitigation plans-an empirical analysis of European cities. *PLoS ONE*, **10**(8), doi:10.1371/journal.pone.0135597.
- Reckien, D., et al., 2018: How are cities planning to respond to climate change? Assessment of local climate plans from 885 cities in the EU-28. *J. Clean. Prod.*, **191**, 207–219, doi:10.1016/j.jclepro.2018.03.220.
- Regos, A., et al., 2016: Predicting the future effectiveness of protected areas for bird conservation in Mediterranean ecosystems under climate change and novel fire regime scenarios. *Divers. Distrib.*, **22**(1), 83–96, doi:10.1111/ddi.12375.
- Reich, P.B., et al., 2016: Boreal and temperate trees show strong acclimation of respiration to warming. *Nature*, **531**(7596), 633–636, doi:10.1038/nature17142.
- Reichstein, M., et al., 2007: Reduction of ecosystem productivity and respiration during the European summer 2003 climate anomaly: a joint flux tower, remote sensing and modelling analysis. *Glob. Change Biol.*, **13**(3), 634–651, doi:10.1111/j.1365-2486.2006.01224.x.
- Reidsma, P., et al., 2015: Climate change impact and adaptation research requires integrated assessment and farming systems analysis: a case study in the Netherlands. *Environ. Res. Lett.*, **10**(4), 45004–45004, doi:10.1088/1748-9326/10/4/045004.
- Reimann, L., et al., 2018: Mediterranean UNESCO World Heritage at risk from coastal flooding and erosion due to sea-level rise. *Nat. Commun.*, **9**(1), 4161, doi:10.1038/s41467-018-06645-9.
- Reischl, C., R. Rauter and A. Posch, 2018: Urban vulnerability and adaptation to heatwaves: a case study of Graz (Austria). *Clim. Policy*, **18**(1), 63–75, doi:10.1080/14693062.2016.1227953.
- Restemeyer, B., M. van den Brink and J. Woltjer, 2018: Resilience unpacked – framing of ‘uncertainty’ and ‘adaptability’ in long-term flood risk management strategies for London and Rotterdam. *Eur. Plan. Stud.*, **26**(8), 1559–1579, doi:10.1080/09654313.2018.1490393.
- Reusch, T.B.H., et al., 2018: The Baltic Sea as a time machine for the future coastal ocean. *Sci. Adv.*, **4**(5), doi:10.1126/sciadv.aar8195.
- Revich, B.A., V.V. Maleev and M.D. Smirnova, 2019: *Climate Change and Public Health: Assessments, Indicators, Predictions* [Revich, B. A. and K. A. O. (eds.)]. INP RAS, Moscow.
- Rey-Valette, H., S. Robert and B. Rulleau, 2019: Resistance to relocation in flood-vulnerable coastal areas: a proposed composite index. *Clim. Policy*, **19**(2), 206–218, doi:10.1080/14693062.2018.1482823.
- Rey, D., I.P. Holman and J.W. Knox, 2017: Developing drought resilience in irrigated agriculture in the face of increasing water scarcity. *Reg. Environ. Chang.*, **17**(5), 1527–1540, doi:10.1007/s10113-017-1116-6.
- Reyer, C., et al., 2014: Projections of regional changes in forest net primary productivity for different tree species in Europe driven by climate change and carbon dioxide. *Ann. For. Sci.*, **71**(2), 211–225, doi:10.1007/s13595-013-0306-8.
- Reyers, M., J. Moemken and J. Pinto, 2016: Future changes of wind energy potentials over Europe in a large CMIP5 multi-model ensemble. *Int. J. Climatol.*, **36**(2), 783–796, doi:10.1002/joc.4382.
- Rezza, G., 2016: Dengue and other Aedes-borne viruses: a threat to Europe? *Euro Surveill.*, **21**(21), 2–4, doi:10.2807/1560-7917.es.2016.21.21.30238.
- Rial-Lovera, K., W.P. Davies and N.D. Cannon, 2017: Implications of climate change predictions for UK cropping and prospects for possible mitigation: a review of challenges and potential responses. *J. Sci. Food Agric.*, **97**(1), 17–32, doi:10.1002/jsfa.7767.
- Ribas, A., J. Olcina and D. Sauri, 2020: More exposed but also more vulnerable? Climate change, high intensity precipitation events and flooding in Mediterranean Spain. *Disaster Prev. Manag. Int. J.*, **29**(3), 229–248, doi:10.1108/DPM-05-2019-0149.
- Ricart, S., J. Olcina and A. Rico, 2019: Evaluating public attitudes and farmers’ beliefs towards climate change adaptation: awareness, perception, and populism at European level. *Land*, **8**(1), 4.
- Richon, C., et al., 2019: Biogeochemical response of the Mediterranean Sea to the transient SRES-A2 climate change scenario. *Biogeosciences*, **16**(1), 135–165, doi:10.5194/bg-16-135-2019.
- Richter, M., 2016: Urban climate change-related effects on extreme heat events in Rostock, Germany. *Urban Ecosyst.*, **19**(2), 849–866, doi:10.1007/s11252-015-0508-y.
- Riebesell, U., et al., 2018: Toxic algal bloom induced by ocean acidification disrupts the pelagic food web. *Nat. Sci. Data*, **8**(12), 1082–1086, doi:10.1038/s41558-018-0344-1.
- Rivetti, I., et al., 2014: Global warming and mass mortalities of benthic invertebrates in the Mediterranean Sea. *PLoS ONE*, **9**(12), doi:10.1371/journal.pone.0115655.
- Rizzetto, S., et al., 2016: Modelling the impact of climate change and atmospheric N deposition on French forests biodiversity. *Environ. Pollut.*, **213**, 1016–1027, doi:10.1016/j.envpol.2015.12.048.
- Rizzoli, A., et al., 2011: Lyme borreliosis in Europe. *Euro Surveill.*, **16**(27), 2–9.
- Roberts, A.M.I., C. Tansey, R.J. Smithers and A.B. Phillimore, 2015: Predicting a change in the order of spring phenology in temperate forests. *Glob. Change Biol.*, **21**(7), 2603–2611, doi:10.1111/gcb.12896.
- Roberts, H.C., et al., 2014: Response to an emerging vector-borne disease: surveillance and preparedness for Schmallenberg virus. *Prev. Vet. Med.*, **116**(4), 341–349, doi:10.1016/j.prevetmed.2014.08.020.
- Roberts, S.P.M., et al., 2011: Assessing continental-scale risks for generalist and specialist pollinating bee species under climate change. *BioRisk*, **6**(1), 1–18, doi:10.3897/biorisk.6.1325.
- Roces-Díaz, J., et al., 2021: Temporal changes in Mediterranean forest ecosystem services are driven by stand development, rather than by climate-related disturbances. *For. Ecol. Manag.*, **480**, 118623, doi:10.1016/j.foreco.2020.118623.
- Rodionov, V.Z., 2016: St. Petersburg Flood Protection Barrier: History and Environmental Problems. *Regional Ecology*, **4**(46) 13–26.
- Rodrigues, A.R., et al., 2020: Addressing soil protection concerns in forest ecosystem management under climate change. *For. Ecosyst.*, **7**(1), doi:10.1186/s40663-020-00247-y.
- Rodriguez-Alarcon, L.G.S., et al., 2021: Unprecedented increase of West Nile virus neuroinvasive disease, Spain, summer 2020. *Euro Surveill.*, **26**(19), doi:10.2807/1560-7917.es.2021.26.19.2002010.
- Roebeling, P.C., L. Costa, L. Magalhaes-Filho and V. Tekken, 2013: Ecosystem service value losses from coastal erosion in Europe: historical trends and future projections. *J. Coast. Conserv.*, **17**(3), 389–395, doi:10.1007/s11852-013-0235-6.
- Roggatz, C.C., et al., 2019: Saxitoxin and tetrodotoxin bioavailability increases in future oceans. *Nat. Clim. Chang.*, **9**(11), 840–844, doi:10.1038/s41558-019-0589-3.
- Rogovskyy, A.S., et al., 2020: Upsurge of Lyme borreliosis in Ukraine: a 20-year survey. *J. Travel Med.*, **27**(6), doi:10.1093/jtm/taaa100.
- Rohat, G., et al., 2019: Influence of changes in socioeconomic and climatic conditions on future heat-related health challenges in Europe. *Glob. Planet. Chang.*, **172**, 45–59, doi:10.1016/j.gloplacha.2018.09.013.
- Rojas-Downing, M.M., A.P. Nejadhashemi, T. Harrigan and S.A. Woznicki, 2017: Climate change and livestock: impacts, adaptation, and mitigation. *Clim. Risk Manag.*, **16**, 145–163, doi:10.1016/j.crm.2017.02.001.
- Rosbakh, S., et al., 2021: Siberian plants shift their phenology in response to climate change. *Glob. Change Biol.*, **27**(18), 4435–4448, doi:10.1111/gcb.15744.

- Rose Vineer, H., et al., 2020: GLOWORM-PARA: a flexible framework to simulate the population dynamics of the parasitic phase of gastrointestinal nematodes infecting grazing livestock. *Int. J. Parasitol.*, **50**(2), 133–144, doi:10.1016/j.ijpara.2019.11.005.
- Roson, R. and R. Damania, 2017: The macroeconomic impact of future water scarcity: An assessment of alternative scenarios. *J. Policy Model.*, **39**(6), 1141–1162, doi:10.1016/j.jpolmod.2017.10.003.
- Roson, R. and M. Sartori, 2016: Estimation of climate change damage functions for 140 regions in the GTAP 9 data base. *J. Glob. Econ. Anal.*, **1**(2), doi:10.21642/JGEA.010202AF.
- Rosqvist, N.I. and P. Eriksson, 2021: Impacts of climate warming on reindeer herding demand new land use strategies. *Ambio*, doi:10.1007/s13280-021-01655-2.
- Rotter, M., E. Hoffmann, A. Pechan and R. Stecker, 2016: Competing priorities: how actors and institutions influence adaptation of the German railway system. *Clim. Chang.*, **137**(3), 609–623, doi:10.1007/s10584-016-1702-5.
- Rotzer, T., et al., 2019: Process based simulation of tree growth and ecosystem services of urban trees under present and future climate conditions. *Sci. Total Environ.*, **676**, 651–664, doi:10.1016/j.scitotenv.2019.04.235.
- Routschek, A., J. Schmidt and F. Kreienkamp, 2014: Impact of climate change on soil erosion - A high-resolution projection on catchment scale until 2100 in Saxony/Germany. *CATENA*, **121**, 99–109, doi:10.1016/j.catena.2014.04.019.
- Royé, D., et al., 2020: Wildfire burnt area patterns and trends in Western Mediterranean Europe via the application of a concentration index. *Land Degrad. Dev.*, **31**(3), 311–324, doi:10.1002/ldr.3450.
- Rubio-Portillo, E., et al., 2016: Effects of the 2015 heat wave on benthic invertebrates in the Tabarca Marine Protected Area (southeast Spain). *Mar. Environ. Res.*, **122**, 135–142, doi:10.1016/j.marenvres.2016.10.004.
- Ruffault, J., V. Moron, R.M. Trigo and T. Curt, 2017: Daily synoptic conditions associated with large fire occurrence in Mediterranean France: evidence for a wind-driven fire regime. *Int. J. Climatol.*, **37**(1), 524–533, doi:10.1002/joc.4680.
- Ruiz-Benito, P., et al., 2014: Diversity increases carbon storage and tree productivity in Spanish forests. *Glob. Ecol. Biogeogr.*, **23**(3), 311–322, doi:10.1111/geb.12126.
- Ruiz-Ramos, M., et al., 2018: Adaptation response surfaces for managing wheat under perturbed climate and CO₂ in a Mediterranean environment. *Agric. Syst.*, **159**, 260–274, doi:10.1016/j.agsy.2017.01.009.
- Rumpf, S.B., et al., 2018: Range dynamics of mountain plants decrease with elevation. *Proc. Natl. Acad. Sci.*, **115**(8), 1848, doi:10.1073/pnas.1713936115.
- Ryan, S.J., C.J. Carlson, E.A. Mordecai and L.R. Johnson, 2019: Global expansion and redistribution of Aedes-borne virus transmission risk with climate change. *PLoS Negl. Trop. Dis.*, **13**(3), doi:10.1371/journal.pntd.0007213.
- Ryan, S.J., et al., 2021: Warming temperatures could expose more than 1.3 billion new people to Zika virus risk by 2050. *Glob. Change Biol.*, **27**(1), 84–93, doi:10.1111/gcb.15384.
- Sáenz-Romero, C., et al., 2017: Adaptive and plastic responses of *Quercus petraea* populations to climate across Europe. *Glob. Change Biol.*, **23**(7), 2831–2847, doi:10.1111/gcb.13576.
- Sairam, N., et al., 2019: Quantifying flood vulnerability reduction via private precaution. *Earth's Future*, **7**(3), 235–249, doi:10.1029/2018EF000994.
- Sajjadian, S.M., J. Lewis and S. Sharples, 2015: The potential of phase change materials to reduce domestic cooling energy loads for current and future UK climates. *Energy Build.*, **93**, 83–89, doi:10.1016/j.enbuild.2015.02.029.
- Sakhel, A., 2017: Corporate climate risk management: Are European companies prepared? *J. Clean. Prod.*, **165**, 103–118, doi:10.1016/j.jclepro.2017.07.056.
- Salinas-Rodríguez, C., et al., 2018: A semi risk-based approach for managing urban drainage systems under extreme rainfall. *Water*, **10**(4), 384, doi:10.3390/w10040384.
- Saltré, F., A. Duputié, C. Gaucherel and I. Chuine, 2015: How climate, migration ability and habitat fragmentation affect the projected future distribution of European beech. *Glob. Change Biol.*, **21**(2), 897–910, doi:10.1111/gcb.12771.
- Samaniego, L., et al., 2018: Anthropogenic warming exacerbates European soil moisture droughts. *Nat. Clim. Chang.*, **8**(5), 421–426, doi:10.1038/s41558-018-0138-5.
- San-Miguel-Ayanz, J., T. Durrant Houston, R. Boca, G. Libertá, A. Branco, D. De Rigo, D. Ferrari, P. Maiani, T. Artes Vivancos, H. Costa, F. Lana, P. Löffler, D. Nuijten, T. Leray and A. Ahlgren, 2018: *Forest Fires in Europe, Middle East and North Africa 2017 2018*. EUR 29318 EN, Publications Office of the European Union, Luxembourg, ISBN 978-9279928321.
- San-Miguel-Ayanz, J., T. Durrant, R. Boca and A. Camia, 2012: *Forest Fire Damage in Natura 2000 Sites 2000-2012*. EUR 25718 EN. Publications Office of the European Union, Luxembourg. JRC77834.
- Sánchez-García, D., C. Rubio-Bellido, M. Trisancho and M. Marrero, 2020: A comparative study on energy demand through the adaptive thermal comfort approach considering climate change in office buildings of Spain. *Build. Simul.*, **13**(1), 51–63, doi:10.1007/s12273-019-0560-2.
- Sanchez-Guevara, C., et al., 2019: Assessing population vulnerability towards summer energy poverty: case studies of Madrid and London. *Energy Build.*, **190**, 132–143, doi:10.1016/j.enbuild.2019.02.024.
- Sandø, A.B., et al., 2020: Climate change and new potential spawning sites for northeast Arctic cod. *Front. Mar. Sci.*, **7**, 28–28.
- Sandström, P., et al., 2016: On the decline of ground lichen forests in the Swedish boreal landscape: implications for reindeer husbandry and sustainable forest management. *Ambio*, **45**(4), 415–429, doi:10.1007/s13280-015-0759-0.
- Sanginés de Cárcer, P., et al., 2018: Vapor–pressure deficit and extreme climatic variables limit tree growth. *Glob. Change Biol.*, **24**(3), 1108–1122, doi:10.1111/gcb.13973.
- Santillán, D., L. Garrote, A. Iglesias and V. Sotes, 2020: Climate change risks and adaptation: new indicators for Mediterranean viticulture. *Mitig. Adapt. Strateg. Glob. Chang.*, **25**(5), 881–899, doi:10.1007/s11027-019-09899-w.
- Santos, A., et al., 2018: Artificial lakes as a climate change adaptation strategy in drylands: evaluating the trade-off on non-target ecosystem services. *Mitig. Adapt. Strateg. Glob. Chang.*, **23**(6), 887–906, doi:10.1007/s11027-017-9764-x.
- Saraiva, S., et al., 2019: Uncertainties in projections of the baltic sea ecosystem driven by an ensemble of global climate models. *Front. Earth Sci.*, **6**, 1, doi:10.3389/feart.2018.00244.
- Schaap, B.F., et al., 2013: Participatory design of farm level adaptation to climate risks in an arable region in The Netherlands. *Eur. J. Agron.*, **48**, 30–42, doi:10.1016/j.eja.2013.02.004.
- Schaffner, F. and A. Mathis, 2014: Dengue and dengue vectors in the WHO European region: past, present, and scenarios for the future. *Lancet Infect. Dis.*, **14**(12), 1271–1280, doi:10.1016/s1473-3099(14)70834-5.
- Schauberger, G., et al., 2020: Efficacy of adaptation measures to alleviate heat stress in confined livestock buildings in temperate climate zones. *Biosyst. Eng.*, **200**, 157–175, doi:10.1016/j.biosystemseng.2020.09.010.
- Scherrer, D. and C. Körner, 2011: Topographically controlled thermal-habitat differentiation buffers alpine plant diversity against climate warming. *J. Biogeogr.*, **38**(2), 406–416, doi:10.1111/j.1365-2699.2010.02407.x.
- Schiemann, F. and A. Sakhel, 2018: Carbon disclosure, contextual factors, and information asymmetry: the case of physical risk reporting. *Eur. Account. Rev.*, **28**(4), 791–818, doi:10.1080/09638180.2018.1534600.
- Schirpke, U., et al., 2017: Future impacts of changing land-use and climate on ecosystem services of mountain grassland and their resilience. *Ecosyst. Serv.*, **26**, 79–94, doi:10.1016/j.ecoser.2017.06.008.
- Schleussner, C.-F., et al., 2016: Differential climate impacts for policy-relevant limits to global warming: the case of 1.5°C and 2°C. *Earth Syst. Dynam.*, **7**(2), 327–351, doi:10.5194/esd-7-327-2016.
- Schöner, W., et al., 2019: Spatiotemporal patterns of snow depth within the Swiss-Austrian Alps for the past half century (1961 to 2012) and linkages to climate change. *Int. J. Climatol.*, **39**(3), 1589–1603, doi:10.1002/joc.5902.
- Schönhart, M., et al., 2014: Integrated analysis of climate change impacts and adaptation measures in Austrian agriculture. *Ger. J. Agric. Econ.*, **6**(3), 156–176, doi:10.22004/AG.ECON.253157.

- Schröder, W., G. Schmidt and S. Schönrock, 2014: Modelling and mapping of plant phenological stages as bio-meteorological indicators for climate change. *Environ. Sci. Eur.*, **26**(1), 5–5, doi:10.1186/2190-4715-26-5.
- Schröter, D., et al., 2005: Ecosystem service supply and vulnerability to global change in Europe. *Science*, **310**(5752), 1333, doi:10.1126/science.1115233.
- Schröter, M., et al., 2014: Ecosystem services and opportunity costs shift spatial priorities for conserving forest biodiversity. *PLoS ONE*, **9**(11), e112557, doi:10.1371/journal.pone.0112557.
- Schubert, S.D., et al., 2016: Global meteorological drought: a synthesis of current understanding with a focus on SST drivers of precipitation deficits. *J. Clim.*, **29**(11), 3989–4019, doi:10.1175/JCLI-D-15-0452.1.
- Schuerch, M., et al., 2018: Future response of global coastal wetlands to sea-level rise. *Nature*, **561**(7722), 231–234, doi:10.1038/s41586-018-0476-5.
- Schuldt, B., et al., 2020: A first assessment of the impact of the extreme 2018 summer drought on Central European forests. *Basic Appl. Ecol.*, **45**, 86–103, doi:10.1016/j.baae.2020.04.003.
- Schulze, E.D., et al., 2009: Importance of methane and nitrous oxide for Europe's terrestrial greenhouse-gas balance. *Nat. Geosci.*, **2**(12), 842–850, doi:10.1038/ngeo686.
- Schuster, C., J. Honold, S. Lauf and T. Lakes, 2017: Urban heat stress: novel survey suggests health and fitness as future avenue for research and adaptation strategies. *Environ. Res. Lett.*, **12**(4), 44021, doi:10.1088/1748-9326/aa5f35.
- Schwalm, C.R., et al., 2017: Global patterns of drought recovery. *Nature*, **548**, 202, doi:10.1038/nature23021.
- Scott, D., M. Rutty, B. Amelung and M. Tang, 2016: An inter-comparison of the Holiday Climate Index (HCI) and the Tourism Climate Index (TCI) in Europe. *Atmosphere*, **7**(6), 80.
- Scott, D., R. Steiger, H. Dannevig and C. Aall, 2019: Climate change and the future of the Norwegian alpine ski industry. *Curr. Issues Tour.*, **23**(19), 2396–2409, doi:10.1080/13683500.2019.1608919.
- Seebauer, S. and C. Winkler, 2020: Should I stay or should I go? Factors in household decisions for or against relocation from a flood risk area. *Glob. Environ. Chang.*, **60**, 102018, doi:10.1016/j.gloenvcha.2019.102018.
- Sehlin MacNeil, K., 2015: Shafted: a case of cultural and structural violence in the power relations between a Sami community and a mining company in northern Sweden. *Ethnol. Scand.*, **45**, 73–88.
- Seibold, S., et al., 2019: Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature*, **574**(7780), 671, doi:10.1038/s41586-019-1684-3.
- Seidl, R., M.-J. Schelhaas, W. Rammer and P.J. Verkerk, 2014: Increasing forest disturbances in Europe and their impact on carbon storage. *Nat. Clim. Chang.*, **4**(9), 806–810, doi:10.1038/nclimate2318.
- Semenza, J. and B. Menne, 2009: Climate change and infectious diseases in Europe. *Lancet Infect. Dis.*, **9**(6), 365–375, doi:10.1016/S1473-3099(09)70104-5.
- Semenza, J. and J. Suk, 2018: Vector-borne diseases and climate change: a European perspective. *FEMS Microbiol. Lett.*, **365**(2), doi:10.1093/femsle/fnx244.
- Semenza, J., et al., 2016: Climate change projections of West Nile virus infections in Europe: implications for blood safety practices. *Environ. Health*, **15**, doi:10.1186/s12940-016-0105-4.
- Semenza, J.C., et al., 2014: International dispersal of dengue through air travel: importation risk for Europe. *PLoS Negl Trop Dis*, **8**(12), doi:10.1371/journal.pntd.0003278.
- Semenza, J.C., et al., 2017: Environmental suitability of vibrio infections in a warming climate: an early warning system. *Environ. Health Perspect.*, **125**(10), 107004.
- Senapati, N., P. Stratonovitch, M.J. Paul and M.A. Semenov, 2019: Drought tolerance during reproductive development is important for increasing wheat yield potential under climate change in Europe. *J. Exp. Bot.*, **70**(9), 2549–2560, doi:10.1093/jxb/ery226.
- Seneviratne, S.I., X. Zhang, M. Adnan, W. Badi, C. Dereczynski, A. Di Luca, S. Ghosh, I. Iskandar, J. Kossin, S. Lewis, F. Otto, I. Pinto, M. Satoh, S.M. Vicente-Serrano, M. Wehner, and B. Zhou, 2021: Weather and Climate Extreme Events in a Changing Climate. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press, Cambridge. In press.
- Serpa, D., et al., 2015: Impacts of climate and land use changes on the hydrological and erosion processes of two contrasting Mediterranean catchments. *Sci. Total Environ.*, **538**, 64–77, doi:10.1016/j.scitotenv.2015.08.033.
- Serra, I.M., M. Edwards and M.J. Genner, 2015: Warming shelf seas drive the subtropicalization of European pelagic fish communities. *Glob. Change Biol.*, **21**(1), 144–153, doi:10.1111/gcb.12747.
- Settele, J., J. Bishop and S.G. Potts, 2016: Climate change impacts on pollination. *Nat. Plants*, **2**(7), 16092–16092, doi:10.1038/nplants.2016.92.
- Sguotti, C., et al., 2019a: Non-linearity in stock–recruitment relationships of Atlantic cod: insights from a multi-model approach. *ICES J. Mar. Sci.*, **77**(4), 1492–1502, doi:10.1093/icesjms/fsz113.
- Sguotti, C., et al., 2019b: Catastrophic dynamics limit Atlantic cod recovery. *Proc. R. Soc. B.*, **286**(1898), 20182877, doi:10.1098/rspb.2018.2877.
- Shaposhnikov, D., et al., 2015: Long-term impact of moscow heat wave and wildfires on mortality. *Epidemiology*, **26**(2), E21–E22, doi:10.1097/EDE.0000000000000251.
- Shen, J., et al., 2020: An early-stage analysis of climate-adaptive designs for multi-family buildings under future climate scenario: case studies in Rome, Italy and Stockholm, Sweden. *J. Build. Eng.*, **27**, 100972, doi:10.1016/j.jobe.2019.100972.
- Shepard, D.S., E.A. Undurraga, Y.A. Halasa and J.D. Stanaway, 2016: The global economic burden of dengue: a systematic analysis. *Lancet Infect. Dis.*, **16**(8), 935–941, doi:10.1016/s1473-3099(16)00146-8.
- Sheverdyaev, I.V., 2019: Integrated mathematical model of the Barents and White Seas large marine ecosystems - a tool for assessing natural risks and efficient use of biological resources. *Rep. Acad. Sci.*, **487**(5), 566–572.
- Sieber, J., 2013: Impacts of, and adaptation options to, extreme weather events and climate change concerning thermal power plants. *Clim. Chang.*, **121**(1), 55–66, doi:10.1007/s10584-013-0915-0.
- Siebert, S., H. Webber, G. Zhao and F. Ewert, 2017: Heat stress is overestimated in climate impact studies for irrigated agriculture. *Environ. Res. Lett.*, **12**(5), 54023, doi:10.1088/1748-9326/aa702f.
- Singh, C., et al., 2020: Assessing the feasibility of adaptation options: methodological advancements and directions for climate adaptation research and practice. *Clim. Chang.*, **162**(2), 255–277, doi:10.1007/s10584-020-02762-x.
- Sirois-Delisle, C. and J.T. Kerr, 2018: Climate change-driven range losses among bumblebee species are poised to accelerate. *Sci. Rep.*, **8**(1), 14464, doi:10.1038/s41598-018-32665-y.
- Sitnov, S.A. and I.I. Mokhov, 2018: A comparative analysis of the characteristics of active fires in the boreal forests of Eurasia and north america based on satellite data. *Izv. Atmos. Ocean. Phys.*, **54**(9), 966–978, doi:10.1134/S0001433818090347.
- Sitnov, S.A., I.I. Mokhov and A.V. Jola, 2017: Influence of Siberian fires on carbon monoxide content in the atmosphere over the European part of Russia in the summer of 2016. *Opt. Atmos. Ocean.*, **30**(2), 146–152.
- Skarin, A. and B. Åhman, 2014: Do human activity and infrastructure disturb domesticated reindeer? The need for the reindeer's perspective. *Polar Biol.*, **37**(7), 1041–1054, doi:10.1007/s00300-014-1499-5.
- Skarin, A. and M. Alam, 2017: Reindeer habitat use in relation to two small wind farms, during preconstruction, construction, and operation. *Ecol. Evol.*, **7**(11), 3870–3882, doi:10.1002/ece3.2941.
- Skarin, A., Ö. Danell, R. Bergström and J. Moen, 2010: Reindeer movement patterns in alpine summer ranges. *Polar Biol.*, **33**(9), 1263–1275, doi:10.1007/s00300-010-0815-y.

- Skarin, A., et al., 2015: Wind farm construction impacts reindeer migration and movement corridors. *Landsc. Ecol.*, **30**(8), 1527–1540, doi:10.1007/s10980-015-0210-8.
- Skougaard Kaspersen, P., et al., 2017: Comparison of the impacts of urban development and climate change on exposing European cities to pluvial flooding. *Hydrol. Earth Syst. Sci.*, **21**(8), 4131–4147, doi:10.5194/hess-21-4131-2017.
- Skuce, P.J., E.R. Morgan, J. van Dijk and M. Mitchell, 2013: Animal health aspects of adaptation to climate change: beating the heat and parasites in a warming Europe. *Animal*, **7**, 333–345, doi:10.1017/S175173111300075X.
- Smale, D.A., et al., 2019a: Marine heatwaves threaten global biodiversity and the provision of ecosystem services. *Nat. Sci. Data*, **9**, 1, doi:10.1038/s41558-019-0412-1.
- Smale, D.A., et al., 2019b: Marine heatwaves threaten global biodiversity and the provision of ecosystem services. *Nat. Clim. Chang.*, **9**(4), 306–312, doi:10.1038/s41558-019-0412-1.
- Smale, D.A., A.L.E. Yunnice, T. Vance and S. Widdicombe, 2015: Disentangling the impacts of heat wave magnitude, duration and timing on the structure and diversity of sessile marine assemblages. *PeerJ*, **3**(1628), doi:10.7717/peerj.863.
- Smith, J.O., et al., 2005: Projected changes in mineral soil carbon of European croplands and grasslands, 1990–2080. *Glob. Chang. Biol.*, **11**(12), 2141–2152, doi:10.1111/j.1365-2486.2005.001075.x.
- Smoliński, S. and Z. Mirny, 2017: Otolith biochronology as an indicator of marine fish responses to hydroclimatic conditions and ecosystem regime shifts. *Ecol. Indic.*, **79**, 286–294, doi:10.1016/j.ecolind.2017.04.028.
- Sneessens, I., L. Sauvé, H. Randrianasolo-Rakotobe and S. Ingrand, 2019: A framework to assess the economic vulnerability of farming systems: application to mixed crop-livestock systems. *Agric. Syst.*, **176**, 102658–102658, doi:10.1016/j.agsy.2019.102658.
- Sobol, N.V., I.M. Gabbasova and M.A. Komissarov, 2015: Impact of climate changes on erosion processes in Republic of Bashkortostan. *Arid Ecosyst.*, **5**(4), 216–221, doi:10.1134/S2079096115040137.
- Solaun, K. and E. Cerdá, 2017: The impact of climate change on the generation of hydroelectric power—A case study in southern Spain. *Energies*, **10**(9), doi:10.3390/en10091343.
- Solaun, K. and E. Cerdá, 2020: Impacts of climate change on wind energy power – Four wind farms in Spain. *Renew. Energy*, **145**, 1306–1316, doi:10.1016/j.renene.2019.06.129.
- Solimini, A.G., et al., 2018: Estimating the risk of Dengue, Chikungunya and Zika outbreaks in a large European city. *Sci. Rep.*, **8**, doi:10.1038/s41598-018-34664-5.
- Soroye, P., T. Newbold and J. Kerr, 2020: Climate change contributes to widespread declines among bumble bees across continents. *Science*, **367**(6478), 685, doi:10.1126/science.aax8591.
- Spandre, P., H. François, E. George-Marcelpoil and S. Morin, 2016: Panel based assessment of snow management operations in French ski resorts. *J. Outdoor Recreat. Tour.*, **16**, 24–36, doi:10.1016/j.jort.2016.09.002.
- Spencer, T., M. Schuerch, R.J. Nicholls, et al., 2016: Global coastal wetland change under sea-level rise and related stresses: the DIVA wetland change model. *Glob. Planet. Chang.*, **139**, 15–30, doi:10.1016/j.gloplacha.2015.12.018.
- Spinoni, J., et al., 2019: A new global database of meteorological drought events from 1951 to 2016. *J. Hydrol. Reg. Stud.*, **22**, 100593, doi:10.1016/j.ejrh.2019.100593.
- Spinoni, J., J. Vogt and P. Barbosa, 2015: European degree-day climatologies and trends for the period 1951–2011. *Int. J. Climatol.*, **35**(1), 25–36, doi:10.1002/joc.3959.
- Spivak, A.C., et al., 2019: Global-change controls on soil-carbon accumulation and loss in coastal vegetated ecosystems. *Nat. Geosci.*, **12**(9), 685–692, doi:10.1038/s41561-019-0435-2.
- Spooner, F.E.B., R.G. Pearson and R. Freeman, 2018: Rapid warming is associated with population decline among terrestrial birds and mammals globally. *Glob. Change Biol.*, **24**(10), 4521–4531, doi:10.1111/gcb.14361.
- Sswat, M., et al., 2018a: Growth performance and survival of larval Atlantic herring, under the combined effects of elevated temperatures and CO₂. *PLoS ONE*, **13**(1), e191947, doi:10.1371/journal.pone.0191947.
- Sswat, M., et al., 2018b: Food web changes under ocean acidification promote herring larvae survival. *Nat. Ecol. Evol.*, **2**(5), 836–840, doi:10.1038/s41559-018-0514-6.
- Stagge, J.H., D.G. Kingston, L.M. Tallaksen and D.M. Hannah, 2017: Observed drought indices show increasing divergence across Europe. *Sci. Rep.*, **7**(1), 14045, doi:10.1038/s41598-017-14283-2.
- Stahl, K., et al., 2016: Impacts of European drought events: insights from an international database of text-based reports. *Nat. Hazards Earth Syst. Sci.*, **16**(3), 801–819, doi:10.5194/nhess-16-801-2016.
- Stamos, I., E. Mitsakis and J. Grau, 2015: Roadmaps for adaptation measures of transportation to climate change. *Transp. Res. Rec.*, **2532**, 1–12, doi:10.3141/2532-01.
- Stanaway, J.D., et al., 2016: The global burden of dengue: an analysis from the Global Burden of Disease Study 2013. *Lancet Infect. Dis.*, **16**(6), 712–723, doi:10.1016/s1473-3099(16)00026-8.
- Stańczuk-Gałowicz, M., K. Sobolewska-Mikulska, H. Ritzema and J.M. van Loon-Steensma, 2018: Integration of water management and land consolidation in rural areas to adapt to climate change: experiences from Poland and the Netherlands. *Land Use Policy*, **77**, 498–511, doi:10.1016/j.landusepol.2018.06.005.
- Staudt, F., et al., 2021: The sustainability of beach nourishments: a review of nourishment and environmental monitoring practice. *J. Coast. Conserv.*, **25**(2), 34, doi:10.1007/s11852-021-00801-y.
- Stecf, 2019: *Monitoring the Performance of the Common Fisheries Policy*. Publications Office of the European Union, Luxembourg.
- Steele, D.J., et al., 2019: *Management and Drivers of Change of Pollinating Insects and Pollination Services*. The Department for Environment, Food and Rural Affairs, UK. January.
- Stefanescu, C., J. Carnicer and J. Peñuelas, 2011: Determinants of species richness in generalist and specialist Mediterranean butterflies: the negative synergistic forces of climate and habitat change. *Ecography*, **34**(3), 353–363, doi:10.1111/j.1600-0587.2010.06264.x.
- Steiger, R., E. Posch, G. Tappeiner and J. Walde, 2020: The impact of climate change on demand of ski tourism – a simulation study based on stated preferences. *Ecol. Econ.*, **170**, 106589, doi:10.1016/j.ecolecon.2019.106589.
- Steiger, R. and D. Scott, 2020: Ski tourism in a warmer world: increased adaptation and regional economic impacts in Austria. *Tour. Manag.*, **77**, 104032, doi:10.1016/j.tourman.2019.104032.
- Steiger, R. and J. Stötter, 2013: Climate change impact assessment of ski tourism in Tyrol. *Tour. Geogr.*, **15**(4), 577–600, doi:10.1080/14616688.2012.762539.
- Steinbauer, M.J., et al., 2018: Accelerated increase in plant species richness on mountain summits is linked to warming. *Nature*, **556**(7700), 231–234, doi:10.1038/s41586-018-0005-6.
- Stepanyan, O.V., 2020: Macrofitobentos in the large ecosystems of the southern seas of Russia. *News Russ. Acad. Sci. Geogr. Ser.*, **84**(2), 228–238.
- Stephens, P.A., et al., 2016: Consistent response of bird populations to climate change on two continents. *Science*, **352**(6281), 84, doi:10.1126/science.aac4858.
- Stiasny, M.H., et al., 2018: Effects of parental acclimation and energy limitation in response to high CO₂ exposure in Atlantic cod. *Sci. Rep.*, **8**(1), 8348, doi:10.1038/s41598-018-26711-y.
- Stiasny, M.H., et al., 2016: Ocean acidification effects on Atlantic cod larval survival and recruitment to the fished population. *PLoS ONE*, **11**(8), e155448, doi:10.1371/journal.pone.0155448.
- Stiasny, M.H., et al., 2019: Divergent responses of Atlantic cod to ocean acidification and food limitation. *Glob. Change Biol.*, **25**(3), 839–849, doi:10.1111/gcb.14554.
- Stocker, B.D., et al., 2018: Quantifying soil moisture impacts on light use efficiency across biomes. *New Phytol.*, **218**(4), 1430–1449, doi:10.1111/nph.15123.

- Stocker, B.D., et al., 2019: Drought impacts on terrestrial primary production underestimated by satellite monitoring. *Nat. Geosci.*, **12**(4), 264, doi:10.1038/s41561-019-0318-6.
- Stoffel, M., D. Tiranti and C. Huggel, 2014: Climate change impacts on mass movements—case studies from the European Alps. *Sci. Total Environ.*, **493**, 1255–1266, doi:10.1016/j.scitotenv.2014.02.102.
- Stojanov, R., et al., 2015: Adaptation to the impacts of climate extremes in Central Europe: a case study in a rural area in the Czech Republic. *Sustainability*, **7**(9), doi:10.3390/su70912758.
- Stoor, P., 2016: *Kunskapssammanställning om samers psykosociala ohälsa*. Sametinget, Giron/Kiruna. 144 pp.
- Straatsma, M.W., et al., 2019: Towards multi-objective optimization of large-scale fluvial landscaping measures. *Nat. Hazards Earth Syst. Sci.*, **19**(6), 1167–1187, doi:10.5194/nhess-19-1167-2019.
- Strasser, U., et al., 2019: Storylines of combined future land use and climate scenarios and their hydrological impacts in an Alpine catchment (Brixental/Austria). *Sci. Total Environ.*, **657**, 746–763, doi:10.1016/j.scitotenv.2018.12.077.
- Stucchi, L., G.M. Bombelli, A. Bianchi and D. Bocchiola, 2019: Hydropower from the alpine cryosphere in the era of climate change: the case of the Sabbione storage plant in Italy. *Water*, **11**(8), doi:10.3390/w11081599.
- Suggitt, A.J., et al., 2018: Extinction risk from climate change is reduced by microclimatic buffering. *Nat. Clim. Chang.*, **8**(8), 713–717, doi:10.1038/s41558-018-0231-9.
- Surminski, S., 2018: Fit for purpose and fit for the future? An evaluation of the UK's new flood reinsurance pool. *Risk Manag. Insur. Rev.*, **21**(1), 33–72, doi:10.1111/rmir.12093.
- Surminski, S., et al., 2015: Reflections on the current debate on how to link flood insurance and disaster risk reduction in the European Union. *Nat. Hazards*, **79**(3), 1451–1479, doi:10.1007/s11069-015-1832-5.
- Surminski, S., et al., 2018: Assessing climate risks across different business sectors and industries: an investigation of methodological challenges at national scale for the UK. *Phil. Trans. R.Soc. A.*, **376**, 20170307, doi:10.1098/rsta.2017.0307.
- Surminski, S. and A.H. Thieken, 2017: Promoting flood risk reduction: the role of insurance in Germany and England. *Earths Future*, **5**(10), 979–1001, doi:10.1002/2017EF000587.
- Sutton, W.R., J.P. Srivastava and J.E. Neumann, 2013: *Looking beyond the horizon: How Climate Change Impacts and Adaptation Responses Will Reshape Agriculture in Eastern Europe and Central Asia*. Directions in Development, The World Bank, Washington, DC, ISBN 978-0821397688. 201 pp.
- Suykens, C., et al., 2016: Dealing with flood damages: will prevention, mitigation, and ex post compensation provide for a resilient triangle? *Ecol. Soc.*, **21**(4), doi:10.5751/ES-08592-210401.
- Svetlitchnyi, O.A., 2020: Long-term forecast of changes in soil erosion losses during spring snowmelt caused by climate within the plain part of Ukraine. *J. Geol. Geogr. Geocol.*, **29**(3), 591–605, doi:10.15421/112054.
- Sámi Parliament in Sweden, 2021: *Boundaries of reindeer herding areas in Sweden*. Sami Parliament in Sweden, Girun. <https://www.sametinget.se/8382>. Accessed 2021.
- Sykes, R.A. and P. Makiello, 2017: An estimate of Lyme borreliosis incidence in Western Europe. *J. Public Health*, **39**(1), 74–81, doi:10.1093/pubmed/fdw017.
- Szabó, B., E. Vincze and B. Czúcz, 2016: Flowering phenological changes in relation to climate change in Hungary. *Int. J. Biometeorol.*, **60**(9), 1347–1356, doi:10.1007/s00484-015-1128-1.
- Szewczyk, W., J.C. Ciscar, I. Mongelli and A. Soria, 2018: *JRC PESETA III Project: Economic Integration and Spillover Analysis*, EUR 29456 EN. Publications Office of the European Union, Luxembourg, ISBN 978-9279974229.
- Szewczyk, W., et al., 2020: *Economic Analysis of Selected Climate Impacts: JRC PESETA IV project: Task 14*. Publications Office of the European Union, Luxembourg, ISBN 978-9276184591.
- Szumelda, A.U., 2019: Agriculture and everyday realities on small farms – An entrepreneurial challenge to farmers between the desire for autonomy and a secure existence. Two examples from east and south-east Poland. *J. Rural Stud.*, **67**, 57–68, doi:10.1016/j.jrurstud.2019.02.008.
- Takakura, J.Y., et al., 2017: Cost of preventing workplace heat-related illness through worker breaks and the benefit of climate-change mitigation. *Environ. Res. Lett.*, **12**(6), 64010, doi:10.1088/1748-9326/aa72cc.
- Talavera, G., X. Espadaler and R. Vila, 2015: Discovered just before extinction? The first endemic ant from the Balearic Islands (*Lasius balearicus* sp. nov.) is endangered by climate change. *J. Biogeogr.*, **42**(3), 589–601, doi:10.1111/jbi.12438.
- Tanner, S.E., et al., 2020: Marine regime shifts impact synchrony of deep-sea fish growth in the northeast Atlantic. *Oikos*, **129**(12), 1781–1794, doi:10.1111/oik.07332.
- Tanner, S.E., et al., 2019: Regional climate, primary productivity and fish biomass drive growth variation and population resilience in a small pelagic fish. *Ecol. Indic.*, **103**, 530–541, doi:10.1016/j.ecolind.2019.04.056.
- Taylor, J., et al., 2018: Comparison of built environment adaptations to heat exposure and mortality during hot weather, West Midlands region, UK. *Environ. Int.*, **111**, 287–294, doi:10.1016/j.envint.2017.11.005.
- TCFD, 2017: *Implementing the Recommendations of the Task Force on Climate Related Financial Disclosures*. <https://www.fsb-tcfd.org/wp-content/uploads/2017/12/FINAL-TCFD-Annex-Amended-121517.pdf>. Accessed 2019.
- Tedim, F., G. Xanthopoulos and V. Leone, 2015: Forest fires in Europe. In: *Wildfire Hazards, Risks and Disasters*. Elsevier, Amsterdam, pp. 77–99. ISBN 978-0124104341.
- Teixeira, C.M., et al., 2016: Environmental influence on commercial fishery landings of small pelagic fish in Portugal. *Reg. Environ. Chang.*, **16**(3), 709–716, doi:10.1007/s10113-015-0786-1.
- Temmerman, S., et al., 2013: Ecosystem-based coastal defence in the face of global change. *Nature*, **504**(7478), 79–83, doi:10.1038/nature12859.
- Teotónio, C., M. Rodríguez, P. Roebeling and P. Fortes, 2020: Water competition through the 'water-energy' nexus: assessing the economic impacts of climate change in a Mediterranean context. *Energy Econ.*, **85**, 104539, doi:10.1016/j.eneco.2019.104539.
- Termaat, T., et al., 2019: Distribution trends of European dragonflies under climate change. *Divers. Distrib.*, **25**(6), 936–950, doi:10.1111/ddi.12913.
- Teuling, A.J., et al., 2017: Observational evidence for cloud cover enhancement over western European forests. *Nat. Commun.*, **8**, 14065, doi:10.1038/ncomms14065. <https://www.nature.com/articles/ncomms14065#supplementary-information>.
- Thacker, S., S. Kelly, R. Pant and J.W. Hall, 2018: Evaluating the benefits of adaptation of critical infrastructures to hydrometeorological risks. *Risk Anal.*, **38**(1), 134–150, doi:10.1111/risa.12839.
- Thackeray, S.J., et al., 2016a: Phenological sensitivity to climate across taxa and trophic levels. *Nature*, **535**(7611), 241–245, doi:10.1038/nature18608.
- Thackeray, S.J., et al., 2016b: Phenological sensitivity to climate across taxa and trophic levels. *Nature*, **535**(7611), 241–294, doi:10.1038/nature18608.
- Thackeray, S.J., et al., 2013: Quantifying uncertainties in biologically-based water quality assessment: a pan-European analysis of lake phytoplankton community metrics. *Ecol. Indic.*, **29**, 34–47, doi:10.1016/j.ecolind.2012.12.010.
- Thaler, T., 2021: Just retreat—how different countries deal with it: examples from Austria and England. *J. Environ. Stud. Sci.*, **3**(14), 412–419, doi:10.1007/s13412-021-00694-1.
- Thieken, A.H., et al., 2016: Estimating changes in flood risks and benefits of non-structural adaptation strategies - a case study from Tyrol, Austria. *Mitig. Adapt. Strateg. Glob. Chang.*, **21**(3), 343–376, doi:10.1007/s11027-014-9602-3.
- Thomsen, J., et al., 2017: Naturally acidified habitat selects for ocean acidification tolerant mussels. *Sci. Adv.*, **3**(4), e1602411, doi:10.1126/sciadv.1602411.

- Thomson, H., S. Bouzarovski and C. Snell, 2017: Rethinking the measurement of energy poverty in Europe: A critical analysis of indicators and data. *Indoor Built Environ.*, **26**(7), 879–901, doi:10.1177/1420326X17699260.
- Thomson, H., N. Simcock, S. Bouzarovski and S. Petrova, 2019: Energy poverty and indoor cooling: an overlooked issue in Europe. *Energy Build.*, **196**, 21–29, doi:10.1016/j.enbuild.2019.05.014.
- Tian, H., et al., 2016: The terrestrial biosphere as a net source of greenhouse gases to the atmosphere. *Nature*, **531**(7593), 225–228, doi:10.1038/nature16946.
- Tillson, A.-A., T. Oreszczyn and J. Palmer, 2013: Assessing impacts of summertime overheating: some adaptation strategies. *Build. Res. Inf.*, **41**(6), 652–661, doi:10.1080/09613218.2013.808864.
- Tirado, D., W. Nilsson, B. Deyà-Tortella and C. García, 2019: Implementation of water-saving measures in hotels in Mallorca. *Sustainability*, **11**(23), doi:10.3390/su11236880.
- Tishkov, A.A., et al., 2019: Biotic significant climate trends and biota dynamics of the Russian Arctic. *Arctic Ecol. Econ.*, **1**(33), 71–87, doi:10.25283/2223-4594-2019-1-71-87.
- Tjaden, N., et al., 2017: Modelling the effects of global climate change on Chikungunya transmission in the 21st century. *Sci. Rep.*, **7**, doi:10.1038/s41598-017-03566-3.
- Tjaden, N.B., C. Caminade, C. Beierkuhnlein and S.M. Thomas, 2018: Mosquito-borne diseases: advances in modelling climate-change impacts. *Trends Parasitol.*, **34**(3), 227–245, doi:10.1016/j.pt.2017.11.006.
- Tobin, I., et al., 2018a: Vulnerabilities and resilience of European power generation to 1.5°C, 2°C and 3°C warming. *Environ. Res. Lett.*, **13**(4), 44024, doi:10.1088/1748-9326/aab211.
- Tobin, I., et al., 2018b: Vulnerabilities and resilience of European power generation to 1.5°C, 2°C and 3°C warming. *Environ. Res. Lett.*, **13**(4), 44024–44024, doi:10.1088/1748-9326/aab211.
- Tobin, I., et al., 2016: Climate change impacts on the power generation potential of a European mid-century wind farms scenario. *Environ. Res. Lett.*, **11**(3), doi:10.1088/1748-9326/11/3/034013.
- Toimil, A., P. Diaz-Simal, I. Losada and P. Camus, 2018: Estimating the risk of loss of beach recreation value under climate change. *Tour. Manag.*, **68**, 387–400, doi:10.1016/j.tourman.2018.03.024.
- Tokarevich, N., et al., 2017: Impact of air temperature variation on the ixodid ticks habitat and tick-borne encephalitis incidence in the Russian Arctic: the case of the Komi Republic. *Int. J. Circumpolar Health*, **76**, doi:10.1080/22423982.2017.1298882.
- Toreti, A., et al., 2019: The exceptional 2018 European water seesaw calls for action on adaptation. *Earths Future*, **7**(6), 652–663, doi:10.1029/2019EF001170.
- Torrallba, M., et al., 2018: A social-ecological analysis of ecosystem services supply and trade-offs in European wood-pastures. *Sci. Adv.*, **4**(5), eaar2176, doi:10.1126/sciadv.aar2176.
- Townhill, B., et al., 2017: Non-native marine species in north-west Europe: developing an approach to assess future spread using regional downscaled climate projections. *Aquat. Conserv. Mar. Freshw. Ecosyst.*, **27**(5), 1035–1050, doi:10.1002/aqc.2764.
- Townhill, B.L., et al., 2018: Harmful algal blooms and climate change: exploring future distribution changes. *ICES J. Mar. Sci.*, **75**(6), 1882–1893, doi:10.1093/icesjms/fsy113.
- Trnka, M., P. Hlavinka and M. A. Semenov, 2015: Adaptation options for wheat in Europe will be limited by increased adverse weather events under climate change. *J. R. Soc. Interface*, **12**(112), doi:10.1098/rsif.2015.0721.
- Trnka, M., et al., 2014: Adverse weather conditions for European wheat production will become more frequent with climate change. *Nat. Clim. Chang.*, **4**(7), 637–643, doi:10.1038/nclimate2242.
- Tryland, M., et al., 2019: Infectious disease outbreak associated with supplementary feeding of semi-domesticated reindeer. *Front. Vet. Sci.*, **6**, doi:10.3389/fvets.2019.00126.
- Tsikliras, A.C., et al., 2019: Synchronization of Mediterranean pelagic fish populations with the North Atlantic climate variability. *Deep. Sea Res. Part II Top. Stud. Oceanogr.*, **159**, 143–151, doi:10.1016/j.dsr2.2018.07.005.
- Turco, M., et al., 2016: Decreasing fires in Mediterranean Europe. *PLoS ONE*, **11**(3), e150663, doi:10.1371/journal.pone.0150663.
- Turco, M., et al., 2018a: Skilful forecasting of global fire activity using seasonal climate predictions. *Nat. Commun.*, **9**(1), 2718, doi:10.1038/s41467-018-05250-0.
- Turco, M., M.-C. Llasat, J. von Hardenberg and A. Provenzale, 2014: Climate change impacts on wildfires in a Mediterranean environment. *Clim. Chang.*, **125**(3), 369–380, doi:10.1007/s10584-014-1183-3.
- Turco, M., et al., 2018b: Exacerbated fires in Mediterranean Europe due to anthropogenic warming projected with non-stationary climate-fire models. *Nat. Commun.*, **9**(1), 3821, doi:10.1038/s41467-018-06358-z.
- Turco, M., et al., 2017: On the key role of droughts in the dynamics of summer fires in Mediterranean Europe. *Sci. Rep.*, **7**(1), 81, doi:10.1038/s41598-017-00116-9.
- Tzilivakis, J., D.J. Warner, A. Green and K. A. Lewis, 2015: Adapting to climate change: assessing the vulnerability of ecosystem services in Europe in the context of rural development. *Mitig. Adapt. Strateg. Glob. Chang.*, **20**(4), 547–572, doi:10.1007/s11027-013-9507-6.
- Umgiesser, G., 2020: The impact of operating the mobile barriers in Venice (MOSE) under climate change. *J. Nat. Conserv.*, **54**, 125783, doi:10.1016/j.jnc.2019.125783.
- Umgiesser, G., et al., 2021: The prediction of floods in Venice: methods, models and uncertainties. *Nat. Hazards Earth Syst. Sci.*, **21**, 2679–2704, doi:10.5194/nhess-21-2679-2021.
- Urban, M.C., 2015: Accelerating extinction risk from climate change. *Science*, **348**(6234), 571, doi:10.1126/science.aaa4984.
- Ürge-Vorsatz, D., et al., 2018: Locking in positive climate responses in cities. *Nat. Clim. Chang.*, **8**(3), 174–177, doi:10.1038/s41558-018-0100-6.
- Uriarte, I., et al., 2021: Opposite phenological responses of zooplankton to climate along a latitudinal gradient through the European Shelf. *ICES J. Mar. Sci.*, doi:10.1093/icesjms/fsab008.
- Urvois, T., et al., 2021: Climate change impact on the potential geographical distribution of two invading *Xylosandrus ambrosia* beetles. *Sci. Rep.*, **11**(1), 1339, doi:10.1038/s41598-020-80157-9.
- Valade, A., V. Bellassen, C. Magand and S. Luyssaert, 2017: Sustaining the sequestration efficiency of the European forest sector. *For. Ecol. Manag.*, **405**, 44–55, doi:10.1016/j.foreco.2017.09.009.
- Valta, K., et al., 2016: Adaptation measures for the food and beverage industry to the impact of climate change on water availability. *Desalin. Water Treat.*, **57**(5), 2336–2343, doi:10.1080/19443994.2015.1049407.
- van der Kooij, J., G.H. Engelhard and D.A. Righton, 2016: Climate change and squid range expansion in the North Sea. *J. Biogeogr.*, **43**(11), 2285–2298, doi:10.1111/jbi.12847.
- van der Plas, F., et al., 2016: Biotic homogenization can decrease landscape-scale forest multifunctionality. *Proc. Natl. Acad. Sci.*, **113**(13), 3557, doi:10.1073/pnas.1517903113.
- van der Spek, A.J.F., 2018: The development of the tidal basins in the Dutch Wadden Sea until 2100: the impact of accelerated sea-level rise and subsidence on their sediment budget – a synthesis. *Neth. J. Geosci.*, **97**(3), 71–78, doi:10.1017/njg.2018.10.
- van der Velde, M., et al., 2018: In-season performance of European Union wheat forecasts during extreme impacts. *Sci. Rep.*, **8**, doi:10.1038/s41598-018-33688-1.
- Van Dooren, T.J.M., 2019: Assessing species richness trends: Declines of bees and bumblebees in the Netherlands since 1945. *Ecol. Evol.*, **9**(23), 13056–13068, doi:10.1002/ece3.5717.
- van Duinen, R., T. Filatova, P. Geurts and A. van der Veen, 2015: Coping with drought risk: empirical analysis of farmers' drought adaptation in the south-west Netherlands. *Reg. Environ. Change*, **15**(6), 1081–1093, doi:10.1007/s10113-014-0692-y.

- van Hooff, T., B. Blocken, J.L.M. Hensen and H.J.P. Timmermans, 2014: On the predicted effectiveness of climate adaptation measures for residential buildings. *Build. Environ.*, **82**, 300–316, doi:10.1016/j.buildenv.2014.08.027.
- van Klink, R., et al., 2020a: Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances. *Science*, **368**(6489), 417, doi:10.1126/science.aax9931.
- van Klink, R., et al., 2020b: Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances. *Science*, **368**(6489), 417–420, doi:10.1126/science.aax9931.
- van Slobbe, E., et al., 2016: The future of the Rhine: stranded ships and no more salmon? *Reg. Environ. Chang.*, **16**(1), 31–41, doi:10.1007/s10113-014-0683-z.
- van Strien, A., et al., 2019: Over a century of data reveal more than 80% decline in butterflies in the Netherlands. *Biol. Conserv.*, **234**, 116–122, doi:10.1016/j.biocon.2019.03.023.
- van Vliet, A.J.H., et al., 2014: Observed climate-induced changes in plant phenology in the Netherlands. *Reg. Environ. Chang.*, **14**(3), 997–1008.
- van Vliet, M., et al., 2016a: Multi-model assessment of global hydropower and cooling water discharge potential under climate change. *Glob. Environ. Chang. Policy Dimens.*, **40**, 156–170, doi:10.1016/j.gloenvcha.2016.07.007.
- van Vliet, M.T.H., J. Sheffield, D. Wiberg and E.F. Wood, 2016b: Impacts of recent drought and warm years on water resources and electricity supply worldwide. *Environ. Res. Lett.*, **11**(12), 124021, doi:10.1088/1748-9326/11/12/124021.
- van Vliet, M.T.H., D. Wiberg, S. Leduc and K. Riahi, 2016c: Power-generation system vulnerability and adaptation to changes in climate and water resources. *Nat. Clim. Chang.*, **6**, 375–375.
- van Vuuren, D.P., et al., 2015: Pathways to achieve a set of ambitious global sustainability objectives by 2050: explorations using the IMAGE integrated assessment model. *Technol. Forecast. Soc. Chang.*, **98**, 303–323, doi:10.1016/j.techfore.2015.03.005.
- van Vuuren, D.P., et al., 2017: The shared socio-economic pathways: trajectories for human development and global environmental change. *Glob. Environ. Chang.*, **42**, 148–152.
- Vandekerckhove, O., E. De Buck and E. Van Wijngaerden, 2021: Lyme disease in Western Europe: an emerging problem? A systematic review. *Acta Clin. Belg.*, **76**(3), 244–252, doi:10.1080/17843286.2019.1694293.
- Vanos, J.K., J.W. Baldwin, O. Jay and K.L. Ebi, 2020: Simplicity lacks robustness when projecting heat-health outcomes in a changing climate. *Nat. Commun.*, **11**(1), 6079, doi:10.1038/s41467-020-19994-1.
- Varela-Ortega, C., et al., 2016: How can irrigated agriculture adapt to climate change? Insights from the Guadiana Basin in Spain. *Reg. Environ. Chang.*, **16**(1), 59–70, doi:10.1007/s10113-014-0720-y.
- Varrani, A. and M. Nones, 2018: Vulnerability, impacts and assessment of climate change on Jakarta and Venice. *Int. J. River Basin Manag.*, **16**(4), 439–447, doi:10.1080/15715124.2017.1387125.
- Vasilakopoulos, P., D.E. Raitos, E. Tzanatos and C.D. Maravelias, 2017: Resilience and regime shifts in a marine biodiversity hotspot. *Sci. Rep.*, **7**(1), 13647, doi:10.1038/s41598-017-13852-9.
- Vasiliev, D. and S. Greenwood, 2021: The role of climate change in pollinator decline across the Northern Hemisphere is underestimated. *Sci. Total Environ.*, **775**, 145788–145788, doi:10.1016/j.scitotenv.2021.145788.
- Vazquez, D.P., E. Gianoli, W.F. Morris and F. Bozinovic, 2017: Ecological and evolutionary impacts of changing climatic variability. *Biol. Rev.*, **92**(1), 22–42, doi:10.1111/brv.12216.
- Venter, Z.S., N.H. Krog and D.N. Barton, 2020: Linking green infrastructure to urban heat and human health risk mitigation in Oslo, Norway. *Sci. Total Environ.*, **709**, 136193, doi:10.1016/j.scitotenv.2019.136193.
- Verezemskaya, P.S., et al., 2019: *Forecast and analysis of climate change in the Russian Part of the Barents Sea*. WWF Russia, Moscow. 611p.
- Verhagen, W., A.J.A. van Teeffelen and P.H. Verburg, 2018: Shifting spatial priorities for ecosystem services in Europe following land use change. *Ecol. Indic.*, **89**, 397–410, doi:10.1016/j.ecolind.2018.01.019.
- Verkerk, P., et al., 2017: A participatory approach for adapting river basins to climate change. *Water*, **9**(12), 958, doi:10.3390/w9120958.
- Vermaat, J.E., et al., 2017: Differentiating the effects of climate and land use change on European biodiversity: a scenario analysis. *Ambio*, **46**(3), 277–290, doi:10.1007/s13280-016-0840-3.
- Vicedo-Cabrera, A.M., et al., 2021: The burden of heat-related mortality attributable to recent human-induced climate change. *Nat. Clim. Chang.*, **11**(6), 492–500, doi:10.1038/s41558-021-01058-x.
- Vieira, A.R., S. Dores, M. Azevedo and S.E. Tanner, 2019: Otolith increment width-based chronologies disclose temperature and density-dependent effects on demersal fish growth. *ICES J. Mar. Sci.*, **77**(2), 633–644, doi:10.1093/icesjms/fsz243.
- Vilà-Cabrera, A., A.C. Premoli and A.S. Jump, 2019: Refining predictions of population decline at species' rear edges. *Glob. Change Biol.*, **0**(0), doi:10.1111/gcb.14597.
- Virk, G., et al., 2014: The effectiveness of retrofitted green and cool roofs at reducing overheating in a naturally ventilated office in London: direct and indirect effects in current and future climates. *Indoor Built Environ.*, **23**(3), 504–520, doi:10.1177/1420326X14527976.
- Virk, G., et al., 2015: Microclimatic effects of green and cool roofs in London and their impacts on energy use for a typical office building. *Energy Build.*, **88**, 214–228, doi:10.1016/j.enbuild.2014.11.039.
- Visser, M.E., L. te Marvelde and M.E. Lof, 2012: Adaptive phenological mismatches of birds and their food in a warming world. *J. Ornithol.*, **153**(1), 75–84, doi:10.1007/s10336-011-0770-6.
- Vitali, A., et al., 2015: The effect of heat waves on dairy cow mortality. *J. Dairy Sci.*, **98**(7), 4572–4579, doi:10.3168/jds.2015-9331.
- Vitali, V., U. Büntgen and J. Bauhus, 2018: Seasonality matters—The effects of past and projected seasonal climate change on the growth of native and exotic conifer species in Central Europe. *Dendrochronologia*, **48**, 1–9, doi:10.1016/j.dendro.2018.01.001.
- Vitasse, Y., C. Signarbieux and Y.H. Fu, 2018: Global warming leads to more uniform spring phenology across elevations. *Proc. Natl. Acad. Sci. U.S.A.*, **115**(5), 1004–1008, doi:10.1073/pnas.1717342115.
- Viskamp, D.R.M., et al., 2020: First autochthonous human West Nile virus infections in the Netherlands, July to August 2020. *Euro Surveill.*, **25**(46), doi:10.2807/1560-7917.es.2020.25.46.2001904.
- Vodá, R., et al., 2016: Historical and contemporary factors generate unique butterfly communities on islands. *Sci. Rep.*, **6**(1), 28828, doi:10.1038/srep28828.
- Vogel, M.M., et al., 2019: Concurrent 2018 hot extremes across northern hemisphere due to human-induced climate change. *Earth's Future*, **7**(7), 692–703, doi:10.1029/2019ef001189.
- Vogels, C., N. Hartemink and C. Koenraadt, 2017: Modelling West Nile virus transmission risk in Europe: effect of temperature and mosquito biotypes on the basic reproduction number. *Sci. Rep.*, **7**, doi:10.1038/s41598-017-05185-4.
- Voss, R., et al., 2019: Ecological-economic sustainability of the Baltic cod fisheries under ocean warming and acidification. *J. Environ. Manag.*, **238**, 110–118, doi:10.1016/j.jenvman.2019.02.105.
- Vousdoukas, M.I., et al., 2020: Economic motivation for raising coastal flood defenses in Europe. *Nat. Commun.*, **11**(1), 2119, doi:10.1038/s41467-020-15665-3.
- Vousdoukas, M.I., et al., 2018: Climatic and socioeconomic controls of future coastal flood risk in Europe. *Nat. Clim. Chang.*, **8**(9), 776–780, doi:10.1038/s41558-018-0260-4.
- Vuik, V., B.W. Borsje, P.W.J.M. Willemsen and S.N. Jonkman, 2019: Salt marshes for flood risk reduction: quantifying long-term effectiveness and life-cycle costs. *Ocean Coast. Manag.*, **171**, 96–110, doi:10.1016/j.ocecoaman.2019.01.010.
- Wahl, T., et al., 2015: Increasing risk of compound flooding from storm surge and rainfall for major US cities. *Nat. Clim. Chang.*, **5**(12), 1093–1097, doi:10.1038/nclimate2736.

- Waits, A., et al., 2018: Human infectious diseases and the changing climate in the Arctic. *Environ. Int.*, **121**, 703–713, doi:10.1016/j.envint.2018.09.042.
- Wall, M., et al., 2015: pH Up-regulation as a potential mechanism for the cold-water coral *Lophelia pertusa* to sustain growth in aragonite undersaturated conditions. *Biogeosciences*, **12**(23), 6869–6880, doi:10.5194/bg-12-6869-2015.
- Wamelink, G.W.W., et al., 2020: Prediction of plant species occurrence as affected by nitrogen deposition and climate change on a European scale. *Environ. Pollut.*, **266**, 115257, doi:10.1016/j.envpol.2020.115257.
- Wang, H., et al., 2017a: Impacts of global warming on phenology of spring leaf unfolding remain stable in the long run. *Int. J. Biometeorol.*, **61**(2), 287–292, doi:10.1007/s00484-016-1210-3.
- Wang, J., et al., 2020: Anthropogenically-driven increases in the risks of summertime compound hot extremes. *Nat. Commun.*, **11**(1), 528, doi:10.1038/s41467-019-14233-8.
- Wang, S., 2020: Recent global decline of CO₂ fertilization effects on vegetation photosynthesis. *Science*, **370**(6522), 1295–1300, doi:10.1126/science.abb7772.
- Wang, T., et al., 2019: How can the UK road system be adapted to the impacts posed by climate change? By creating a climate adaptation framework. *Transp. Res. Part D. Transp. Environ.*, **77**, 403–424, doi:10.1016/j.trd.2019.02.007.
- Wang, X., R. Brown, G. Prudent-Richard and K. O'Mara, 2016: *Enhancing Power Sector Resilience: Emerging Practices to Manage Weather and Geological Risks*. Energy Sector Management Assistance Program, World Bank Group, Washington, D.C. 125 pp.
- Wang, Y., I. Hendy and T. Napier, 2017b: Climate and anthropogenic controls of coastal deoxygenation on Interannual to centennial timescales. *Geophys. Res. Lett.*, **44**(22), 11528–11536, doi:10.1002/2017GL075443.
- Wang, Z. B., E.P.L. Elias, A.J.F. van der Spek and Q.J. Lodder, 2018: Sediment budget and morphological development of the Dutch Wadden Sea: impact of accelerated sea-level rise and subsidence until 2100. *Neth. J. Geosci.*, **97**(3), 183–214, doi:10.1017/njg.2018.8.
- Ward, K., S. Lauf, B. Kleinschmit and W. Endlicher, 2016: Heat waves and urban heat islands in Europe: a review of relevant drivers. *Sci. Total. Environ.*, **569**, 527–539, doi:10.1016/j.scitotenv.2016.06.119.
- Warren, R., et al., 2018: The projected effect on insects, vertebrates, and plants of limiting global warming to 1.5°C rather than 2°C. *Science*, **360**(6390), 791–795, doi:10.1126/science.aar3646.
- Wasmund, N., et al., 2019: Extension of the growing season of phytoplankton in the western Baltic Sea in response to climate change. *Mar. Ecol. Prog. Ser.*, **622**, 1–16.
- Webber, H., et al., 2018: Diverging importance of drought stress for maize and winter wheat in Europe. *Nat. Commun.*, **9**(1), 4249, doi:10.1038/s41467-018-06525-2.
- Webber, H., et al., 2016: Uncertainty in future irrigation water demand and risk of crop failure for maize in Europe. *Environ. Res. Lett.*, **11**(7), 1–10, doi:10.1088/1748-9326/11/7/074007.
- Wiens, J.J., 2016: Climate-related local extinctions are already widespread among plant and animal species. *PLoS Biol.*, **14**(12), e2001104, doi:10.1371/journal.pbio.2001104.
- Williges, K., R. Mechler, P. Bowyer and J. Balkovic, 2017: Towards an assessment of adaptive capacity of the European agricultural sector to droughts. *Clim. Serv.*, **7**, 47–63, doi:10.1016/j.cliser.2016.10.003.
- Wilson, R.J., D. Gutiérrez, J. Gutiérrez and V.J. Monserrat, 2007: An elevational shift in butterfly species richness and composition accompanying recent climate change. *Glob. Change Biol.*, **13**(9), 1873–1887, doi:10.1111/j.1365-2486.2007.01418.x.
- Wimmer, F., et al., 2014: Modelling the effects of cross-sectoral water allocation schemes in Europe. *Clim. Change*, **128**(3-4), 229–244, doi:10.1007/s10584-014-1161-9.
- Wiréhn, L., 2018: Nordic agriculture under climate change: a systematic review of challenges, opportunities and adaptation strategies for crop production. *Land Use Policy*, **77**, 63–74, doi:10.1016/j.landusepol.2018.04.059.
- Wiréhn, L., J. Käyhkö, T.-S. Neset and S. Juhola, 2020: Analysing trade-offs in adaptation decision-making—agricultural management under climate change in Finland and Sweden. *Reg. Environ. Chang.*, **20**(1), 18–18, doi:10.1007/s10113-020-01585-x.
- Wreford, A. and C.F.E. Topp, 2020: Impacts of climate change on livestock and possible adaptations: a case study of the United Kingdom. *Agric. Syst.*, **178**, 102737–102737, doi:10.1016/j.agsy.2019.102737.
- Wu, C., et al., 2018: Contrasting responses of autumn-leaf senescence to daytime and night-time warming. *Nat. Clim. Chang.*, **8**(12), 1092–1096, doi:10.1038/s41558-018-0346-z.
- Wu, M., et al., 2015a: Sensitivity of burned area in Europe to climate change, atmospheric CO₂ levels, and demography: a comparison of two fire-vegetation models. *J. Geophys. Res. Biogeosci.*, **120**(11), 2256–2272, doi:10.1002/2015JG003036.
- Wu, M., et al., 2015b: Sensitivity of burned area in Europe to climate change, atmospheric CO₂ levels, and demography: a comparison of two fire-vegetation models. *J. Geophys. Res. Biogeosci.*, **120**(11), 2256–2272, doi:10.1002/2015JG003036.
- Xi, Y., 2020: Future impacts of climate change on inland Ramsar wetlands. *Nat. Clim. Chang.*, **11**(1), 45–51.
- Xi, Y., S. Peng, P. Ciais and Y. Chen, 2021: Future impacts of climate change on inland Ramsar wetlands. *Nat. Clim. Chang.*, **11**(1), 45–51, doi:10.1038/s41558-020-00942-2.
- Xu, C., et al., 2019: Increasing impacts of extreme droughts on vegetation productivity under climate change. *Nat. Clim. Chang.*, **9**(12), 948–953, doi:10.1038/s41558-019-0630-6.
- Yigini, Y. and P. Panagos, 2016: Assessment of soil organic carbon stocks under future climate and land cover changes in Europe. *Sci. Total. Environ.*, **557–558**, 838–850, doi:10.1016/j.scitotenv.2016.03.085.
- Young, J.J., et al., 2021: Epidemiology of human West Nile virus infections in the European Union and European Union enlargement countries, 2010 to 2018. *Euro Surveill.*, **26**(19), doi:10.2807/1560-7917.es.2021.26.19.2001095.
- Yu, J., P. Berry, B.P. Guillod and T. Hickler, 2021: Climate change impacts on the future of forests in Great Britain. *Front. Environ. Sci.*, **9**, 83.
- Yuan, W., et al., 2019: Increased atmospheric vapor pressure deficit reduces global vegetation growth. *Sci. Adv.*, **5**(8), 1–13, doi:10.1126/sciadv.aax1396.
- Zakharov, A.I. and R.B. Sharipova, 2017: Agro climate potential and basic problems of influence of climate changes on agricultural crop production in Ulyanovsk region. *Вестник Ульяновской Государственной Сельскохозяйственной Академии*, **1**(37), 25–30, doi:10.18286/1816-4501-2017-1-25-30.
- Zattara, E.E. and M.A. Aizen, 2021: Worldwide occurrence records reflect a global decline in bee species richness. *One Earth*, **4**(1): 114–123, doi:10.1101/869784.
- Zellweger, F., et al., 2020: Forest microclimate dynamics drive plant responses to warming. *Science*, **368**(6492), 772, doi:10.1126/science.aba6880.
- Zeng, Z.L., et al., 2021: Global, regional, and national dengue burden from 1990 to 2017: a systematic analysis based on the global burden of disease study 2017. *Eclinicalmedicine*, **32**, doi:10.1016/j.eclinm.2020.100712.
- Zhang, Y., et al., 2020: Large and projected strengthening moisture limitation on end-of-season photosynthesis. *Proc. Natl. Acad. Sci.*, **117**(17), 9216, doi:10.1073/pnas.1914436117.
- Zhang, Y.L., et al., 2017: Global loss of aquatic vegetation in lakes. *Earth Sci. Rev.*, **173**, 259–265, doi:10.1016/j.earscirev.2017.08.013.
- Zhao, C., et al., 2017: Temperature increase reduces global yields of major crops in four independent estimates. *Proc. Natl. Acad. Sci.*, **114**(35), 9326–9331, doi:10.1073/pnas.1701762114.
- Zhou, S., et al., 2019: Land–atmosphere feedbacks exacerbate concurrent soil drought and atmospheric aridity. *Proc. Natl. Acad. Sci. U.S.A.*, **116**(38), 18848–18853, doi:10.1073/pnas.1904955116.

- Zinzi, M., 2016: Characterisation and assessment of near infrared reflective paintings for building facade applications. *Energy Build.*, **114**, 206–213, doi:10.1016/j.enbuild.2015.05.048.
- Zinzi, M., et al., 2017: Assessing the overheating risks in Italian existing school buildings renovated with nZEB targets. *Energy Procedia*, **142**, 2517–2524, doi:10.1016/j.egypro.2017.12.192.
- Zlatanov, T., C. Elkin, F. Irauschek and M.J. Lexer, 2017: Impact of climate change on vulnerability of forests and ecosystem service supply in Western Rhodopes Mountains. *Reg. Environ. Chang.*, **17**(1), 79–91, doi:10.1007/s10113-015-0869-z.
- Zografou, K., et al., 2014: Signals of climate change in butterfly communities in a Mediterranean protected area. *PLoS ONE*, **9**(1), e87245, doi:10.1371/journal.pone.0087245.
- Zölch, T., L. Henze, P. Keilholz and S. Pauleit, 2017: Regulating urban surface runoff through nature-based solutions – an assessment at the micro-scale. *Environ. Res.*, **157**, 135–144, doi:10.1016/j.envres.2017.05.023.
- Zommers, Z., et al., 2020: Burning embers: towards more transparent and robust climate-change risk assessments. *Nat. Rev. Earth Environ.*, **1**(10), 516–529, doi:10.1038/s43017-020-0088-0.