

# Sea Level Change

## Supplementary Material

### Coordinating Lead Authors:

John A. Church (Australia), Peter U. Clark (USA)

### Lead Authors:

Anny Cazenave (France), Jonathan M. Gregory (UK), Svetlana Jevrejeva (UK), Anders Levermann (Germany), Mark A. Merrifield (USA), Glenn A. Milne (Canada), R. Steven Nerem (USA), Patrick D. Nunn (Australia), Antony J. Payne (UK), W. Tad Pfeffer (USA), Detlef Stammer (Germany), Alakkat S. Unnikrishnan (India)

### Contributing Authors:

David Bahr (USA), Jason E. Box (Denmark/USA), David H. Bromwich (USA), Mark Carson (Germany), William Collins (UK), Xavier Fettweis (Belgium), Piers Forster (UK), Alex Gardner (USA), W. Roland Gehrels (UK), Rianne Giesen (Netherlands), Peter J. Gleckler (USA), Peter Good (UK), Rune Grand Graversen (Sweden), Ralf Greve (Japan), Stephen Griffies (USA), Edward Hanna (UK), Mark Hemer (Australia), Regine Hock (USA), Simon J. Holgate (UK), John Hunter (Australia), Philippe Huybrechts (Belgium), Gregory Johnson (USA), Ian Joughin (USA), Georg Kaser (Austria), Caroline Katsman (Netherlands), Leonard Konikow (USA), Gerhard Krinner (France), Anne Le Brocq (UK), Jan Lenaerts (Netherlands), Stefan Ligtenberg (Netherlands), Christopher M. Little (USA), Ben Marzeion (Austria), Kathleen L. McInnes (Australia), Sebastian H. Mernild (USA), Didier Monselesan (Australia), Ruth Mottram (Denmark), Tavi Murray (UK), Gunnar Myhre (Norway), J.P. Nicholas (USA), Faezeh Nick (Norway), Mahé Perrette (Germany), David Pollard (USA), Valentina Radić (Canada), Jamie Rae (UK), Markku Rummukainen (Sweden), Christian Schoof (Canada), Aimée Slangen (Australia/Netherlands), Jan H. van Angelen (Netherlands), Willem Jan van de Berg (Netherlands), Michiel van den Broeke (Netherlands), Miren Vizcaino (Netherlands), Yoshihide Wada (Netherlands), Neil J. White (Australia), Ricarda Winkelmann (Germany), Jianjun Yin (USA), Masakazu Yoshimori (Japan), Kirsten Zickfeld (Canada)

### Review Editors:

Jean Jouzel (France), Roderik van de Wal (Netherlands), Philip L. Woodworth (UK), Cunde Xiao (China)

### This chapter supplementary material should be cited as:

Church, J.A., P.U. Clark, A. Cazenave, J.M. Gregory, S. Jevrejeva, A. Levermann, M.A. Merrifield, G.A. Milne, R.S. Nerem, P.D. Nunn, A.J. Payne, W.T. Pfeffer, D. Stammer and A.S. Unnikrishnan, 2013: Sea Level Change Supplementary Material. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Available from [www.climate-change2013.org](http://www.climate-change2013.org) and [www.ipcc.ch](http://www.ipcc.ch).

# Table of Contents

13.SM.1	Methods of Global Mean Sea Level Projections for the 21st Century.....	13SM-3
13.SM.2	Computation of Regional Maps of Sea Level Change from Coupled Model Intercomparison Project Phase 5 Model Output .....	13SM-5
References	.....	13SM-8

### 13.SM.1 Methods of Global Mean Sea Level Projections for the 21st Century

This section summarizes the methods used to produce the projections shown in Section 13.5.1 for the Representative Concentration Pathway (RCP) scenarios and the Special Report on Emission Scenarios (SRES) A1B scenario. The Supplementary Material includes files of the annual time series of median, 5th percentile and 95th percentile for each of the contributions to global mean sea level rise and the sum, corresponding to the results shown in Table 13.5. The data files are named as follows:

*scenario \_ quantity statistic . suffix*

for instance *rcp45\_summid.nc*. In each name,

*scenario* is **rcp26**, **rcp45**, **rcp60** or **rcp85**, corresponding to the four representative concentration pathways used in CMIP5, or **sresa1b** for SRES A1B used in CMIP3.

*quantity* is **temperature** for global mean surface temperature change, **expansion** for thermal expansion (sections 13.4.1 and 13.SM.1.2), **glacier** for glaciers (13.4.2 and 13.SM.1.3), **greensmb** for Greenland ice-sheet SMB (13.4.3.1 and 13.SM.1.4), **antsmb** for Antarctic ice-sheet SMB (13.4.4.1 and 13.SM.1.5), **greendyn** for Greenland ice-sheet rapid dynamical change (13.4.3.2 and 13.SM.1.6), **antdyn** for Antarctic ice-sheet rapid dynamical change (13.4.4.2 and 13.SM.1.6), **landwater** for anthropogenic intervention in water storage on land (13.4.5 and 13.SM.1.6), **greennet** for the sum of SMB and rapid dynamical contributions from the Greenland ice-sheet, **antnet** for the sum of SMB and rapid dynamical contributions from the Antarctic ice-sheet, **sheetdyn** for the sum of the rapid dynamical contributions from the Greenland and Antarctic ice-sheets, or **sum** for the sea level projection including all contributions. Except for temperature, these are the quantities shown in Table 13.5.

*statistic* is **mid** for the median, or **lower** or **upper** for the limits of the range.

*suffix* is **txt** for plain ASCII text, or **nc** for netCDF.

The text files have two columns, year and sea level change in metres. The netCDF files describe their contents using the CF convention.

#### 13.SM.1.1 Derivation of Global Surface Temperature and Thermal Expansion Time Series from Coupled Model Intercomparison Project Phase 5

Annual time series for change in global mean surface air temperature (SAT) ('tas' in the CMIP5 archive) and global-mean sea level (GMSL) rise due to thermal expansion ('zostoga') in the historical period and during the 21st century under RCP scenarios (Section 13.4.1) were obtained from a set of 21 CMIP5 AOGCMs (ACCESS1-0, ACCESS1-3, CCSM4, CNRM-CM5, CSIRO-Mk3-6-0, CanESM2, GFDL-CM3, GFDL-ES-M2G, GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR, IPSL-CM5A-MR, MIROC-ESM, MIROC-ESM-CHEM, MIROC5, MPI-ESM-LR, MPI-ESM-MR, MRI-CGCM3, NorESM1-M, NorESM1-ME, Inmcm4). These were all those for which thermal expansion was available, including from a parallel pre-industrial control experiment, which is required to remove the thermal expansion due to climate drift in deep-ocean temperatures (Gleckler et al., 2012). The drift was removed by subtracting a polynomial fit as a function of time to the control thermal expansion time series. Where CMIP5 results were not available for a particular Atmosphere–Ocean General Circulation Model (AOGCM) and scenario, they were estimated by the method of Good et al. (2011) and Good et al. (2013) using the response of that AOGCM to an instantaneous quadrupling of carbon dioxide (CO<sub>2</sub>) concentration. The same method was used to estimate the CMIP5 projections for scenario SRES A1B. The method gives estimates of change in global mean surface air temperature and net radiative flux at the top of the atmosphere. The

**Table 13.SM.1 |** Median values and *likely* ranges for projections of global-mean sea level rise and its contributions in metres in 2100 relative to 1986–2005 for the four RCP scenarios and SRES A1B. See Section 13.5.1 concerning how the *likely* range is defined. Because some of the uncertainties in modelling the contributions are treated as uncorrelated, the sum of the lower bound of contributions does not equal the lower bound of the sum, and similarly for the upper bound. Because of imprecision from rounding, the sum of the medians of contributions may not exactly equal the median of the sum.

	SRES A1B	RCP2.6	RCP4.5	RCP6.0	RCP8.5
Thermal expansion	0.24 [0.18 to 0.30]	0.15 [0.11 to 0.20]	0.20 [0.15 to 0.25]	0.22 [0.17 to 0.27]	0.32 [0.25 to 0.39]
Glaciers	0.16 [0.09 to 0.23]	0.11 [0.05 to 0.17]	0.13 [0.07 to 0.20]	0.14 [0.07 to 0.20]	0.18 [0.10 to 0.26]
Greenland Ice Sheet SMB <sup>a</sup>	0.07 [0.03 to 0.15]	0.03 [0.01 to 0.08]	0.05 [0.02 to 0.11]	0.05 [0.02 to 0.12]	0.10 [0.04 to 0.22]
Antarctic Ice Sheet SMB <sup>b</sup>	−0.04 [−0.07 to −0.01]	−0.02 [−0.05 to −0.00]	−0.03 [−0.06 to −0.01]	−0.03 [−0.06 to −0.01]	−0.05 [−0.09 to −0.02]
Greenland Ice Sheet Rapid Dynamics	0.04 [0.01 to 0.06]	0.04 [0.01 to 0.06]	0.04 [0.01 to 0.06]	0.04 [0.01 to 0.06]	0.05 [0.02 to 0.09]
Antarctic Ice Sheet Rapid Dynamics	0.08 [−0.02 to 0.19]	0.08 [−0.02 to 0.19]	0.08 [−0.02 to 0.19]	0.08 [−0.02 to 0.19]	0.08 [−0.02 to 0.19]
Land Water Storage	0.05 [−0.01 to 0.11]	0.05 [−0.01 to 0.11]	0.05 [−0.01 to 0.11]	0.05 [−0.01 to 0.11]	0.05 [−0.01 to 0.11]
Sea Level Rise	0.60 [0.42 to 0.80]	0.44 [0.28 to 0.61]	0.53 [0.36 to 0.71]	0.55 [0.38 to 0.73]	0.74 [0.52 to 0.98]
Greenland Ice Sheet	0.11 [0.07 to 0.19]	0.08 [0.04 to 0.12]	0.09 [0.05 to 0.16]	0.09 [0.06 to 0.16]	0.15 [0.09 to 0.28]
Antarctic Ice Sheet	0.05 [−0.06 to 0.15]	0.06 [−0.04 to 0.16]	0.05 [−0.05 to 0.15]	0.05 [−0.05 to 0.15]	0.04 [−0.08 to 0.14]
Ice-Sheet Rapid Dynamics	0.12 [0.03 to 0.22]	0.12 [0.03 to 0.22]	0.12 [0.03 to 0.22]	0.12 [0.03 to 0.22]	0.14 [0.04 to 0.24]

Only the collapse of the marine-based sectors of the Antarctic Ice Sheet could cause GMSL to rise substantially above the *likely* range during the 21st century. This potential additional contribution cannot be precisely quantified but there is *medium confidence* that it would not exceed several tenths of a meter of sea level rise.

Notes:

<sup>a</sup> Including the height-SMB feedback.

<sup>b</sup> Including the interaction between SMB change and outflow.

latter was integrated in time to obtain the estimated change in heat content of the climate system, and converted to thermal expansion using the expansion efficiency of heat appropriate to each AOGCM, as diagnosed from all the available RCPs for that AOGCM. The correlation between heat content change and thermal expansion is very high and the relationship can be accurately treated as linear (Kuhlbrodt and Gregory, 2012).

### 13.SM.1.2 Interpretation and Combination of Uncertainties

Uncertainties were derived from the CMIP5 ensemble by treating the model spread as a normal distribution, and following Section 12.4.1.2 it was assumed that the 5 to 95% interval of CMIP5 projections for the 21st century for each RCP scenario can be interpreted as a *likely* range (Section 13.5.1). The CMIP5 timeseries of thermal expansion  $X$  and global mean surface air temperature  $T$  were expressed as anomalies as a function of time  $t$  with respect to their time-means for 1986–2005, and the timeseries of ensemble means  $X_M(t)$  and  $T_M(t)$  and ensemble standard deviations  $X_S(t)$  and  $T_S(t)$  were calculated. As in the AR4, a Monte Carlo was used to generate distributions of timeseries of  $X$  and  $T$  in a perfectly correlated way; for each member of the ensemble, a random number  $r$  was chosen from a normal distribution, giving  $X(t) = X_M(t) + r X_S(t)$  and  $T(t) = T_M(t) + r T_S(t)$ , and  $T(t)$  was used to estimate land ice contributions to GMSLR, as described in the following sections. As in the AR4, all the uncertainties described by the land ice methods were assumed to be independent of the climate change uncertainty represented by the variation of  $r$  and of one another, except where stated, and were combined by Monte Carlo. Because of the use of Monte Carlo, the results for GMSLR have a random uncertainty. For different random samples of the sizes used to compute the results in Table 13.5, the results vary by up to 0.01 m in GMSLR and its contributions, and 0.1 mm yr<sup>-1</sup> in the rate of GMSLR. The projections are shown for 2081–2100 in Table 13.5, and for 2100 in Table 13.SM.1.

### 13.SM.1.3 Glaciers

Changes in glacier mass in all regions excluding Antarctica from 2006 onwards were projected using a parameterized scheme which was fitted separately to results from each of the global glacier models of Giesen and Oerlemans (2013), Marzeion et al. (2012), Radić et al. (2014) and Slangen and van de Wal (2011). For the model of Giesen and Oerlemans (2013), only the dependence on temperature was considered; the dependences on precipitation and atmospheric transmissivity were not included. All of these global glacier models have been used to make projections using output from several AOGCMs. Giesen and Oerlemans used results from CMIP3 AOGCMs for scenario SRES A1B, and the other authors used results from different sets of CMIP5 AOGCMs for RCPs. The RCP results of Slangen and van de Wal (2011) are not included in their published paper, but use the same glacier model as in the paper. The parameterized scheme enables estimates to be made for the glacier contribution to GMSL rise  $g_i$  as a function of time  $t$  for the consistent set of CMIP3 and CMIP5 AOGCMs across all RCPs and SRES A1B. The scheme gives  $g_i(t)$  in millimetres with respect to 2006 as  $fI(t)^p$ , where  $I(t)$  is the time integral of  $T$  from 2006 to time  $t$  in degrees Celsius year, and the constants  $f$  and  $p$  used for each glacier model are shown in Table 13.SM.2. The constants were fitted by linear regression of  $\log(g)$  against  $\log(I)$ . The global glacier models on

which this formula is based calculate their results from geographically dependent climate change with detailed treatments of glacier surface mass balance (SMB) and the evolution of hypsometry; their complexity cannot be accurately reproduced by a simple formula, and the spread of their results around the prediction of this formula has a coefficient of variation (standard deviation divided by mean) of 20% or less for decadal means for all glacier models and RCPs, except for the early decades of the 21st century under RCP2.6 for the model of Slangen and van de Wal (2011) for which there are fractional errors of up to 40%, but the absolute error is small. Therefore we take 20% of the projection of the formula made using the CMIP5 ensemble mean  $I(t)$  as the standard deviation of a normally distributed methodological uncertainty in the glacier projection for each global glacier model. In order to incorporate this uncertainty into the projections, for each member of the Monte Carlo ensemble of glacier time-series, a normally distributed random number was chosen, independent of time, as a factor by which the time-dependent standard deviation should be multiplied, giving the uncertainty to be added to the glacier time-series. We give the four global glacier models equal weight in the projections. Because the time integration began in 2006, a constant 9.5 mm was added to the projections to account for the glacier contribution from 1996 (the centre of the reference period for projections) to 2005; this is the mean result from the model of Marzeion et al. (2012) using input from CMIP5 AOGCM historical experiments. The formula is not applicable beyond 2100 because it does not represent the tendency of global glacier mass to reach a new steady value when global climate stabilizes, although the global glacier models on which it is based can predict this as a consequence of the evolution of hypsometry. Glaciers on Antarctica were not included in the global glacier projections because they are included in the projections for the Antarctic ice sheet.

**Table 13.SM.2** | Parameters for the fits to the global glacier models.

Global Glacier Model	$f$ (mm °C <sup>-1</sup> yr <sup>-1</sup> )	$p$ (no unit)
Giesen and Oerlemans (2013)	3.02	0.733
Marzeion et al. (2012)	4.96	0.685
Radić et al. (2013)	5.45	0.676
Slangen and van de Wal (2011)	3.44	0.742

### 13.SM.1.4 Greenland Ice Sheet Surface Mass Balance

The change in Greenland ice sheet SMB  $G_e(t)$ , excluding changes in ice sheet topography, was computed from  $T(t)$  using the cubic polynomial formula, Equation (2) of Fettweis et al., which predicts the Greenland SMB anomaly as a function of  $T$ , and was obtained by fitting results from an RCM using input from several CMIP5 AOGCMs for RCP4.5 and RCP8.5. Their Equation (2)  $G_e = -71.5T - 20.4T^2 - 2.8T^3$  gives  $G_e$  in Gt yr<sup>-1</sup>, which we convert to mm yr<sup>-1</sup> SLE. In this formula,  $T$  is relative to the time mean of 1980–1999, rather than 1986–2005; in the CMIP5 AOGCM results, the former period is cooler by 0.15°C. The results of this formula were compared with those for the same AOGCMs and RCP from Equation (1) of Fettweis et al. (2013), which predicts  $G(t)$  from summer (June to August) air temperature at 600 hPa over Greenland. Equation (1) reproduces the RCM results more accurately but cannot be used for the consistent set of CMIP5 AOGCMs and all RCPs because

their required input data are not available. The results of Equation (2) were also compared with those for the same AOGCMs and RCPs with results obtained from the models of Gregory and Huybrechts (2006) and Yoshimori and Abe-Ouchi (2012), the former being the one used in the AR4. As a result of this comparison of projections (Section 13.4.3.1, Table 13.4),  $G_e(t)$  was estimated as  $FG_2(t)$ , where  $G_2(t)$  is calculated from  $T$  using Fettweis et al. Equation (2), and  $F$  is a factor representing methodological uncertainty. This factor is taken to have a log-normal distribution i.e. one of the form  $F = e^N$ , where  $N$  is a normal distribution having a mean of zero. A log-normal distribution is used because the distributions of  $G_e(t)$  from the various Greenland ice sheet SMB models are positively skewed. None of these models simulates the change in SMB caused by the evolution of the ice sheet surface topography, which gives a positive feedback on mass loss (Section 13.4.3.2). To allow for this effect, the Greenland ice sheet SMB change  $G(t)$  with respect to 1986–2005 was estimated as  $EG_e(t)$ , where  $E$  is a randomly varying factor with a uniform probability distribution in the range 1.00 to 1.15. The uncertainties of  $E$  and  $F$  were assumed not be correlated, and independent of time. The ice sheet SMB change  $G(t)$  was integrated in time to obtain the change in ice sheet mass, starting in 2006. A constant 1.5 mm was added to the projections to account for the Greenland SMB contribution from 1996 (the centre of the reference period for projections) to 2005; this is half of the central observational estimate of the rate of Greenland ice sheet mass loss during this period (Section 13.3.3.2, using data presented in Figure 4.15).

### 13.SM.1.5 Antarctic Ice Sheet Surface Mass Balance

The change in Antarctic ice sheet SMB  $A(t)$  with respect to 1986–2005 was assumed to be due solely to an increase in accumulation (thus,  $A < 0$  in units of sea level equivalent, because accumulation on the ice sheet removes mass from the ocean), which was estimated using the results of Gregory and Huybrechts (2006) from CMIP3 AOGCMs. Accumulation was taken to increase at  $5.1 \pm 1.5\% \text{ } ^\circ\text{C}^{-1}$  of warming in Antarctica relative to 1985–2005, the ratio of warming in Antarctic to  $T$  was taken to be  $1.1 \pm 0.2$ , and the accumulation for the reference period was taken to be  $1923 \text{ Gt yr}^{-1}$  (Section 13.3.3.2). Both of these uncertainties (standard deviations) were treated as normally distributed methodological uncertainties in the projections. The resulting spread of projections is very close to the spread of the results from the high-resolution Antarctic SMB models of Krinner et al. (2007), Bengtsson et al. (2011) and Ligtenberg et al. (2013) assessed in Section 13.4.4.1. The effect of increased accumulation on the dynamics of the Antarctic ice sheet (Section 13.4.4.2) was taken into account by adding a rate  $-SA(t)$  (a positive number in units of sea level equivalent, because the increase in outflow opposes the increase in accumulation and adds mass to the ocean) to the GMSL projections, where  $S$  is a randomly varying factor with a uniform probability distribution in the range 0.00 to 0.35. The uncertainties in accumulation sensitivity, Antarctic warming ratio, and the factor  $S$  were assumed not to be correlated, but  $S$  was perfectly correlated with the distribution of Antarctic rapid ice sheet dynamics (next paragraph), in the sense that when the rapid dynamical increase in outflow is large, the increase in outflow due to the dynamical reaction to increased accumulation is also large. The mass balance changes  $A$  and  $-SA$  were integrated in time to obtain the change in the ice sheet mass, starting from 2006. Unlike for Greenland ice sheet SMB, no addition to the projections was made to account for the period

1996–2005 for the contribution from Antarctic ice-sheet SMB, because changes during this period are judged to be due solely to dynamical change (Section 13.3.3.2).

### 13.SM.1.6 Rapid Ice Sheet Dynamics and Anthropogenic Change in Land Water Storage

Following Section 13.3.3.2, the contributions from rapid ice-sheet dynamics at the start of the projections were taken to be half of the observed rate of loss for 2005–2010 from Greenland (half of  $0.46\text{--}0.80 \text{ mm yr}^{-1}$  from Table 4.6) and all of that from Antarctica ( $0.21\text{--}0.61 \text{ mm yr}^{-1}$  from Table 4.6). The contributions reach  $0.020$  to  $0.085 \text{ m}$  at 2100 from Greenland for RCP8.5,  $0.014$  to  $0.063 \text{ m}$  for the other RCPs and  $-0.020$  to  $0.185 \text{ m}$  from Antarctica for all RCPs; these are the likely ranges from our assessment of existing studies (Sections 13.4.3.2 and 13.4.4.2). For each ice sheet, a quadratic function of time was fitted which begins at the minimal initial rate and reaches the minimum final amount, and another for the maxima. Time series for the rapid dynamic contribution lying between these extremes were constructed as combinations of the extreme time series assuming a uniform probability density between the extremes. Finally, a constant 1.5 mm was added to the contribution from the Greenland ice sheet, and 2.5 mm to the contribution from the Antarctic ice sheet, these being the estimates of those contributions from 1996 to 2005 (using the data presented in Figures 4.15 and 4.16).

The same method was followed for the anthropogenic land water storage contribution (initial rates as for 1993–2010 from Table 13.1 and amounts for the time-mean of 2081–2100 from Section 13.4.5, with no additional amount for land water storage from 1996 to 2005). These contributions are treated as uncorrelated with the magnitude of global climate change and as independent of scenario (except for the higher rate of change for Greenland ice sheet outflow under RCP8.5). This treatment does not imply that the contributions concerned will not depend on the scenario followed, only that the current state of knowledge does not permit a quantitative assessment of the dependence.

## 13.SM.2 Computation of Regional Maps of Sea Level Change from Coupled Model Intercomparison Project Phase 5 Model Output

Several results and figures in Section 13.6 are based on published methods as referred to in the main text but have not been published independently. This document details all information that led to numbers and figures shown in Section 13.6 on regional sea level projections. Data files for each figure are available.

For each figure or each step involved, the underlying technical details that were used are described. The Supplementary Material includes files containing the data in each case.

Figures 13.15, 13.16 and 13.24 show maps of regional sea level changes computed from CMIP5 coupled climate models. The following steps were pursued in the preparation of those figures.

### 13.SM.2.1 Sea Surface Height from Coupled Climate Models

Sea surface height (SSH) data, labeled the 'zos' variable, from the CMIP5 AOGCM database, are used to show regional changes in SSH over time, and include the regional variability of dynamic topography changes due to water mass advection, thermohaline circulation and to the wind-driven circulation (see Table 13.SM.3). These regional changes are corrected for regional control drift by removing the linearly fitted control run drift from each latitude–longitude grid box individually, on a per-model basis. After this correction, the global average of this regional SSH field (a function of  $x, y, t$ ) is forced to be the global thermal expansion ('zostoga' variable) at each time step by first subtracting the globally averaged regional SSH field at each time step from each grid box, and then adding the global thermal expansion time series to each grid box (the same number at every grid box, for a given time). The global thermal expansion time series was also corrected for control drift by removing a quadratic fit to the control run's thermal expansion time series before being added to the regional SSH data. As not all models had multiple ensemble forced runs for the various RCP scenarios, only one run from each model (in each RCP scenario) was used to compute the multi-model ensemble means (i.e., the results for each individual model are only a single realization per scenario, as shown in Figure 13.24).

### 13.SM.2.2 Interpolation

All of the steps outlined above were performed on each model's own grid, with interpolation to a common  $1^\circ \times 1^\circ$  grid only being applied after statistical analyses, to each model's relative sea level changes, means and variances. The interpolation procedure involves applying a nearest-neighbour interpolation and a bilinear interpolation, with the nearest-neighbour interpolation chosen close to the coasts where the bilinear interpolation loses grid boxes.

### 13.SM.2.3 Masking

Some of the models, on their original grids, had detached marginal seas (e.g., the Mediterranean, Hudson Bay, Baltic Sea, etc.), and in most cases, the SSH in the marginal seas behaved differently than in the nearby ocean, with some models having significant numerical instability, and others undergoing a different SSH evolution in these seas. To remove large and obvious errors from the ensemble mean (and other ensemble statistics) and to treat all the models consistently, marginal seas were masked out from individual models, if they were detached from the adjacent ocean basin, on the common  $1^\circ \times 1^\circ$  grid. This results in a final ensemble mean product that consists of, for example, for the RCP4.5 run, a 21-model mean over most of the ocean, but has only as few as 12 ensemble members contributing to the mean for some marginal seas (9 is the lowest number of RCP4.5/8.5 members for which regional data are shown for ensemble statistics).

### 13.SM.2.4 Combining All Sea Level Rise Components

Figures 13.18, 13.19, 13.22 and 13.23 show projected sea level changes as they result after combining various different contributions to sea level change in addition to those available from CMIP5 models. The following steps were necessary to obtain those maps and figures.

Contributions to regional sea level change due to changes in other components of the climate system were added to the thermosteric/dynamic SSH from the AOGCMs. These components include surface mass balance and dynamic ice sheet contributions from Greenland and Antarctica, a glacier contribution, a land water storage contribution, glacial isostatic adjustment (GIA), and the inverse barometer effect (IBE). The projections of the various land ice contributions and the land water storage contribution are described elsewhere (Sections 13.4, 13.5.1 and 13.SM.1 in the Supplementary Material). These global estimates were turned into regional maps of sea level response, due to the addition of mass increasing the global ocean volume (the barystatic contribution) plus the resultant gravitational and rotational changes, through application of an iterative sea level equation solver (Slangen et al., 2012). The groundwater storage change contribution to regional sea level rise was also found similarly by taking estimates of its geographical distribution from Wada et al. (2012) and applying the same sea level equation solver. The GIA contribution was calculated from the mean of the ICE-5G model (Peltier 2004) and the ANU ice sheet model (Lambeck et al. 1998 and subsequent improvements) with the SELEN code for the sea level equation (Farrell and Clark 1976; Spada and Stocchi 2006, 2007), including updates to allow for coastline variation through time, near-field meltwater damping and Earth rotation in a self-consistent manner (Milne and Mitrovica, 1998; Kendall et al., 2006). The IBE contribution was found by using an ensemble of atmospheric results from the atmospheric component of the same CMIP5 models used for the SSH data. All of these components were calculated 'offline' (i.e., were not part of diagnostic 'zos' and 'zostoga' variables in the models) and then added to the regional sea level rise results previously derived from CMIP5 'zos' and 'zostoga' variables.

### 13.SM.2.5 Uncertainties

Figures 13.19, 13.21 and 13.23 show uncertainty measures for sea level projections. Those uncertainties were computed as follows.

The uncertainties in the results directly from the CMIP5 model data are estimated with the ensemble spread: one standard deviation of the members' means is treated as the standard error for the ensemble mean. This applies to the dynamic/thermosteric SSH ocean data, and the IBE atmospheric data. The ice sheet, glacier and land water storage uncertainties are found regionally from the global uncertainties of the sources using the same iterative sea level equation solver used to obtain the regional distribution from their means. The one standard error of the GIA uncertainty is evaluated as the departures of the two different GIA estimates (from ICE-5G and ANU/SELEN models) from their mean value. To combine these uncertainties, for both maps of uncertainty as well as time series of uncertainty at individual stations, it is assumed that contributions that correlate with global air temperature have correlated uncertainties, which are therefore added linearly. This combined uncertainty is then added to the other



components' uncertainties in quadrature. The uncertainties in the projected ice sheet SMB changes were assumed to be dominated by the magnitude of climate change, rather than their methodological uncertainty, while the uncertainty in the projected glacier change was assumed to be dominated by its methodological uncertainty. The formula shown below for the regional error, when applied to the global contributions, estimates a global uncertainty close to that given in Table 13.5. The estimated squared uncertainty (standard error) at each grid box is found as follows:

$$\sigma_{\text{tot}}^2 = (\sigma_{\text{steric/dyn}} + \sigma_{\text{smb\_a}} + \sigma_{\text{smb\_g}})^2 + \sigma_{\text{glac}}^2 + \sigma_{\text{IBE}}^2 + \sigma_{\text{GIA}}^2 + \sigma_{\text{LW}}^2 + \sigma_{\text{dyn\_a}}^2 + \sigma_{\text{dyn\_g}}^2 \quad (13.SM.1)$$

where:

*steric/dyn* = global thermal expansion uncertainty + dynamic SSH (ensemble spread)

*smb\_a* = Antarctic ice sheet SMB uncertainty (including interaction of SMB and dynamics)

*smb\_g* = Greenland ice sheet SMB uncertainty (including interaction of SMB and dynamics)

*glac* = Glacier uncertainty

*IBE* = inverse barometer effect uncertainty (ensemble spread)

*GIA* = glacial isostatic adjustment uncertainty

*LW* = land water storage uncertainty

*dyn\_a* = Antarctica ice sheet rapid dynamics uncertainty

*dyn\_g* = Greenland ice sheet rapid dynamics uncertainty

The 90% confidence limits for the ice components are asymmetric and were combined with the 90% confidence limit uncertainties of the CMIP5 ocean components to find the lower and upper uncertainty limits separately (Figures 13.19 and 13.23), using the given equation. In Figure 13.21, in which a single standard error at each location is used, the  $\sigma$  used in the equation were standard deviations for all components except *LW*, *dyn\_a* and *dyn\_g*; these latter had uniform PDFs in the global projections, and the half-range of the distribution was used for  $\sigma$ . To find the 90% confidence limits of the ocean components, regional uncertainties were multiplied by 1.645, thus treating them as methodological, normally distributed uncertainties.

**Table 13.SM.3** | Availability of 'zos' variable from CMIP5.

Model	RCP2.6	RCP6.0	RCP4.5 / RCP8.5
ACCESS-1.0			X
BCC-CSM1.1	X	X	X
CanESM2			X
CNRM-CM5			X
CSIRO-MK3.6.0	X	X	X
GFDL-ESM2G	X	X	X
GFDL-ESM2M	X	X	X
GISS-E2-R	X	X	X
HadGEM2-CC			X
HadGEM2-ES	X		X
INM-CM4			X
IPSL-CM5A-LR	X	X	X
IPSL-CM5A-MR	X		X
MIROC5	X	X	X
MIROC-ESM	X	X	X
MIROC-ESM-CHEM	X	X	X
MPI-ESM-LR	X		X
MPI-ESM-MR	X		X
MRI-CGCM3	X	X	X
NorESM1-M	X	X	X
NorESM1-ME	X	X	X

## References

- Bengtsson, L., S. Koumoutsaris, and K. Hodges, 2011: Large-scale surface mass balance of ice sheets from a comprehensive atmosphere model. *Surv. Geophys.*, **32**, 459–474.
- Farrell, W.E., and Clark, J.A., 1976. On postglacial sea-level, *Geophys. J. R. Astr. Soc.*, **46**, 647–667.
- Fettweis, X., B. Franco, M. Tedesco, J. H. van Angelen, J. T. M. Lenaerts, M. R. van den Broeke, and H. Gallee, 2013: Estimating Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric model MAR. *Cryosphere*, **7**, 469–489.
- Giesen, R. H., and J. Oerlemans, 2013: Climate-model induced differences in the 21st century global and regional glacier contributions to sea-level rise. *Clim. Dyn.*, **41**, 3283–3300.
- Gleckler, P. J., et al., 2012: Human-induced global ocean warming on multidecadal timescales. *Nature Clim. Change*, **2**, 524–529.
- Good, P., J. M. Gregory, and J. A. Lowe, 2011: A step-response simple climate model to reconstruct and interpret AOGCM projections. *Geophys. Res. Lett.*, **38**, L01703.
- Good, P., J. M. Gregory, J. A. Lowe, and T. Andrews, 2013: Abrupt CO<sub>2</sub> experiments as tools for predicting and understanding CMIP5 representative concentration pathway projections. *Clim. Dyn.*, **40**, 1041–1053.
- Gregory, J. M., and P. Huybrechts, 2006: Ice-sheet contributions to future sea-level change. *Philos. Trans. R. Soc. London A*, **364**, 1709–1731.
- Kendall, R., Latychev, K., Mitrovica, J.X., Davis, J.E., and Tamisiea, M., 2006. Decontaminating tide gauge records for the influence of Glacial Isostatic Adjustment: the potential impact of 3-D Earth structure, *Geophys. Res. Lett.*, **33**, L24318, doi:10.1029/2006GL028448.
- Krinner, G., O. Magand, I. Simmonds, C. Genthon, and J. L. Dufresne, 2007: Simulated Antarctic precipitation and surface mass balance at the end of the twentieth and twenty-first centuries. *Clim. Dyn.*, **28**, 215–230.
- Kuhlbrodt, T., and J. M. Gregory, 2012: Ocean heat uptake and its consequences for the magnitude of sea level rise and climate change. *Geophys. Res. Lett.*, **39**, L18608.
- Lambeck, K., C. Smither, and P. Johnston, 1998: Sea-level change, glacial rebound and mantle viscosity for northern Europe. *Geophys. J. Int.*, **134**, 102–144.
- Ligtenberg, S. R. M., W. J. van de Berg, M. R. van den Broeke, J. G. L. Rae, and E. van Meijgaard, 2013: Future surface mass balance of the Antarctic ice sheet and its influence on sea level change, simulated by a regional atmospheric climate model. *Clim. Dyn.*, **41**, 867–884.
- Marzeion, B., A. H. Jarosch, and M. Hofer, 2012: Past and future sea-level changes from the surface mass balance of glaciers. *Cryosphere*, **6**, 1295–1322.
- Milne, G.A., and Mitrovica, J.X., 1998. Postglacial sea-level change on a rotating Earth, *Geophys. J. Int.*, **133**, 1–19.
- Peltier, W. R., 2004: Global glacial isostasy and the surface of the ice-age earth: The ICE-5G (VM2) model and GRACE. *Annu. Rev. Earth Planet. Sci.*, **32**, 111–149.
- Radić, V., A. Bliss, A. D. Beedlow, R. Hock, E. Miles, and J. G. Cogley, 2014: Regional and global projections of twenty-first century glacier mass changes in response to climate scenarios from global climate models. *Clim. Dyn.*, **42**, 37–58.
- Slangen, A. B. A., and R. S. W. van de Wal, 2011: An assessment of uncertainties in using volume-area modelling for computing the twenty-first century glacier contribution to sea-level change. *Cryosphere*, **5**, 673–686.
- Slangen, A. B. A., C. A. Katsman, R. S. W. van de Wal, L. L. A. Vermeersen, and R. E. M. Riva, 2012: Towards regional projections of twenty-first century sea-level change based on IPCC SRES scenarios. *Clim. Dyn.*, **38**, 1191–1209.
- Spada, G., and Stocchi, P., 2006. The Sea Level Equation, Theory and Numerical Examples, Aracne, Roma, p. 96, ISBN: 88-548-0384-7.
- Spada, G., and Stocchi, P., 2007. SELEN: a Fortran 90 program for solving the 'Sea Level Equation', *Comput. Geosci.*, **33**(4), 538–562, doi:10.1016/j.cageo.2006.08.006.
- Wada, Y., L. P. H. van Beek, F. C. S. Weiland, B. F. Chao, Y. H. Wu, and M. F. P. Bierkens, 2012: Past and future contribution of global groundwater depletion to sea-level rise. *Geophys. Res. Lett.*, **39**, L09402.
- Yoshimori, M., and A. Abe-Ouchi, 2012: Sources of spread in multi-model projections of the Greenland ice-sheet surface mass balance. *J. Clim.*, **25**, 1157–1175.